



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:17 pm GMT

PDB ID : 2C13
Title : 5-HYDROXY-LEVULINIC ACID BOUND TO PORPHOBILINOGEN SYNTHASE FROM PSEUDOMONAS AERUGINOSA
Authors : Frere, F.; Nentwich, M.; Gacond, S.; Heinz, D.W.; Neier, R.; Frankenberg-Dinkel, N.
Deposited on : 2005-09-11
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 16%, green 80%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 80% 16% .. </div> </div>
1	B	337	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 12%, green 83%, grey 3%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 83% 12% ... </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PGE	B	1338	-	-	-	X
6	PGE	B	1339	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6020 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DELTA-AMINOLEVULINIC ACID DEHYDRATASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	334	Total	C	N	O	S	0	21	1
			2777	1740	492	533	12			
1	B	331	Total	C	N	O	S	0	22	1
			2753	1723	482	536	12			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	VAL	ILE	ENGINEERED MUTATION	UNP Q59643
B	199	VAL	ILE	ENGINEERED MUTATION	UNP Q59643


- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	1	Total	Cl	0	0
			1	1		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	K	0	0
			1	1		
5	A	1	Total	K	0	0
			1	1		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			10	6	4		
6	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	228	Total	O	0	0
			228	228		
7	B	229	Total	O	0	0
			229	229		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



• Molecule 1: DELTA-AMINOLEVULINIC ACID DEHYDRATASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	126.31Å 126.31Å 85.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	89.80 – 2.15 39.94 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (89.80-2.15) 99.2 (39.94-2.15)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.14 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.152 , 0.204 0.153 , 0.204	Depositor DCC
R_{free} test set	1931 reflections (5.38%)	DCC
Wilson B-factor (Å ²)	19.5	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6020	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FHL, MG, PGE, CL, K, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/2791	0.87	9/3782 (0.2%)
1	B	0.83	0/2766	0.88	10/3749 (0.3%)
All	All	0.82	0/5557	0.88	19/7531 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	4
All	All	0	7

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	198	ARG	NE-CZ-NH2	-11.13	114.73	120.30
1	A	25[A]	ARG	NE-CZ-NH2	-8.98	115.81	120.30
1	A	25[B]	ARG	NE-CZ-NH2	-8.98	115.81	120.30
1	B	198	ARG	NE-CZ-NH1	7.57	124.08	120.30
1	A	97	ASP	CB-CG-OD1	6.80	124.42	118.30
1	B	113	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	25[A]	ARG	NE-CZ-NH1	6.15	123.38	120.30
1	A	25[B]	ARG	NE-CZ-NH1	6.15	123.38	120.30
1	B	138	GLN	N-CA-C	5.80	126.67	111.00
1	B	277	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	A	333	LEU	N-CA-C	5.39	125.56	111.00
1	B	277	ARG	NE-CZ-NH2	-5.37	117.61	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	256[A]	MET	CG-SD-CE	5.34	108.74	100.20
1	B	256[B]	MET	CG-SD-CE	5.34	108.74	100.20
1	A	26	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	A	157	LEU	CA-CB-CG	5.29	127.48	115.30
1	B	271	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	A	274	ASP	CB-CG-OD1	5.14	122.93	118.30
1	B	235	ASP	CB-CG-OD1	5.02	122.81	118.30

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	260	FHL	Mainchain,Peptide
1	A	333	LEU	Peptide
1	B	137	GLY	Peptide
1	B	205	FHL	Mainchain
1	B	260	FHL	Mainchain,Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2777	0	2717	43	0
1	B	2753	0	2669	48	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	7	0	10	2	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	B	20	0	28	1	0
7	A	228	0	0	16	0
7	B	229	0	0	16	0
All	All	6020	0	5424	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 8.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25[B]:ARG:NH1	7:A:2037:HOH:O	1.71	1.22
1:A:25[A]:ARG:HD2	7:A:2037:HOH:O	1.57	1.04
1:B:335:ARG:HG3	1:B:336:GLY:N	1.80	0.96
1:B:187[B]:GLU:OE2	7:B:2152:HOH:O	1.84	0.96
1:A:166[B]:GLU:OE2	7:A:2138:HOH:O	1.88	0.89
1:A:32[A]:VAL:CG2	7:A:2045:HOH:O	2.21	0.87
1:B:293[B]:MET:HE2	1:B:325:PHE:HE1	1.42	0.84
1:B:118[A]:ARG:NE	7:B:2112:HOH:O	2.11	0.83
1:B:118[A]:ARG:NH2	7:B:2112:HOH:O	2.13	0.80
1:A:330:ALA:O	1:A:334:ARG:HB2	1.83	0.78
1:B:293[B]:MET:CE	1:B:325:PHE:HE1	1.97	0.76
1:B:222[A]:SER:C	7:B:2173:HOH:O	2.23	0.75
7:A:2002:HOH:O	1:B:244[B]:HIS:HD2	1.72	0.73
1:B:198:ARG:N	1:B:198:ARG:HD3	2.04	0.73
1:B:71[A]:GLU:OE2	7:B:2078:HOH:O	2.06	0.72
1:B:331[B]:GLU:OE2	7:B:2224:HOH:O	2.10	0.70
1:B:222[A]:SER:O	7:B:2171:HOH:O	2.12	0.68
1:B:335:ARG:CG	1:B:336:GLY:N	2.53	0.67
1:A:32[B]:VAL:HG13	7:A:2045:HOH:O	1.94	0.67
1:A:166[A]:GLU:OE2	7:A:2139:HOH:O	2.12	0.67
1:B:194:HIS:HB3	1:B:197[B]:VAL:CG1	2.24	0.67
1:B:154[B]:ILE:HD11	1:B:178:MET:HB3	1.78	0.66
1:B:154[B]:ILE:HD11	1:B:178:MET:CB	2.26	0.65
1:B:222[B]:SER:HA	7:B:2173:HOH:O	1.97	0.65
1:A:32[A]:VAL:HG23	7:A:2045:HOH:O	1.91	0.63
1:B:118[A]:ARG:CZ	7:B:2112:HOH:O	2.40	0.62
1:B:213:PRO:HB2	1:B:286:SER:HB2	1.81	0.62
1:A:136[A]:HIS:HD2	1:A:138:GLN:H	1.48	0.62
1:B:2:SER:HB2	7:B:2001:HOH:O	1.98	0.61
1:B:293[B]:MET:HE2	1:B:325:PHE:CE1	2.30	0.60
1:A:301:LEU:HG	1:A:305:VAL:HG11	1.83	0.60
1:B:105[B]:GLU:OE1	7:B:2103:HOH:O	2.17	0.60
1:B:143:ASP:HB3	1:B:149:LEU:HD21	1.84	0.59
1:A:252:GLU:HA	4:A:1338:PEG:H42	1.84	0.58
1:A:327:LYS:HE2	1:A:331:GLU:OE2	2.03	0.58
1:A:273[B]:LYS:NZ	7:A:2196:HOH:O	2.37	0.57
1:B:301:LEU:HG	1:B:305:VAL:HG11	1.87	0.56
1:A:252:GLU:HA	4:A:1338:PEG:C4	2.37	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:38:LEU:O	1:B:81:PRO:HD2	2.06	0.55
1:B:293[B]:MET:CE	1:B:325:PHE:CE1	2.86	0.55
1:A:289:TYR:CZ	1:A:293[A]:MET:HG3	2.42	0.54
1:A:45:LEU:HD21	1:A:62[A]:ARG:HD3	1.89	0.54
1:B:130:LEU:HD12	1:B:140:GLY:HA2	1.88	0.54
1:B:134:THR:OG1	1:B:137:GLY:HA2	2.06	0.54
1:A:35[A]:VAL:HG11	7:A:2074:HOH:O	2.09	0.53
1:B:48:VAL:O	1:B:50:GLN:HG2	2.09	0.52
1:A:136[A]:HIS:CD2	1:A:138:GLN:HB2	2.44	0.52
1:A:143:ASP:HB3	1:A:149:LEU:HD21	1.90	0.52
1:B:132:PRO:HB3	7:B:2084:HOH:O	2.10	0.52
1:A:62[B]:ARG:NH2	7:A:2064:HOH:O	2.39	0.52
1:A:130:LEU:HD12	1:A:140:GLY:HA2	1.92	0.51
1:B:195[A]:THR:HG21	7:B:2157:HOH:O	2.11	0.51
1:A:105:GLU:O	1:A:110[A]:ARG:HD3	2.10	0.51
1:B:154[A]:ILE:HD11	1:B:179:ASP:O	2.11	0.50
1:A:332[B]:GLN:HG2	1:A:335:ARG:NH2	2.26	0.50
1:B:2:SER:CB	7:B:2001:HOH:O	2.58	0.50
1:B:222[A]:SER:HA	7:B:2173:HOH:O	2.11	0.50
1:A:136[A]:HIS:CD2	1:A:138:GLN:H	2.30	0.49
1:B:198:ARG:HA	1:B:255:ASP:OD2	2.12	0.49
1:A:38:LEU:O	1:A:81:PRO:HD2	2.12	0.49
1:A:45:LEU:HG	1:A:62[A]:ARG:HB3	1.95	0.48
1:A:55:PRO:HB3	6:B:1338:PGE:H52	1.96	0.48
1:B:154[B]:ILE:HD11	1:B:178:MET:HB2	1.95	0.47
1:A:198:ARG:HA	1:A:255:ASP:OD2	2.15	0.47
1:A:273[A]:LYS:NZ	7:A:2196:HOH:O	2.36	0.47
1:A:131:ASP:HB3	1:A:139:ASP:OD1	2.16	0.46
1:A:136[A]:HIS:HE1	7:A:2084:HOH:O	1.98	0.46
1:A:90:PRO:HD2	1:A:93:LYS:HD2	1.98	0.46
1:B:214:PHE:CG	1:B:260:FHL:H2C1	2.52	0.45
1:A:42[B]:VAL:HG21	1:A:68:LEU:HD13	1.99	0.45
1:B:260:FHL:H3C1	1:B:260:FHL:HEC2	1.81	0.45
1:A:97:ASP:HB3	7:A:2088:HOH:O	2.16	0.44
1:B:154[B]:ILE:CD1	1:B:178:MET:CB	2.93	0.44
1:B:256[A]:MET:HB2	1:B:281:PHE:CD1	2.53	0.43
1:A:42[B]:VAL:HG21	1:A:68:LEU:CD1	2.48	0.43
1:B:198:ARG:N	1:B:198:ARG:CD	2.79	0.42
1:A:205:FHL:H3C2	1:A:260:FHL:NZ	2.35	0.42
1:A:154:ILE:HD11	1:A:179:ASP:O	2.19	0.41
1:B:154[B]:ILE:HD12	1:B:181:ARG:CB	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:ILE:HG21	1:A:303[B]:GLU:HG2	2.01	0.41
1:A:248:ALA:O	1:A:252:GLU:HG3	2.21	0.41
1:A:92[A]:GLU:HG3	7:A:2081:HOH:O	2.20	0.41
1:A:270:ARG:HA	1:A:316:ALA:O	2.19	0.41
1:A:287:GLY:HA3	1:B:301:LEU:HD11	2.03	0.41
1:A:32[A]:VAL:HG21	7:A:2045:HOH:O	2.05	0.41
1:B:222[A]:SER:CA	7:B:2173:HOH:O	2.62	0.41
1:B:154[B]:ILE:CD1	1:B:178:MET:HB2	2.51	0.41
1:B:51:ARG:HA	1:B:62:ARG:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/337 (104%)	340 (97%)	9 (3%)	2 (1%)	28	20
1	B	346/337 (103%)	331 (96%)	11 (3%)	4 (1%)	15	8
All	All	697/674 (103%)	671 (96%)	20 (3%)	6 (1%)	20	12

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	261	PRO
1	B	137	GLY
1	B	261	PRO
1	A	334	ARG
1	B	138	GLN
1	B	335	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/267 (107%)	272 (95%)	13 (5%)	31	27
1	B	284/267 (106%)	270 (95%)	14 (5%)	29	24
All	All	569/534 (107%)	542 (95%)	27 (5%)	33	26

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	PHE
1	A	49	ASN
1	A	53	SER
1	A	96	LEU
1	A	117	GLU
1	A	118	ARG
1	A	157	LEU
1	A	181	ARG
1	A	198	ARG
1	A	226	LYS
1	A	283	TYR
1	A	301	LEU
1	A	334	ARG
1	B	48	VAL
1	B	74[A]	GLU
1	B	74[B]	GLU
1	B	181	ARG
1	B	198	ARG
1	B	222[A]	SER
1	B	222[B]	SER
1	B	256[A]	MET
1	B	256[B]	MET
1	B	283	TYR
1	B	293[A]	MET
1	B	293[B]	MET
1	B	301	LEU
1	B	307	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	FHL	A	205	1	12,16,17	1.04	1 (8%)	10,18,20	1.02	0
1	FHL	A	260	1	12,16,17	0.80	0	10,18,20	3.39	5 (50%)
1	FHL	B	205	1	12,16,17	0.80	0	10,18,20	1.52	1 (10%)
1	FHL	B	260	1	12,16,17	0.97	1 (8%)	10,18,20	2.60	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	FHL	A	205	1	-	0/12/17/19	0/0/0/0
1	FHL	A	260	1	-	0/12/17/19	0/0/0/0
1	FHL	B	205	1	-	0/12/17/19	0/0/0/0
1	FHL	B	260	1	-	0/12/17/19	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	260	FHL	CA-C	2.32	1.53	1.50
1	A	205	FHL	CA-C	3.14	1.54	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	FHL	O-C-CA	-4.54	112.50	125.02
1	B	260	FHL	O-C-CA	-3.92	114.19	125.02
1	A	260	FHL	CD-CE-NZ	-3.13	105.80	110.71
1	A	260	FHL	C2-C3-C4	-2.31	109.99	114.58
1	A	260	FHL	CE-NZ-C4	2.18	128.72	122.49
1	B	205	FHL	O3-C5-C4	3.35	116.05	111.70
1	B	260	FHL	O3-C5-C4	6.59	120.25	111.70
1	A	260	FHL	O3-C5-C4	8.29	122.46	111.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	205	FHL	1	0
1	A	260	FHL	1	0
1	B	260	FHL	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 6 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	1338	-	6,6,6	0.46	0	5,5,5	0.80	0
6	PGE	B	1338	-	9,9,9	0.64	0	8,8,8	0.45	0
6	PGE	B	1339	-	9,9,9	0.68	0	8,8,8	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	1338	-	-	0/4/4/4	0/0/0/0
6	PGE	B	1338	-	-	0/7/7/7	0/0/0/0
6	PGE	B	1339	-	-	0/7/7/7	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1338	PEG	2	0
6	B	1338	PGE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/337 (98%)	-0.63	2 (0%) 89 91	11, 16, 27, 58	0
1	B	329/337 (97%)	-0.65	3 (0%) 84 87	9, 15, 29, 50	3 (0%)
All	All	661/674 (98%)	-0.64	5 (0%) 86 89	9, 15, 28, 58	3 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	335	ARG	3.2
1	A	334	ARG	3.1
1	B	144	ASP	2.5
1	B	334	ARG	2.1
1	B	335	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	FHL	A	205	17/18	0.96	0.11	-	10,13,19,22	0
1	FHL	A	260	17/18	0.95	0.12	-	2,12,14,15	0
1	FHL	B	260	17/18	0.95	0.10	-	5,12,14,16	0
1	FHL	B	205	17/18	0.95	0.14	-	15,20,29,33	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	PGE	B	1338	10/10	0.79	0.35	8.45	23,30,33,34	10
6	PGE	B	1339	10/10	0.77	0.24	6.97	36,43,47,47	0
3	CL	B	1337	1/1	1.00	0.08	0.25	23,23,23,23	0
4	PEG	A	1338	7/7	0.92	0.11	-0.19	29,31,35,36	0
5	K	B	1340	1/1	0.99	0.04	-1.90	22,22,22,22	0
2	MG	A	1336	1/1	0.99	0.06	-2.87	10,10,10,10	0
5	K	A	1339	1/1	0.99	0.02	-3.20	20,20,20,20	0
2	MG	B	1336	1/1	0.99	0.03	-3.90	10,10,10,10	0
3	CL	A	1337	1/1	0.99	0.08	-	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.