



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 10:58 AM EST

PDB ID : 6C12  
Title : SDHA-SDHE complex  
Authors : Maher, M.J.  
Deposited on : 2018-01-03  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030736  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	588	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 86%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>1%</span> <span>86%</span> <span>5%</span> <span>9%</span> </div> </div>
1	B	588	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 86%, yellow 6%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>86%</span> <span>6%</span> <span>9%</span> </div> </div>
2	C	101	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 77%, yellow 7%, grey 16%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>77%</span> <span>7%</span> <span>16%</span> </div> </div>
2	D	101	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 78%, yellow 7%, grey 15%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>78%</span> <span>7%</span> <span>15%</span> </div> </div>

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 10122 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase flavoprotein subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	537	Total	C	N	O	S	0	0	0
			4123	2566	743	786	28			
1	A	536	Total	C	N	O	S	0	0	0
			4100	2552	738	782	28			

- Molecule 2 is a protein called FAD assembly factor SdhE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	86	Total	C	N	O	S	0	0	0
			714	446	130	131	7			
2	C	85	Total	C	N	O	S	0	1	0
			720	450	132	130	8			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-12	MET	-	expression tag	UNP P64561
D	-11	GLY	-	expression tag	UNP P64561
D	-10	SER	-	expression tag	UNP P64561
D	-9	SER	-	expression tag	UNP P64561
D	-8	HIS	-	expression tag	UNP P64561
D	-7	HIS	-	expression tag	UNP P64561
D	-6	HIS	-	expression tag	UNP P64561
D	-5	HIS	-	expression tag	UNP P64561
D	-4	HIS	-	expression tag	UNP P64561
D	-3	HIS	-	expression tag	UNP P64561
D	-2	SER	-	expression tag	UNP P64561
D	-1	GLN	-	expression tag	UNP P64561
D	0	ASP	-	expression tag	UNP P64561
D	1	PRO	-	expression tag	UNP P64561
C	-12	MET	-	expression tag	UNP P64561
C	-11	GLY	-	expression tag	UNP P64561

*Continued on next page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	-10	SER	-	expression tag	UNP P64561
C	-9	SER	-	expression tag	UNP P64561
C	-8	HIS	-	expression tag	UNP P64561
C	-7	HIS	-	expression tag	UNP P64561
C	-6	HIS	-	expression tag	UNP P64561
C	-5	HIS	-	expression tag	UNP P64561
C	-4	HIS	-	expression tag	UNP P64561
C	-3	HIS	-	expression tag	UNP P64561
C	-2	SER	-	expression tag	UNP P64561
C	-1	GLN	-	expression tag	UNP P64561
C	0	ASP	-	expression tag	UNP P64561
C	1	PRO	-	expression tag	UNP P64561

- # FAD

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Na 1	0	0
4	A	1	Total 1	Na 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	182	Total 182	O 182	0	0
5	D	15	Total 15	O 15	0	0
5	A	140	Total 140	O 140	0	0
5	C	20	Total 20	O 20	0	0



MET	GLY	SER	SER	HIS	HIS	HIS	HIS	HIS	HIS	SER	GLN	ASP	PRO	D2	H10	R14	F27	H30	M59	R81	E82	R83	V86	ALA	ILE
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.98Å 100.03Å 232.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.89 – 2.15 48.89 – 2.15	Depositor EDS
% Data completeness (in resolution range)	74.0 (48.89-2.15) 74.0 (48.89-2.15)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.44 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.199 , 0.258 0.204 , 0.258	Depositor DCC
$R_{free}$ test set	2561 reflections (5.35%)	DCC
Wilson B-factor (Å <sup>2</sup> )	20.0	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 34.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	10122	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/4177	0.73	3/5651 (0.1%)
1	B	0.48	0/4202	0.77	6/5685 (0.1%)
2	C	0.48	0/738	0.86	3/992 (0.3%)
2	D	0.49	0/729	0.74	1/981 (0.1%)
All	All	0.48	0/9846	0.76	13/13309 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	460	ARG	NE-CZ-NH1	8.38	124.49	120.30
1	B	510	ARG	NE-CZ-NH1	7.74	124.17	120.30
1	B	510	ARG	NE-CZ-NH2	-7.43	116.59	120.30
1	A	510	ARG	NE-CZ-NH2	-6.25	117.17	120.30
1	A	510	ARG	NE-CZ-NH1	6.13	123.37	120.30
1	B	460	ARG	NE-CZ-NH2	-6.10	117.25	120.30
2	C	14	ARG	NE-CZ-NH1	5.88	123.24	120.30
1	A	183	LEU	CA-CB-CG	5.83	128.71	115.30
1	B	443	ARG	NE-CZ-NH1	5.82	123.21	120.30
1	B	533	ARG	NE-CZ-NH1	5.68	123.14	120.30
2	D	15	ARG	NE-CZ-NH2	-5.58	117.51	120.30
2	C	83	ARG	NE-CZ-NH1	5.39	123.00	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	81	ARG	NE-CZ-NH1	5.22	122.91	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	392	VAL	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4100	0	4016	15	0
1	B	4123	0	4032	19	0
2	C	720	0	702	3	0
2	D	714	0	687	4	0
3	A	53	0	30	0	0
3	B	53	0	30	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	140	0	0	0	0
5	B	182	0	0	0	0
5	C	20	0	0	0	0
5	D	15	0	0	0	0
All	All	10122	0	9497	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:207:ARG:HD2	1:B:216:HIS:HA	1.68	0.75
1:A:555:CYS:HA	1:A:571:VAL:HG23	1.83	0.60
2:D:80:ASN:OD1	2:D:83:ARG:NH2	2.31	0.59

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:27:PHE:O	2:C:30:HIS:O	2.21	0.59
1:B:139:ASP:HB3	1:B:141:THR:HG23	1.86	0.57
1:B:11:VAL:HG23	1:B:195:ALA:HB2	1.87	0.55
1:B:490:ILE:HG22	1:B:520:MET:CE	2.36	0.55
1:B:296:GLU:OE1	1:B:301:ARG:NE	2.35	0.53
1:A:26:ILE:HG12	1:A:419:LEU:HD22	1.91	0.51
1:A:298:ARG:NH2	1:A:470:ASN:OD1	2.43	0.51
1:B:6:ARG:HD2	1:B:191:VAL:HG11	1.93	0.50
1:B:336:THR:HG21	2:D:59:MET:HE1	1.94	0.50
1:A:20:MET:HG2	1:A:35:LEU:HD21	1.95	0.49
1:B:490:ILE:HG22	1:B:520:MET:HE1	1.95	0.49
1:B:2:LYS:HD3	1:A:275:ARG:HB3	1.95	0.48
1:B:139:ASP:CB	1:B:141:THR:HG23	2.46	0.45
1:B:454:GLU:OE2	1:B:493:ARG:NE	2.44	0.45
1:A:508:THR:HG21	2:C:10:HIS:CE1	2.51	0.45
1:A:340:VAL:HG13	1:A:345:GLU:HB2	1.97	0.45
1:A:537:ARG:NH1	1:A:548:ARG:HD2	2.32	0.45
1:B:83:ASP:OD1	1:B:84:ALA:N	2.49	0.45
1:B:310:HIS:CD2	1:B:348:PRO:HB3	2.52	0.44
1:A:255:GLU:OE1	1:A:286:ARG:NH2	2.50	0.44
1:A:472:SER:OG	1:A:473:VAL:N	2.49	0.44
2:D:25:MET:HB3	2:D:26:PRO:HD3	2.00	0.44
1:B:48:SER:HB3	3:B:601:FAD:HM72	1.99	0.43
1:B:79:ILE:HA	1:B:571:VAL:HG21	2.00	0.43
1:A:257:CYS:HB3	1:A:315:LEU:HD21	2.00	0.42
1:B:239:TRP:CE3	1:B:353:CYS:SG	3.13	0.42
1:B:490:ILE:HG22	1:B:520:MET:HE3	2.01	0.42
1:A:50:GLN:HE22	1:A:138:ALA:HA	1.84	0.42
1:A:336:THR:HG21	2:C:59:MET:HE2	2.02	0.42
1:A:207:ARG:HD2	1:A:216:HIS:HA	2.01	0.42
1:B:337:PHE:CZ	2:D:59:MET:HE1	2.55	0.42
1:A:227:ILE:HG23	1:A:561:PRO:HB3	2.02	0.41
1:B:26:ILE:HG12	1:B:419:LEU:HD22	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/588 (90%)	518 (98%)	11 (2%)	1 (0%)	51	50
1	B	531/588 (90%)	518 (98%)	12 (2%)	1 (0%)	51	50
2	C	84/101 (83%)	83 (99%)	1 (1%)	0	100	100
2	D	84/101 (83%)	81 (96%)	3 (4%)	0	100	100
All	All	1229/1378 (89%)	1200 (98%)	27 (2%)	2 (0%)	51	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	393	SER
1	A	393	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	432/473 (91%)	427 (99%)	5 (1%)	75	80
1	B	434/473 (92%)	432 (100%)	2 (0%)	91	94
2	C	79/92 (86%)	79 (100%)	0	100	100
2	D	77/92 (84%)	76 (99%)	1 (1%)	73	79
All	All	1022/1130 (90%)	1014 (99%)	8 (1%)	85	89

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	393	SER
1	B	404	SER
2	D	71	MET
1	A	183	LEU
1	A	275	ARG
1	A	333	LEU
1	A	393	SER
1	A	429	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	56	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	FAD	A	601	1	51,58,58	1.51	8 (15%)	54,89,89	2.17	11 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FAD	B	601	1	51,58,58	1.55	8 (15%)	54,89,89	2.17	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	A	601	1	-	0/28/50/50	0/6/6/6
3	FAD	B	601	1	-	0/28/50/50	0/6/6/6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	FAD	C2A-N3A	2.05	1.35	1.32
3	A	601	FAD	C4X-N5	2.13	1.36	1.33
3	B	601	FAD	C10-N1	2.39	1.36	1.33
3	A	601	FAD	C10-N1	2.58	1.36	1.33
3	B	601	FAD	C5A-C4A	2.95	1.47	1.40
3	A	601	FAD	C5A-C4A	2.99	1.47	1.40
3	A	601	FAD	C9A-N10	3.25	1.43	1.38
3	B	601	FAD	C9A-N10	3.62	1.43	1.38
3	A	601	FAD	C8-C7	3.66	1.50	1.41
3	A	601	FAD	C9A-C5X	3.76	1.50	1.42
3	B	601	FAD	C8-C7	3.76	1.50	1.41
3	A	601	FAD	C4-C4X	3.97	1.48	1.41
3	B	601	FAD	C9A-C5X	4.06	1.50	1.42
3	B	601	FAD	C4-C4X	4.07	1.49	1.41
3	A	601	FAD	C4X-C10	4.29	1.48	1.41
3	B	601	FAD	C4X-C10	4.46	1.48	1.41

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	FAD	N3A-C2A-N1A	-7.42	122.40	128.86
3	A	601	FAD	N3A-C2A-N1A	-7.15	122.63	128.86
3	B	601	FAD	C4-C4X-C10	-4.52	116.30	119.96
3	A	601	FAD	C4-C4X-C10	-4.36	116.44	119.96
3	A	601	FAD	C4X-C4-N3	-3.03	119.16	123.48
3	B	601	FAD	C4X-C4-N3	-2.95	119.29	123.48
3	A	601	FAD	C1B-N9A-C4A	-2.82	121.76	126.64
3	A	601	FAD	C4A-C5A-N7A	-2.76	106.74	109.41

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	FAD	C4X-C10-N10	-2.29	118.93	120.52
3	B	601	FAD	C4A-C5A-N7A	-2.27	107.22	109.41
3	A	601	FAD	C2A-N1A-C6A	2.17	122.56	118.77
3	B	601	FAD	C4-C4X-N5	3.51	122.53	118.68
3	A	601	FAD	C4-C4X-N5	3.58	122.61	118.68
3	B	601	FAD	C4X-N5-C5X	4.42	121.42	116.76
3	A	601	FAD	C4X-N5-C5X	4.58	121.60	116.76
3	A	601	FAD	C1'-N10-C9A	5.45	123.34	118.35
3	B	601	FAD	C1'-N10-C9A	5.45	123.34	118.35
3	A	601	FAD	C4-N3-C2	7.95	122.11	115.16
3	B	601	FAD	C4-N3-C2	8.01	122.17	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	601	FAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	536/588 (91%)	-0.22	7 (1%) 77 82	13, 25, 47, 82	0
1	B	537/588 (91%)	-0.36	2 (0%) 92 93	13, 22, 37, 48	0
2	C	85/101 (84%)	-0.26	0 100 100	17, 26, 38, 48	0
2	D	86/101 (85%)	-0.17	0 100 100	18, 25, 37, 39	0
All	All	1244/1378 (90%)	-0.28	9 (0%) 87 90	13, 24, 42, 82	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	PHE	5.2
1	A	577	LEU	4.6
1	A	428	ALA	3.2
1	A	51	GLY	2.8
1	A	576	LYS	2.3
1	B	67	TRP	2.2
1	A	580	ALA	2.2
1	A	326	ARG	2.1
1	B	157	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	FAD	B	601	53/53	0.97	0.10	-0.12	13,16,18,19	0
4	NA	B	602	1/1	0.97	0.08	-0.27	14,14,14,14	0
3	FAD	A	601	53/53	0.97	0.09	-0.44	16,18,24,26	0
4	NA	A	602	1/1	0.98	0.04	-2.55	16,16,16,16	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.