



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2017 – 05:21 PM EDT

PDB ID : 5C2Z  
Title : Molecular insights into the specificity of exfoliative toxins from *Staphylococcus aureus*  
Authors : Mariutti, R.B.; Souza, T.A.C.B.; Ullah, A.; Zaphorlin, L.M.; Murakami, M.T.; Arni, R.K.  
Deposited on : unknown  
Resolution : 1.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	<div> <div>2%</div> <div>88%</div> <div>9%</div> <div>.</div> </div>
1	B	255	<div> <div>%</div> <div>79%</div> <div>18%</div> <div>.</div> </div>

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4213 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exfoliative toxin D2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	22	2	0
			1928	1228	320	379	1			
1	B	248	Total	C	N	O		13	0	0
			1921	1219	321	381				

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	25	HIS	-	expression tag	UNP L0RUV7
A	26	HIS	-	expression tag	UNP L0RUV7
A	27	HIS	-	expression tag	UNP L0RUV7
A	28	HIS	-	expression tag	UNP L0RUV7
A	29	HIS	-	expression tag	UNP L0RUV7
A	30	HIS	-	expression tag	UNP L0RUV7
A	31	MET	-	expression tag	UNP L0RUV7
B	25	HIS	-	expression tag	UNP L0RUV7
B	26	HIS	-	expression tag	UNP L0RUV7
B	27	HIS	-	expression tag	UNP L0RUV7
B	28	HIS	-	expression tag	UNP L0RUV7
B	29	HIS	-	expression tag	UNP L0RUV7
B	30	HIS	-	expression tag	UNP L0RUV7
B	31	MET	-	expression tag	UNP L0RUV7

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	180	Total	O	0	0
			180	180		
2	B	184	Total	O	0	0
			184	184		



- Molecule 1: Exfoliative toxin D2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.41Å 93.14Å 50.48Å 90.00° 91.23° 90.00°	Depositor
Resolution (Å)	35.69 – 1.96 35.69 – 1.96	Depositor EDS
% Data completeness (in resolution range)	96.2 (35.69-1.96) 96.2 (35.69-1.96)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.39 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.177 , 0.253 0.181 , 0.256	Depositor DCC
$R_{free}$ test set	1614 reflections (5.08%)	DCC
Wilson B-factor (Å <sup>2</sup> )	25.8	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 38.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.065 for l,k,-h 0.064 for h,-k,-l 0.049 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4213	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.98% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	3/1973 (0.2%)	0.58	2/2668 (0.1%)
1	B	0.49	0/1960	0.58	0/2649
All	All	0.54	3/3933 (0.1%)	0.58	2/5317 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	131[A]	ILE	CA-C	5.64	1.67	1.52
1	A	131[B]	ILE	CA-C	5.64	1.67	1.52
1	A	66	PRO	N-CD	5.15	1.55	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	131[A]	ILE	CA-C-O	5.14	130.89	120.10
1	A	131[B]	ILE	CA-C-O	5.14	130.89	120.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1928	0	1928	23	0
1	B	1921	0	1907	40	0
2	A	180	0	0	2	1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	184	0	0	3	1
All	All	4213	0	3835	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:PRO:CA	1:B:51:SER:HB2	1.95	0.96
1:B:50:PRO:CB	1:B:51:SER:HB2	2.01	0.90
1:A:49:ARG:HA	1:A:213:TYR:CE1	2.16	0.81
1:B:50:PRO:HA	1:B:51:SER:HB2	1.62	0.79
1:B:49:ARG:HG3	1:B:213:TYR:OH	1.83	0.78
1:B:50:PRO:HB3	1:B:51:SER:HB2	1.66	0.77
1:B:49:ARG:HG3	1:B:213:TYR:CZ	2.22	0.74
1:A:75:VAL:HG22	1:A:108:VAL:HG12	1.73	0.68
1:A:49:ARG:CA	1:A:213:TYR:CE1	2.79	0.66
1:B:173:VAL:HB	1:B:250:ASN:HB3	1.78	0.64
1:B:74:PHE:CZ	1:B:76:LYS:HA	2.33	0.63
1:A:50:PRO:HD3	1:A:213:TYR:CG	2.36	0.60
1:B:203:PHE:CD1	1:B:211:PHE:CE2	2.90	0.59
1:B:203:PHE:CD1	1:B:211:PHE:CZ	2.93	0.57
1:A:49:ARG:HG3	1:A:213:TYR:CE2	2.41	0.56
1:B:171:ASP:OD2	2:B:301:HOH:O	2.18	0.55
1:B:200:ILE:HA	1:B:212:ALA:HB2	1.89	0.55
1:B:215:GLU:HB3	1:B:216:PRO:HD2	1.88	0.55
1:B:125:LYS:NZ	1:B:129:GLU:OE2	2.31	0.55
1:A:60:GLN:H	1:A:60:GLN:CD	2.10	0.55
1:B:203:PHE:HB2	1:B:211:PHE:CE2	2.42	0.55
1:B:279:GLN:O	2:B:302:HOH:O	2.18	0.55
1:A:50:PRO:HD3	1:A:213:TYR:CD1	2.45	0.52
1:B:170:PRO:HD3	1:B:229:GLU:HB3	1.92	0.52
1:A:170:PRO:HD3	1:A:229:GLU:HB3	1.93	0.51
1:B:184:LEU:HD12	1:B:224:PHE:CD1	2.45	0.50
1:A:123[A]:VAL:HG13	1:A:125:LYS:HE3	1.94	0.49
1:A:49:ARG:HA	1:A:213:TYR:CD1	2.46	0.49
1:B:88:LYS:HA	1:B:161:GLY:HA3	1.94	0.49
1:A:32:ILE:HD13	1:A:178:GLY:HA3	1.95	0.49
1:B:95:LYS:NZ	1:B:142:GLY:O	2.46	0.49
1:B:45:PHE:O	1:B:48:THR:HG22	2.13	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:ARG:HA	1:A:213:TYR:CZ	2.48	0.48
1:B:52:ASP:HB3	1:B:55:LEU:HG	1.95	0.48
1:A:49:ARG:HG3	1:A:213:TYR:CZ	2.49	0.47
1:B:175:VAL:HG22	1:B:248:LEU:HD11	1.96	0.47
1:B:158:LYS:HD2	1:B:163:LEU:HD11	1.97	0.47
1:A:49:ARG:N	1:A:213:TYR:CE1	2.83	0.46
1:B:211:PHE:CE1	1:B:243:LEU:HD11	2.51	0.46
1:B:180:LYS:HA	1:B:200:ILE:O	2.16	0.46
1:A:96:HIS:NE2	1:B:279:GLN:HG2	2.31	0.45
1:A:49:ARG:HB2	1:A:213:TYR:OH	2.16	0.45
1:B:50:PRO:CA	1:B:51:SER:CB	2.82	0.45
1:A:74:PHE:O	1:A:108:VAL:HA	2.18	0.44
1:B:215:GLU:O	1:B:218:ASN:HB2	2.18	0.44
1:B:264:THR:OG1	1:B:265:THR:N	2.50	0.43
1:A:60:GLN:HG2	1:A:61:ASP:H	1.83	0.42
1:B:203:PHE:HD1	1:B:211:PHE:HE2	1.67	0.42
1:B:203:PHE:CD1	1:B:211:PHE:HE2	2.37	0.42
1:B:50:PRO:HA	1:B:51:SER:CB	2.43	0.42
1:B:78:LYS:HG3	1:B:100:LEU:HB3	2.02	0.41
1:B:225:ASN:HB2	2:B:418:HOH:O	2.20	0.41
1:B:49:ARG:HD3	1:B:190:THR:OG1	2.20	0.41
1:B:156:TYR:HB2	1:B:158:LYS:HE2	2.02	0.41
1:A:150:LYS:NZ	2:A:318:HOH:O	2.53	0.41
1:A:99:ARG:NH2	2:A:317:HOH:O	2.53	0.41
1:A:49:ARG:CB	1:A:213:TYR:CZ	3.04	0.41
1:B:95:LYS:HA	1:B:147:SER:OG	2.20	0.41
1:B:207:THR:O	1:B:209:GLN:HG3	2.22	0.40
1:A:236:GLY:O	1:A:244:PRO:HA	2.21	0.40
1:A:76:LYS:HB3	1:A:76:LYS:HE2	1.93	0.40
1:B:50:PRO:CB	1:B:51:SER:CB	2.88	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:470:HOH:O	2:B:466:HOH:O[1_554]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/255 (97%)	235 (95%)	13 (5%)	0	100	100
1	B	246/255 (96%)	234 (95%)	11 (4%)	1 (0%)	38	25
All	All	494/510 (97%)	469 (95%)	24 (5%)	1 (0%)	51	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	51	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/219 (98%)	213 (100%)	1 (0%)	91	90
1	B	212/219 (97%)	211 (100%)	1 (0%)	91	90
All	All	426/438 (97%)	424 (100%)	2 (0%)	91	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	64	ARG
1	B	51	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	248/255 (97%)	-0.04	4 (1%)	72 80	16, 28, 50, 68	9 (3%)
1	B	248/255 (97%)	-0.13	2 (0%)	86 91	15, 25, 44, 63	6 (2%)
All	All	496/510 (97%)	-0.08	6 (1%)	79 85	15, 26, 46, 68	15 (3%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	211	PHE	4.6
1	B	108	VAL	2.7
1	A	213	TYR	2.6
1	A	46	PHE	2.5
1	A	135	ILE	2.4
1	A	261	LYS	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.