



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 11, 2018 – 10:39 PM EDT

PDB ID : 6C2G
Title : Human triosephosphate isomerase mutant V231M
Authors : Torres, L.A.; Enriquez, F.S.
Deposited on : 2018-01-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 98%; height: 10px; background-color: green;"></div> <div style="width: 98%; text-align: right;">98%</div> </div>
1	B	249	<div> <div style="width: 4%; height: 10px; background-color: red;"></div> <div style="width: 98%; height: 10px; background-color: green;"></div> <div style="width: 98%; text-align: right;">98%</div> </div>
1	C	249	<div> <div style="width: 3%; height: 10px; background-color: red;"></div> <div style="width: 98%; height: 10px; background-color: green;"></div> <div style="width: 98%; text-align: right;">98%</div> </div>
1	D	249	<div> <div style="width: 16%; height: 10px; background-color: red;"></div> <div style="width: 97%; height: 10px; background-color: green;"></div> <div style="width: 97%; text-align: right;">97%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7536 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Triosephosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	244	Total	C	N	O	S	0	0	0
			1841	1165	319	349	8			
1	B	244	Total	C	N	O	S	0	0	0
			1841	1165	319	349	8			
1	C	243	Total	C	N	O	S	0	0	0
			1834	1161	318	347	8			
1	D	241	Total	C	N	O	S	0	0	0
			1822	1154	315	345	8			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	231	MET	VAL	engineered mutation	UNP P60174
B	231	MET	VAL	engineered mutation	UNP P60174
C	231	MET	VAL	engineered mutation	UNP P60174
D	231	MET	VAL	engineered mutation	UNP P60174


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	67	Total	O	0	0
			67	67		
2	B	58	Total	O	0	0
			58	58		
2	C	52	Total	O	0	0
			52	52		
2	D	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Triosephosphate isomerase




- Molecule 1: Triosephosphate isomerase



- Molecule 1: Triosephosphate isomerase



- Molecule 1: Triosephosphate isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.07Å 63.45Å 334.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.89 – 2.30 41.89 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (41.89-2.30) 99.9 (41.89-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.84 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.215 , 0.226 0.222 , 0.232	Depositor DCC
R_{free} test set	2419 reflections (5.45%)	DCC
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.17$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7536	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/1875	0.65	0/2538
1	B	0.44	0/1875	0.65	0/2538
1	C	0.49	0/1867	0.67	0/2525
1	D	0.43	0/1853	0.67	0/2503
All	All	0.46	0/7470	0.66	0/10104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1841	0	1851	0	0
1	B	1841	0	1851	0	0
1	C	1834	0	1843	0	0
1	D	1822	0	1830	0	0
2	A	67	0	0	0	0
2	B	58	0	0	0	0
2	C	52	0	0	0	0
2	D	21	0	0	0	0
All	All	7536	0	7375	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/249 (97%)	235 (97%)	7 (3%)	0	100	100
1	B	242/249 (97%)	234 (97%)	8 (3%)	0	100	100
1	C	239/249 (96%)	231 (97%)	8 (3%)	0	100	100
1	D	233/249 (94%)	225 (97%)	8 (3%)	0	100	100
All	All	956/996 (96%)	925 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	192/196 (98%)	192 (100%)	0	100	100
1	B	192/196 (98%)	192 (100%)	0	100	100
1	C	191/196 (97%)	191 (100%)	0	100	100
1	D	191/196 (97%)	191 (100%)	0	100	100
All	All	766/784 (98%)	766 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/249 (97%)	-0.09	3 (1%) 79 82	23, 35, 58, 70	0
1	B	244/249 (97%)	0.21	10 (4%) 38 45	24, 38, 74, 93	0
1	C	243/249 (97%)	0.15	8 (3%) 47 54	28, 43, 66, 83	0
1	D	241/249 (96%)	0.81	40 (16%) 2 3	31, 72, 115, 128	0
All	All	972/996 (97%)	0.27	61 (6%) 21 27	23, 43, 97, 128	0

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	176	ALA	4.9
1	D	33	VAL	4.9
1	C	215	ALA	4.8
1	D	170	ILE	4.4
1	D	140	GLU	4.4
1	C	175	THR	4.3
1	C	170	ILE	4.0
1	D	28	LEU	3.6
1	D	36	ASP	3.5
1	D	220	LEU	3.4
1	D	212	VAL	3.3
1	B	136	ALA	3.2
1	D	243	ILE	3.1
1	D	181	ALA	3.1
1	D	227	ASP	3.1
1	C	173	GLY	3.0
1	D	30	ALA	3.0
1	B	175	THR	2.9
1	D	172	THR	2.9
1	D	177	THR	2.9
1	B	174	LYS	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	35	ALA	2.8
1	D	238	PRO	2.8
1	D	242	ASP	2.8
1	B	137	GLY	2.8
1	C	168	TRP	2.8
1	B	183	GLU	2.7
1	B	171	GLY	2.7
1	D	199	ALA	2.7
1	D	246	ALA	2.7
1	D	8	VAL	2.7
1	D	188	LEU	2.6
1	D	186	GLU	2.6
1	D	175	THR	2.6
1	D	216	THR	2.6
1	D	225	ASP	2.6
1	D	222	SER	2.5
1	C	169	ALA	2.5
1	D	226	VAL	2.5
1	D	174	LYS	2.5
1	D	151	ALA	2.4
1	D	168	TRP	2.4
1	A	173	GLY	2.4
1	D	202	GLN	2.3
1	B	131	LEU	2.3
1	A	175	THR	2.3
1	D	217	CYS	2.3
1	C	174	LYS	2.2
1	D	169	ALA	2.2
1	D	233	GLY	2.2
1	B	172	THR	2.2
1	B	168	TRP	2.2
1	B	180	GLN	2.2
1	D	32	LYS	2.1
1	A	32	LYS	2.1
1	C	224	PRO	2.1
1	D	204	THR	2.1
1	D	214	GLY	2.1
1	D	198	ASP	2.1
1	D	218	LYS	2.0
1	D	182	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.