



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 11:37 pm GMT

PDB ID : 2C6I  
Title : Crystal structure of the human CDK2 complexed with the triazolopyrimidine inhibitor  
Authors : Richardson, C.M.; Dokurno, P.; Murray, J.B.; Surgenor, A.E.  
Deposited on : 2005-11-10  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	<div> <div>52%</div> <div>71%</div> <div>25%</div> <div>..</div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2606 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	292	Total	C	N	O	S	0	0	0
			2345	1529	399	409	8			

- Molecule 2 is 4-{[5-(CYCLOHEXYLMETHOXY)[1,2,4]TRIAZOLO[1,5-A]PYRIMIDIN-7-YL]AMINO}BENZENESULFONAMIDE (three-letter code: DT1) (formula: C<sub>18</sub>H<sub>22</sub>N<sub>6</sub>O<sub>3</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			28	18	6	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	233	Total 233	0	0
			O 233		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: CELL DIVISION PROTEIN KINASE 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.58Å 71.61Å 72.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 1.80 23.70 – 1.78	Depositor EDS
% Data completeness (in resolution range)	97.1 (25.00-1.80) 94.1 (23.70-1.78)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 1.78Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.202 , 0.275 0.327 , 0.377	Depositor DCC
$R_{free}$ test set	2135 reflections (8.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	28.1	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.004 for -h,l,k	Xtriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	2606	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DT1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	0/2406	0.92	3/3264 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	ASP	CB-CG-OD2	7.19	124.77	118.30
1	A	92	ASP	CB-CG-OD2	7.14	124.73	118.30
1	A	262	LEU	CB-CG-CD2	5.50	120.35	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2345	0	2396	45	3
2	A	28	0	22	3	0
3	A	233	0	0	14	1
All	All	2606	0	2418	46	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:PRO:HD2	3:A:2106:HOH:O	1.57	1.04
1:A:177:CYS:SG	3:A:2134:HOH:O	2.34	0.85
1:A:180:TYR:O	3:A:2135:HOH:O	1.99	0.78
1:A:81:GLU:O	2:A:1299:DT1:H11	1.89	0.72
1:A:15:TYR:OH	1:A:149:ALA:HA	1.90	0.70
1:A:18:VAL:HG22	1:A:33:LYS:HG2	1.74	0.69
1:A:60:HIS:HD2	1:A:62:ASN:H	1.42	0.65
1:A:7:VAL:HG12	1:A:8:GLU:HG2	1.79	0.65
1:A:15:TYR:CD1	1:A:152:PHE:HB2	2.32	0.64
1:A:97:THR:HG21	3:A:2080:HOH:O	1.98	0.64
1:A:60:HIS:CD2	1:A:62:ASN:H	2.17	0.61
1:A:62:ASN:OD1	3:A:2049:HOH:O	2.17	0.61
1:A:293:VAL:HG12	3:A:2223:HOH:O	2.03	0.59
1:A:15:TYR:CE1	1:A:152:PHE:HB2	2.39	0.58
1:A:245:ARG:NH2	1:A:248:PHE:HE1	2.01	0.58
1:A:153:GLY:O	1:A:155:PRO:HD3	2.07	0.55
1:A:137:THR:C	3:A:2106:HOH:O	2.45	0.55
1:A:12:GLU:HB3	1:A:159:TYR:HE2	1.72	0.54
1:A:167:TRP:CD1	1:A:204:PRO:HA	2.43	0.53
1:A:2:GLU:HA	3:A:2020:HOH:O	2.08	0.53
1:A:156:VAL:HG12	1:A:157:ARG:HD3	1.91	0.51
1:A:44:VAL:HG21	1:A:76:LEU:HB2	1.91	0.51
1:A:106:SER:HB2	1:A:290:THR:O	2.10	0.51
1:A:245:ARG:NH2	1:A:248:PHE:CE1	2.78	0.51
1:A:3:ASN:ND2	3:A:2001:HOH:O	2.40	0.50
1:A:36:ARG:CB	3:A:2011:HOH:O	2.60	0.50
1:A:60:HIS:CD2	1:A:61:PRO:HD2	2.48	0.49
1:A:270:ASP:HB3	1:A:273:LYS:HD2	1.93	0.49
1:A:56:LYS:HD2	1:A:69:VAL:HG23	1.95	0.48
1:A:33:LYS:NZ	1:A:145:ASP:OD1	2.41	0.48
1:A:255:LEU:O	1:A:260:ARG:HD3	2.14	0.47
1:A:46:SER:O	1:A:50:ARG:HG3	2.15	0.46
1:A:257:GLU:OE1	1:A:260:ARG:NH2	2.48	0.46
1:A:132:ASN:HB3	2:A:1299:DT1:H6C2	1.98	0.45
1:A:205:GLY:HA2	1:A:210:ASP:OD2	2.16	0.45
1:A:262:LEU:HD22	1:A:266:MET:HG3	1.97	0.45
1:A:139:GLY:HA2	1:A:294:PRO:HD3	2.00	0.44
1:A:294:PRO:CD	3:A:2106:HOH:O	2.37	0.43
1:A:9:LYS:HD2	1:A:12:GLU:OE1	2.19	0.43
1:A:56:LYS:HZ2	1:A:69:VAL:H	1.67	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1299:DT1:C18	2:A:1299:DT1:C8	2.98	0.41
1:A:88:LYS:HG3	3:A:2075:HOH:O	2.19	0.41
1:A:181:SER:HB2	3:A:2138:HOH:O	2.21	0.41
1:A:176:GLY:O	1:A:234:PRO:HG2	2.21	0.41
1:A:29:VAL:HB	3:A:2025:HOH:O	2.19	0.40
1:A:5:GLN:HE21	1:A:22:ARG:HD2	1.85	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:ASN:ND2	1:A:272:ASN:O[2_464]	1.99	0.21
1:A:20:LYS:NZ	1:A:258:ASP:OD1[2_564]	2.00	0.20
1:A:74:ASN:N	1:A:272:ASN:OD1[2_464]	2.09	0.11
3:A:2109:HOH:O	3:A:2190:HOH:O[2_564]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/298 (97%)	281 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	256/263 (97%)	234 (91%)	22 (9%)	12 3

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	12	GLU
1	A	32	LEU
1	A	51	GLU
1	A	54	LEU
1	A	56	LYS
1	A	57	GLU
1	A	72	THR
1	A	75	LYS
1	A	96	LEU
1	A	124	LEU
1	A	126	ARG
1	A	146	PHE
1	A	150	ARG
1	A	165	THR
1	A	178	LYS
1	A	200	ARG
1	A	247	ASP
1	A	262	LEU
1	A	278	LYS
1	A	293	VAL
1	A	295	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	59	ASN
1	A	60	HIS
1	A	62	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	DT1	A	1299	-	25,31,31	1.47	2 (8%)	33,44,44	2.56	8 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DT1	A	1299	-	-	0/15/23/23	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1299	DT1	C16-S1	-2.88	1.72	1.77
2	A	1299	DT1	C7-N1	5.23	1.38	1.30

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1299	DT1	O2-S1-O3	-6.38	107.00	118.70
2	A	1299	DT1	C20-O1-C7	-6.00	109.91	117.27

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1299	DT1	C18-C17-C16	-2.97	116.25	119.46
2	A	1299	DT1	C17-C16-S1	-2.05	116.70	119.76
2	A	1299	DT1	C17-C16-C15	2.23	123.61	120.43
2	A	1299	DT1	O3-S1-C16	2.76	110.62	107.39
2	A	1299	DT1	O2-S1-N6	4.30	114.09	107.34
2	A	1299	DT1	N5-C9-N2	8.02	120.85	113.84

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1299	DT1	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	292/298 (97%)	2.31	156 (53%) <b>0</b> <b>0</b>	25, 38, 58, 72	21 (7%)

All (156) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	44	VAL	12.8
1	A	155	PRO	9.3
1	A	14	THR	8.6
1	A	96	LEU	8.4
1	A	154	VAL	8.3
1	A	160	THR	7.2
1	A	159	TYR	6.9
1	A	54	LEU	6.3
1	A	180	TYR	6.3
1	A	153	GLY	6.3
1	A	243	TRP	5.9
1	A	15	TYR	5.8
1	A	152	PHE	5.6
1	A	103	LEU	5.2
1	A	297	ARG	5.2
1	A	25	LEU	5.1
1	A	69	VAL	5.1
1	A	45	PRO	4.9
1	A	296	LEU	4.2
1	A	227	TRP	4.2
1	A	240	PHE	4.2
1	A	67	LEU	4.1
1	A	281	LEU	4.1
1	A	66	LEU	4.1
1	A	95	ALA	4.0
1	A	97	THR	4.0
1	A	262	LEU	3.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	80	PHE	3.8
1	A	46	SER	3.7
1	A	179	TYR	3.7
1	A	71	HIS	3.7
1	A	229	GLY	3.6
1	A	150	ARG	3.6
1	A	24	LYS	3.6
1	A	32	LEU	3.5
1	A	112	LEU	3.5
1	A	79	VAL	3.4
1	A	50	ARG	3.4
1	A	292	PRO	3.4
1	A	34	LYS	3.3
1	A	53	SER	3.3
1	A	193	PHE	3.3
1	A	287	GLN	3.3
1	A	146	PHE	3.2
1	A	63	ILE	3.2
1	A	98	GLY	3.2
1	A	108	LEU	3.2
1	A	233	MET	3.2
1	A	133	LEU	3.2
1	A	284	PRO	3.2
1	A	111	LEU	3.1
1	A	143	LEU	3.1
1	A	298	LEU	3.1
1	A	114	GLY	3.1
1	A	73	GLU	3.0
1	A	225	VAL	3.0
1	A	267	LEU	3.0
1	A	158	THR	3.0
1	A	18	VAL	3.0
1	A	184	VAL	2.9
1	A	246	GLN	2.9
1	A	293	VAL	2.9
1	A	247	ASP	2.9
1	A	94	SER	2.9
1	A	248	PHE	2.9
1	A	273	LYS	2.9
1	A	104	ILE	2.9
1	A	279	ALA	2.9
1	A	2	GLU	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	77	TYR	2.8
1	A	1	MET	2.8
1	A	263	LEU	2.8
1	A	194	ALA	2.8
1	A	186	ILE	2.8
1	A	278	LYS	2.8
1	A	282	ALA	2.8
1	A	121	HIS	2.8
1	A	230	VAL	2.8
1	A	144	ALA	2.7
1	A	78	LEU	2.7
1	A	13	GLY	2.7
1	A	161	HIS	2.7
1	A	19	TYR	2.7
1	A	170	ALA	2.7
1	A	280	ALA	2.7
1	A	156	VAL	2.7
1	A	148	LEU	2.7
1	A	215	ILE	2.7
1	A	59	ASN	2.7
1	A	64	VAL	2.7
1	A	37	LEU	2.7
1	A	87	LEU	2.7
1	A	116	ALA	2.6
1	A	90	PHE	2.6
1	A	60	HIS	2.6
1	A	226	VAL	2.6
1	A	101	LEU	2.6
1	A	51	GLU	2.6
1	A	16	GLY	2.6
1	A	117	PHE	2.6
1	A	151	ALA	2.6
1	A	22	ARG	2.6
1	A	289	VAL	2.6
1	A	35	ILE	2.6
1	A	72	THR	2.6
1	A	83	LEU	2.5
1	A	29	VAL	2.5
1	A	271	PRO	2.5
1	A	285	PHE	2.5
1	A	5	GLN	2.5
1	A	141	ILE	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	124	LEU	2.4
1	A	119	HIS	2.4
1	A	36	ARG	2.4
1	A	93	ALA	2.4
1	A	11	GLY	2.4
1	A	245	ARG	2.4
1	A	178	LYS	2.4
1	A	189	LEU	2.4
1	A	109	PHE	2.4
1	A	168	TYR	2.4
1	A	75	LYS	2.4
1	A	251	VAL	2.3
1	A	115	LEU	2.3
1	A	183	ALA	2.3
1	A	137	THR	2.3
1	A	290	THR	2.3
1	A	49	ILE	2.3
1	A	70	ILE	2.3
1	A	177	CYS	2.3
1	A	254	PRO	2.3
1	A	190	GLY	2.3
1	A	191	CYS	2.3
1	A	10	ILE	2.3
1	A	217	ARG	2.2
1	A	74	ASN	2.2
1	A	157	ARG	2.2
1	A	241	PRO	2.2
1	A	238	PRO	2.2
1	A	232	SER	2.2
1	A	173	ILE	2.1
1	A	286	PHE	2.1
1	A	128	LEU	2.1
1	A	162	GLU	2.1
1	A	4	PHE	2.1
1	A	255	LEU	2.1
1	A	26	THR	2.1
1	A	294	PRO	2.1
1	A	231	THR	2.1
1	A	187	TRP	2.1
1	A	99	ILE	2.1
1	A	269	TYR	2.0
1	A	30	VAL	2.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	192	ILE	2.0
1	A	295	HIS	2.0
1	A	234	PRO	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DT1	A	1299	28/28	0.86	0.18	-0.78	31,35,44,48	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.