



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2018 – 02:12 PM EST

PDB ID : 2C6O
Title : Crystal structure of the human CDK2 complexed with the triazolopyrimidine inhibitor
Authors : Richardson, C.M.; Dokurno, P.; Murray, J.B.; Surgenor, A.E.
Deposited on : 2005-11-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2409 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	290	Total	C	N	O	S	0	0	0
			2321	1514	393	406	8			

- Molecule 2 is O6-CYCLOHEXYLMETHOXY-2-(4'-SULPHAMOYLANILINO) PURINE (three-letter code: 4SP) (formula: C₁₈H₂₂N₆O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			28	18	6	3	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	60	Total	O	0
			60	60	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CELL DIVISION PROTEIN KINASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.04Å 71.76Å 71.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.75 – 2.10 9.97 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (29.75-2.10) 97.4 (9.97-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.82 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.1.09	Depositor
R, R_{free}	0.233 , 0.288 0.234 , 0.281	Depositor DCC
R_{free} test set	604 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.122	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.009 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2409	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 4SP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.26	4/2382 (0.2%)	1.26	17/3235 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	183	ALA	CA-CB	13.45	1.80	1.52
1	A	85	GLN	CB-CG	-5.34	1.38	1.52
1	A	294	PRO	CA-C	5.33	1.63	1.52
1	A	186	ILE	CA-CB	5.12	1.66	1.54

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	145	ASP	CB-CG-OD2	8.47	125.93	118.30
1	A	262	LEU	CB-CG-CD2	8.41	125.31	111.00
1	A	256	ASP	CB-CG-OD2	7.06	124.66	118.30
1	A	185	ASP	CB-CG-OD2	6.92	124.52	118.30
1	A	210	ASP	CB-CG-OD2	6.77	124.39	118.30
1	A	68	ASP	CB-CG-OD2	6.61	124.25	118.30
1	A	92	ASP	CB-CG-OD2	6.60	124.24	118.30
1	A	255	LEU	CB-CG-CD2	-6.50	99.94	111.00
1	A	235	ASP	CB-CG-OD2	6.44	124.10	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296	LEU	CA-CB-CG	6.15	129.44	115.30
1	A	128	LEU	CB-CG-CD1	-6.14	100.57	111.00
1	A	111	LEU	CB-CG-CD1	5.90	121.03	111.00
1	A	245	ARG	NE-CZ-NH2	-5.47	117.56	120.30
1	A	206	ASP	CB-CG-OD2	5.41	123.17	118.30
1	A	216	PHE	CB-CG-CD1	5.40	124.58	120.80
1	A	143	LEU	CA-CB-CG	-5.24	103.25	115.30
1	A	54	LEU	CA-CB-CG	5.23	127.33	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	161	HIS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2321	0	2361	109	0
2	A	28	0	22	4	0
3	A	60	0	0	9	0
All	All	2409	0	2383	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

All (111) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:VAL:HG22	1:A:45:PRO:CD	1.22	1.61
1:A:183:ALA:CA	1:A:183:ALA:CB	1.80	1.58
1:A:44:VAL:CG2	1:A:45:PRO:HD3	1.01	1.47
1:A:177:CYS:HB2	3:A:2029:HOH:O	1.23	1.25
1:A:44:VAL:CB	1:A:45:PRO:HD3	1.70	1.21

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:VAL:HG13	1:A:45:PRO:HD2	1.29	1.15
1:A:44:VAL:HG13	1:A:45:PRO:CD	1.76	1.13
1:A:165:THR:HG22	3:A:2025:HOH:O	1.49	1.10
1:A:1:MET:CE	1:A:6:LYS:HZ3	1.63	1.10
1:A:17:VAL:HG21	1:A:19:TYR:CE1	1.89	1.08
1:A:44:VAL:CG1	1:A:45:PRO:CD	2.32	1.07
1:A:157:ARG:HB3	1:A:161:HIS:HA	1.40	1.04
1:A:44:VAL:CG2	1:A:45:PRO:CD	1.95	1.03
1:A:44:VAL:CG1	1:A:45:PRO:HD2	1.91	1.00
1:A:1:MET:HE3	1:A:6:LYS:NZ	1.76	0.99
1:A:129:LYS:H	1:A:132:ASN:HD21	1.01	0.99
1:A:44:VAL:CB	1:A:45:PRO:CD	2.32	0.97
1:A:44:VAL:HG22	1:A:45:PRO:CG	1.95	0.97
1:A:1:MET:CE	1:A:6:LYS:NZ	2.32	0.90
1:A:1:MET:HE3	1:A:6:LYS:HZ3	1.27	0.88
1:A:60:HIS:HD2	1:A:62:ASN:H	1.22	0.88
1:A:1:MET:HE1	1:A:6:LYS:HZ3	1.40	0.86
1:A:60:HIS:CD2	1:A:62:ASN:H	1.94	0.85
1:A:83:LEU:HD21	1:A:142:LYS:HD2	1.57	0.85
1:A:125:HIS:HD2	1:A:127:ASP:H	1.25	0.80
2:A:1297:4SP:H22	2:A:1297:4SP:N1	1.97	0.80
1:A:47:THR:HB	3:A:2004:HOH:O	1.81	0.78
1:A:17:VAL:CG2	1:A:19:TYR:CE1	2.67	0.78
1:A:172:GLU:O	1:A:177:CYS:SG	2.44	0.74
1:A:129:LYS:H	1:A:132:ASN:ND2	1.83	0.74
1:A:129:LYS:N	1:A:132:ASN:HD21	1.82	0.73
1:A:44:VAL:HG13	1:A:45:PRO:N	2.01	0.73
1:A:132:ASN:HD22	1:A:132:ASN:H	1.36	0.73
1:A:18:VAL:HG22	1:A:33:LYS:HG2	1.72	0.71
1:A:213:PHE:O	1:A:217:ARG:HG3	1.91	0.70
1:A:129:LYS:O	1:A:132:ASN:ND2	2.24	0.70
1:A:125:HIS:CD2	1:A:127:ASP:H	2.12	0.66
1:A:177:CYS:SG	1:A:177:CYS:O	2.54	0.65
1:A:197:VAL:HG21	1:A:255:LEU:HD13	1.79	0.64
1:A:228:PRO:HB2	3:A:2049:HOH:O	1.99	0.63
1:A:158:THR:O	1:A:161:HIS:N	2.24	0.62
1:A:10:ILE:HG22	1:A:10:ILE:O	1.99	0.62
1:A:125:HIS:HE1	1:A:144:ALA:O	1.84	0.61
1:A:26:THR:HG23	1:A:28:GLU:HB2	1.83	0.61
1:A:18:VAL:HG22	1:A:33:LYS:CG	2.29	0.61
1:A:18:VAL:HG21	2:A:1297:4SP:H162	1.82	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:52:ILE:CG2	1:A:69:VAL:HG21	2.33	0.58
1:A:126:ARG:HG3	1:A:180:TYR:HE1	1.66	0.58
1:A:17:VAL:CG2	1:A:19:TYR:CD1	2.88	0.57
1:A:251:VAL:HG12	1:A:252:VAL:HG13	1.85	0.57
1:A:222:PRO:HG3	1:A:269:TYR:CE1	2.39	0.57
1:A:26:THR:CG2	1:A:28:GLU:HB2	2.35	0.56
1:A:124:LEU:CD2	1:A:182:THR:HG22	2.36	0.56
1:A:47:THR:CB	3:A:2004:HOH:O	2.48	0.56
1:A:15:TYR:HB2	1:A:33:LYS:HE2	1.89	0.54
1:A:180:TYR:HB2	1:A:184:VAL:HG11	1.90	0.54
1:A:52:ILE:HG21	1:A:69:VAL:HG21	1.90	0.53
1:A:125:HIS:HD2	1:A:127:ASP:N	2.00	0.53
1:A:183:ALA:C	1:A:183:ALA:CB	2.70	0.53
1:A:17:VAL:HG22	1:A:19:TYR:CD1	2.45	0.52
1:A:129:LYS:HA	1:A:192:ILE:HD11	1.90	0.52
2:A:1297:4SP:C22	2:A:1297:4SP:N1	2.60	0.52
1:A:294:PRO:HG2	1:A:294:PRO:O	2.09	0.51
1:A:270:ASP:OD1	1:A:272:ASN:HB2	2.11	0.51
1:A:172:GLU:HG2	1:A:173:ILE:N	2.26	0.51
1:A:60:HIS:HD2	1:A:62:ASN:N	2.02	0.50
1:A:227:TRP:O	1:A:230:VAL:HG12	2.11	0.50
1:A:156:VAL:HG23	1:A:157:ARG:H	1.75	0.49
1:A:181:SER:O	1:A:184:VAL:HG13	2.12	0.49
1:A:172:GLU:C	1:A:177:CYS:SG	2.91	0.49
1:A:44:VAL:HG22	1:A:45:PRO:HD3	0.48	0.48
1:A:126:ARG:HG3	1:A:180:TYR:CE1	2.48	0.48
1:A:89:LYS:HE2	3:A:2010:HOH:O	2.14	0.48
1:A:111:LEU:HD21	1:A:141:ILE:HG12	1.96	0.48
1:A:103:LEU:O	1:A:104:ILE:C	2.53	0.47
1:A:273:LYS:HG2	3:A:2050:HOH:O	2.14	0.47
1:A:160:THR:C	1:A:161:HIS:CD2	2.89	0.47
1:A:34:LYS:HG3	1:A:77:TYR:CE1	2.50	0.46
1:A:183:ALA:CB	1:A:183:ALA:N	2.67	0.46
1:A:139:GLY:HA2	1:A:294:PRO:HD3	1.97	0.46
1:A:125:HIS:CG	1:A:128:LEU:HD23	2.51	0.46
1:A:99:ILE:O	1:A:100:PRO:C	2.53	0.46
1:A:124:LEU:HD22	1:A:182:THR:HG22	1.97	0.45
1:A:163:VAL:HG12	1:A:164:VAL:O	2.16	0.45
1:A:62:ASN:ND2	1:A:110:GLN:HB3	2.31	0.45
1:A:172:GLU:C	1:A:177:CYS:HG	2.19	0.45
1:A:1:MET:HE3	1:A:6:LYS:HZ1	1.75	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:VAL:HG21	1:A:19:TYR:HE1	1.71	0.45
1:A:7:VAL:O	1:A:8:GLU:HB3	2.17	0.44
1:A:257:GLU:OE1	1:A:260:ARG:NH2	2.49	0.44
1:A:222:PRO:HG3	1:A:269:TYR:CZ	2.51	0.44
1:A:180:TYR:HB2	1:A:184:VAL:CG1	2.48	0.44
1:A:1:MET:HE1	1:A:6:LYS:NZ	2.13	0.44
1:A:172:GLU:HG2	1:A:173:ILE:H	1.81	0.44
1:A:60:HIS:CG	1:A:61:PRO:HD2	2.53	0.43
1:A:115:LEU:O	1:A:116:ALA:C	2.57	0.43
1:A:173:ILE:HD13	1:A:173:ILE:HG21	1.81	0.42
1:A:67:LEU:HA	1:A:67:LEU:HD23	1.92	0.42
1:A:72:THR:HB	1:A:73:GLU:OE1	2.19	0.42
1:A:111:LEU:HD13	1:A:133:LEU:HD13	2.02	0.42
1:A:275:ILE:HD13	1:A:275:ILE:HG21	1.64	0.42
1:A:47:THR:N	3:A:2004:HOH:O	2.52	0.41
1:A:132:ASN:HD22	1:A:132:ASN:N	2.02	0.41
1:A:156:VAL:HG22	3:A:2021:HOH:O	2.20	0.41
1:A:187:TRP:CD1	1:A:187:TRP:C	2.94	0.41
1:A:160:THR:C	1:A:161:HIS:CG	2.92	0.41
1:A:194:ALA:O	1:A:198:THR:HG23	2.20	0.41
1:A:13:GLY:HA3	2:A:1297:4SP:H152	2.02	0.41
1:A:198:THR:C	1:A:199:ARG:HG3	2.42	0.41
1:A:1:MET:HE3	1:A:6:LYS:CE	2.51	0.41
1:A:145:ASP:OD1	1:A:145:ASP:N	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	286/298 (96%)	266 (93%)	17 (6%)	3 (1%)	18 12

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	159	TYR
1	A	274	ARG
1	A	27	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	253/263 (96%)	230 (91%)	23 (9%)	11 7

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	3	ASN
1	A	12	GLU
1	A	17	VAL
1	A	26	THR
1	A	33	LYS
1	A	44	VAL
1	A	52	ILE
1	A	54	LEU
1	A	81	GLU
1	A	97	THR
1	A	126	ARG
1	A	132	ASN
1	A	138	GLU
1	A	156	VAL
1	A	161	HIS
1	A	180	TYR
1	A	199	ARG
1	A	200	ARG
1	A	242	LYS
1	A	262	LEU
1	A	272	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	294	PRO

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	60	HIS
1	A	71	HIS
1	A	125	HIS
1	A	132	ASN
1	A	161	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	4SP	A	1297	-	28,31,31	1.70	8 (28%)	35,44,44	2.47	16 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	4SP	A	1297	-	-	0/15/23/23	0/4/4/4

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1297	4SP	C4-N9	-4.16	1.27	1.34
2	A	1297	4SP	C17-N2	-2.71	1.35	1.40
2	A	1297	4SP	O6-C6	-2.63	1.33	1.35
2	A	1297	4SP	C5-N7	-2.50	1.30	1.39
2	A	1297	4SP	C4-N3	-2.01	1.32	1.36
2	A	1297	4SP	O6-C10	2.08	1.50	1.44
2	A	1297	4SP	O24-S23	2.31	1.48	1.43
2	A	1297	4SP	S23-N26	2.97	1.66	1.60

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1297	4SP	C15-C16-C11	-4.34	104.23	112.19
2	A	1297	4SP	O24-S23-C20	-3.97	102.75	107.39
2	A	1297	4SP	O25-S23-O24	-3.43	112.42	118.70
2	A	1297	4SP	N3-C2-N1	-3.22	121.36	126.23
2	A	1297	4SP	C16-C11-C12	-2.91	102.10	109.27
2	A	1297	4SP	C17-N2-C2	-2.45	122.79	129.17
2	A	1297	4SP	O6-C10-C11	-2.41	102.31	107.81
2	A	1297	4SP	O25-S23-N26	-2.26	103.78	107.34
2	A	1297	4SP	O24-S23-N26	3.03	112.10	107.34
2	A	1297	4SP	C6-C5-C4	3.05	120.21	117.26
2	A	1297	4SP	C14-C15-C16	3.23	118.10	111.42
2	A	1297	4SP	C12-C11-C10	3.40	118.43	111.42
2	A	1297	4SP	O25-S23-C20	3.49	111.47	107.39
2	A	1297	4SP	C2-N1-C6	3.88	122.03	115.18
2	A	1297	4SP	C20-S23-N26	4.31	114.64	108.41
2	A	1297	4SP	O6-C6-N1	4.92	125.28	120.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1297	4SP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	290/298 (97%)	0.94	41 (14%) 3 4	24, 38, 61, 90	17 (5%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	156	VAL	8.9
1	A	154	VAL	7.6
1	A	159	TYR	7.1
1	A	158	THR	6.7
1	A	44	VAL	5.8
1	A	155	PRO	5.5
1	A	37	LEU	5.1
1	A	160	THR	5.1
1	A	295	HIS	4.0
1	A	162	GLU	3.9
1	A	26	THR	3.7
1	A	157	ARG	3.7
1	A	230	VAL	3.6
1	A	247	ASP	3.4
1	A	246	GLN	3.4
1	A	164	VAL	3.3
1	A	3	ASN	3.2
1	A	54	LEU	3.2
1	A	296	LEU	3.2
1	A	139	GLY	3.1
1	A	177	CYS	3.1
1	A	25	LEU	3.0
1	A	23	ASN	2.9
1	A	243	TRP	2.8
1	A	287	GLN	2.7
1	A	75	LYS	2.7
1	A	153	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	15	TYR	2.5
1	A	72	THR	2.5
1	A	253	PRO	2.4
1	A	24	LYS	2.3
1	A	1	MET	2.3
1	A	14	THR	2.3
1	A	73	GLU	2.3
1	A	96	LEU	2.2
1	A	18	VAL	2.2
1	A	53	SER	2.2
1	A	74	ASN	2.1
1	A	10	ILE	2.1
1	A	150	ARG	2.0
1	A	47	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	4SP	A	1297	28/28	0.91	0.14	-0.75	31,39,52,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.