



Full wwPDB X-ray Structure Validation Report i

Feb 14, 2017 – 10:46 am GMT

PDB ID : 5C65
Title : Structure of the human glucose transporter GLUT3 / SLC2A3
Authors : Pike, A.C.W.; Quigley, A.; Chu, A.; Tessitore, A.; Xia, X.; Mukhopadhyay, S.; Wang, D.; Kupinska, K.; Strain-Damerell, C.; Chalk, R.; Burgess-Brown, N.A.; Edwards, A.M.; Arrowsmith, C.H.; Bountra, C.; Carpenter, E.P.; Structural Genomics Consortium (SGC)
Deposited on : 2015-06-22
Resolution : 2.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3491 (2.70-2.62)
Clashscore	112137	1026 (2.68-2.64)
Ramachandran outliers	110173	1010 (2.68-2.64)
Sidechain outliers	110143	1010 (2.68-2.64)
RSRZ outliers	101464	3511 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	37X	A	503	-	-	-	X
2	37X	A	506	-	-	-	X
3	Y01	A	507	-	-	-	X
3	Y01	B	504	-	-	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 7231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Solute carrier family 2, facilitated glucose transporter member 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	457	3443	2274	548	601	20	0	0	0
1	B	452	3343	2201	532	592	18	0	0	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	43	GLN	ASN	engineered mutation	UNP P11169
A	475	ALA	-	expression tag	UNP P11169
A	476	GLU	-	expression tag	UNP P11169
A	477	ASN	-	expression tag	UNP P11169
A	478	LEU	-	expression tag	UNP P11169
A	479	TYR	-	expression tag	UNP P11169
A	480	PHE	-	expression tag	UNP P11169
A	481	GLN	-	expression tag	UNP P11169
B	43	GLN	ASN	engineered mutation	UNP P11169
B	475	ALA	-	expression tag	UNP P11169
B	476	GLU	-	expression tag	UNP P11169
B	477	ASN	-	expression tag	UNP P11169
B	478	LEU	-	expression tag	UNP P11169
B	479	TYR	-	expression tag	UNP P11169
B	480	PHE	-	expression tag	UNP P11169
B	481	GLN	-	expression tag	UNP P11169

- Molecule 2 is Octyl Glucose Neopentyl Glycol (three-letter code: 37X) (formula: C₂₇H₅₂O₁₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			36	24	12		
2	A	1	Total	C	O	0	0
			39	27	12		
2	A	1	Total	C	O	0	0
			39	27	12		
2	A	1	Total	C	O	0	0
			39	27	12		
2	A	1	Total	C	O	0	0
			39	27	12		
2	B	1	Total	C	O	0	0
			39	27	12		
2	B	1	Total	C	O	0	0
			39	27	12		

- Molecule 3 is CHOLESTEROL HEMISUCCINATE (three-letter code: Y01) (formula: C₃₁H₅₀O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 35 31 4	0	0
3	B	1	Total C O 35 31 4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	18	Total O 18 18	0	0
4	B	9	Total O 9 9	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Solute carrier family 2, facilitated glucose transporter member 3



- Molecule 1: Solute carrier family 2, facilitated glucose transporter member 3



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.55 Å 116.03 Å 99.75 Å 90.00° 106.91° 90.00°	Depositor
Resolution (Å)	40.00 – 2.65 73.71 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.8 (40.00-2.65) 99.8 (73.71-2.65)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.76 (at 2.65 Å)	Xtriage
Refinement program	BUSTER 2.10.1	Depositor
R , R_{free}	0.205 , 0.223 0.214 , 0.235	Depositor DCC
R_{free} test set	2006 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	74.7	Xtriage
Anisotropy	0.191	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 89.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.031 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7231	wwPDB-VP
Average B, all atoms (Å ²)	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 37X, Y01

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.47	0/3519	0.60	0/4782
1	B	0.46	0/3413	0.60	0/4646
All	All	0.47	0/6932	0.60	0/9428

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3443	0	3502	16	0
1	B	3343	0	3324	13	0
2	A	231	0	303	0	0
2	B	117	0	156	4	0
3	A	35	0	49	1	0
3	B	35	0	49	1	0
4	A	18	0	0	0	0
4	B	9	0	0	0	0
All	All	7231	0	7383	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:ILE:HD13	1:B:395:GLN:HG3	1.75	0.67
1:A:214:ILE:HD13	1:A:395:GLN:HG3	1.78	0.65
1:B:83:LEU:HD11	2:B:503:37X:H51	1.85	0.57
1:B:226:LEU:HD23	1:B:235:VAL:HB	1.88	0.55
1:A:192:ILE:HG21	1:B:196:ILE:HD13	1.89	0.54
1:A:411:THR:HB	3:A:507:Y01:HAQ2	1.90	0.54
1:A:196:ILE:HD13	1:B:192:ILE:HG21	1.91	0.53
1:A:214:ILE:HG23	1:A:246:SER:HA	1.92	0.50
1:A:99:LEU:HG	2:B:501:37X:H19	1.92	0.50
1:A:226:LEU:HD23	1:A:235:VAL:HB	1.94	0.48
1:B:411:THR:HB	3:B:504:Y01:HAQ2	1.94	0.48
1:A:206:PRO:HG3	1:A:229:LEU:HD21	1.97	0.47
1:A:422:SER:O	1:A:426:TYR:HD2	1.97	0.47
1:B:83:LEU:HD21	2:B:503:37X:H52	1.98	0.46
1:A:14:ILE:HG13	1:A:145:ILE:HD11	1.98	0.45
1:B:14:ILE:HG13	1:B:145:ILE:HD11	1.99	0.45
1:A:99:LEU:HG	2:B:501:37X:CBD	2.47	0.45
1:B:272:ILE:HG23	1:B:450:VAL:HG22	1.99	0.44
1:A:272:ILE:HG23	1:A:450:VAL:HG22	2.00	0.43
1:A:342:MET:CE	1:A:381:PRO:HG2	2.49	0.43
1:B:342:MET:CE	1:B:381:PRO:HG2	2.49	0.42
1:A:457:THR:HG22	1:A:459:GLU:H	1.85	0.42
1:B:223:LYS:HA	1:B:239:ILE:HD11	2.02	0.41
1:B:177:ILE:HG22	1:B:178:LEU:HG	2.02	0.41
1:A:223:LYS:HA	1:A:239:ILE:HD11	2.03	0.41
1:A:255:VAL:HG11	1:A:397:PRO:HB3	2.03	0.40
1:B:74:GLY:HA3	1:B:410:TRP:CE3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	451/481 (94%)	441 (98%)	9 (2%)	1 (0%)	51 69
1	B	442/481 (92%)	433 (98%)	8 (2%)	1 (0%)	51 69
All	All	893/962 (93%)	874 (98%)	17 (2%)	2 (0%)	51 69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	380	GLY
1	B	380	GLY

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	361/397 (91%)	357 (99%)	4 (1%)	78 90
1	B	340/397 (86%)	336 (99%)	4 (1%)	75 89
All	All	701/794 (88%)	693 (99%)	8 (1%)	78 90

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	ILE
1	A	245	GLU
1	A	315	ASN
1	A	422	SER
1	B	196	ILE
1	B	239	ILE
1	B	315	ASN
1	B	422	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	37X	A	501	-	37,37,40	0.25	0	49,51,54	1.01	2 (4%)
2	37X	A	502	-	40,40,40	0.23	0	52,54,54	1.12	4 (7%)
2	37X	A	503	-	40,40,40	0.23	0	52,54,54	0.64	2 (3%)
2	37X	A	504	-	40,40,40	0.31	0	52,54,54	0.69	1 (1%)
2	37X	A	505	-	40,40,40	0.29	0	52,54,54	0.93	4 (7%)
2	37X	A	506	-	40,40,40	0.28	0	52,54,54	0.54	1 (1%)
3	Y01	A	507	-	35,38,38	0.43	0	54,57,57	0.56	0
2	37X	B	501	-	40,40,40	0.25	0	52,54,54	0.74	2 (3%)
2	37X	B	502	-	40,40,40	0.25	0	52,54,54	0.60	2 (3%)
2	37X	B	503	-	40,40,40	0.26	0	52,54,54	0.59	0
3	Y01	B	504	-	35,38,38	0.38	0	54,57,57	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	37X	A	501	-	-	0/27/67/70	0/2/2/2
2	37X	A	502	-	-	0/30/70/70	0/2/2/2
2	37X	A	503	-	-	0/30/70/70	0/2/2/2
2	37X	A	504	-	-	0/30/70/70	0/2/2/2
2	37X	A	505	-	-	0/30/70/70	0/2/2/2
2	37X	A	506	-	-	0/30/70/70	0/2/2/2
3	Y01	A	507	-	-	0/17/77/77	0/4/4/4
2	37X	B	501	-	-	0/30/70/70	0/2/2/2
2	37X	B	502	-	-	0/30/70/70	0/2/2/2
2	37X	B	503	-	-	0/30/70/70	0/2/2/2
3	Y01	B	504	-	-	0/17/77/77	0/4/4/4

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	505	37X	CBT-CCM-CBS	-3.43	90.89	109.59
2	A	502	37X	CBR-CCM-CBQ	-3.03	104.29	109.97
2	B	501	37X	CBQ-CCM-CBT	-2.07	104.50	109.40
2	B	502	37X	CBS-O1-C1	2.08	118.99	113.44
2	A	505	37X	CBS-O1-C1	2.11	119.07	113.44
2	A	503	37X	CBS-O1-C1	2.30	119.57	113.44
2	B	502	37X	CBT-OBV-CCJ	2.31	119.60	113.44
2	A	506	37X	CBS-O1-C1	2.34	119.69	113.44
2	A	504	37X	CBS-O1-C1	2.90	121.19	113.44
2	A	505	37X	CBQ-CCM-CBT	2.91	116.29	109.40
2	B	501	37X	CBS-O1-C1	2.94	121.30	113.44
2	A	503	37X	CBT-OBV-CCJ	2.97	121.37	113.44
2	A	505	37X	CBT-OBV-CCJ	3.04	121.56	113.44
2	A	502	37X	CBT-OBV-CCJ	3.20	121.98	113.44
2	A	501	37X	CBT-OBV-CCJ	3.65	123.19	113.44
2	A	502	37X	CBQ-CCM-CBS	4.04	118.98	109.40
2	A	501	37X	CBS-O1-C1	4.16	124.55	113.44
2	A	502	37X	CBS-O1-C1	4.52	125.49	113.44

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	507	Y01	1	0
2	B	501	37X	2	0
2	B	503	37X	2	0
3	B	504	Y01	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	457/481 (95%)	0.69	27 (5%) 23 21	61, 88, 133, 181	0
1	B	452/481 (93%)	0.81	62 (13%) 3 2	66, 108, 164, 220	0
All	All	909/962 (94%)	0.75	89 (9%) 8 6	61, 96, 152, 220	0

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	235	VAL	5.8
1	B	442	PHE	5.8
1	A	363	SER	5.3
1	B	225	ILE	5.2
1	B	295	ILE	5.1
1	A	362	MET	4.7
1	B	239	ILE	4.6
1	B	176	PHE	4.5
1	B	430	TYR	4.5
1	B	83	LEU	4.4
1	A	324	PHE	4.2
1	B	231	GLY	4.1
1	B	318	PHE	3.9
1	B	369	ALA	3.9
1	B	446	THR	3.8
1	A	57	VAL	3.8
1	B	177	ILE	3.7
1	B	232	THR	3.6
1	A	177	ILE	3.6
1	A	351	VAL	3.5
1	B	354	LEU	3.4
1	B	12	PHE	3.4
1	B	226	LEU	3.4
1	B	418	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	9	ALA	3.2
1	B	291	TYR	3.1
1	B	238	ASP	3.0
1	B	275	VAL	3.0
1	B	296	PHE	3.0
1	B	316	THR	3.0
1	A	430	TYR	3.0
1	A	295	ILE	2.9
1	A	367	ILE	2.9
1	B	303	GLU	2.9
1	B	205	CYS	2.9
1	B	234	ASP	2.9
1	A	12	PHE	2.9
1	B	444	ALA	2.9
1	B	459	GLU	2.8
1	A	11	ILE	2.8
1	B	11	ILE	2.7
1	A	317	ILE	2.7
1	B	309	ILE	2.7
1	B	228	ARG	2.7
1	A	200	ALA	2.7
1	A	16	VAL	2.7
1	A	365	VAL	2.6
1	A	444	ALA	2.6
1	A	320	VAL	2.6
1	A	442	PHE	2.6
1	B	240	GLN	2.6
1	B	212	LEU	2.5
1	B	79	PHE	2.5
1	B	376	PHE	2.5
1	B	145	ILE	2.5
1	A	344	PHE	2.4
1	A	54	PRO	2.4
1	A	321	VAL	2.4
1	B	222	ALA	2.4
1	B	14	ILE	2.4
1	B	317	ILE	2.4
1	B	216	ARG	2.4
1	B	276	LEU	2.3
1	B	279	SER	2.3
1	B	349	MET	2.3
1	B	229	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	167	LEU	2.3
1	B	319	THR	2.3
1	B	41	PHE	2.2
1	B	204	PHE	2.2
1	A	235	VAL	2.2
1	B	370	ILE	2.2
1	B	230	TRP	2.2
1	B	172	PHE	2.2
1	B	224	GLN	2.2
1	B	254	GLN	2.2
1	B	314	VAL	2.2
1	B	339	LEU	2.2
1	B	307	ALA	2.2
1	B	171	ILE	2.2
1	A	364	PHE	2.2
1	B	287	ALA	2.1
1	B	45	THR	2.1
1	A	454	ARG	2.1
1	B	458	PHE	2.1
1	B	371	LEU	2.0
1	A	259	GLU	2.0
1	A	433	ILE	2.0
1	A	201	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	37X	A	503	39/39	0.80	0.43	3.24	98,119,145,146	0
2	37X	A	506	39/39	0.53	0.38	3.23	123,158,161,161	0
3	Y01	A	507	35/35	0.78	0.36	2.79	122,130,140,142	0
3	Y01	B	504	35/35	0.73	0.34	2.16	142,151,158,159	0
2	37X	A	504	39/39	0.81	0.32	1.79	101,137,144,147	0
2	37X	A	502	39/39	0.87	0.29	0.86	95,101,116,118	0
2	37X	B	501	39/39	0.92	0.28	0.27	94,99,102,108	0
2	37X	A	505	39/39	0.84	0.27	0.06	103,112,128,130	0
2	37X	B	503	39/39	0.82	0.28	-0.04	110,136,153,154	0
2	37X	B	502	39/39	0.85	0.26	-0.11	115,148,153,154	0
2	37X	A	501	36/39	0.89	0.22	-0.58	86,95,106,112	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.