



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 02:12 am GMT

PDB ID : 1C7G
Title : TYROSINE PHENOL-LYASE FROM ERWINIA HERBICOLA
Authors : Mikami, B.; Yamamoto, Y.; Katayama, T.; Suzuki, H.
Deposited on : 2000-02-18
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	456	
1	B	456	
1	C	456	
1	D	456	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PLP	A	1000	-	X	-	-
2	PLP	B	1000	-	X	-	-
2	PLP	C	1000	-	X	-	-
2	PLP	D	1000	-	X	-	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 14905 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TYROSINE PHENOL-LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			
1	B	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			
1	C	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			
1	D	456	Total	C	N	O	S	0	0	0
			3605	2282	618	678	27			

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		
3	B	77	Total	O	0	0
			77	77		
3	C	158	Total	O	0	0
			158	158		
3	D	87	Total	O	0	0
			87	87		

Note EDS was not executed.

Chain A:

76% 21%

400


100

0

M1 V16 S17 M18 I19 S20 R21 D22 E23 R24 V25 K26 K27 M28 Q29 E30 F36 L37 K41 L45 L48 S51 D58 E75 K91 H97 Q98 G99 R100 E103 M104 L105 Q108 I111 M121 Y122 F123 T124 T125 T126 K132 M133 G134 A135 R142 D143 E144 L150 M151 L152 K155 L166 I167 K168 E169 E173 L180 A181 V182 L186 A187 G188 K210 Y213 D214 A215 T216 G255 K256 K257 C258 C259 M269 E272 E273 F274 F275 K279 L284 R307 R318 L325 G341 F346 P350 H355 L356 P362 A363 Q364 S365 A367 A368 Y371 M372 E373 T374 G375 V376 M379 E380 R381 R388 S389 K390 E391 T392 G393 E394 K395 H396 R397 P398 K399 L400 E401 T402 Y403 R404 L405 T406 R410 V420 G423 E432 V440 Y441 E442 P443 L446 R447 T450


Chain B:

[illegible]



● Molecule 1: TYROSINE PHENOL-LYASE

Chain D: 73% 23% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	163.49Å 113.04Å 101.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.10	Depositor
% Data completeness (in resolution range)	78.6 (8.00-2.10)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
Refinement program	X-PLOR 3.85	Depositor
R, R_{free}	0.186 , 0.242	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	14905	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3678	0.60	1/4955 (0.0%)
1	B	0.35	0/3678	0.59	1/4955 (0.0%)
1	C	0.36	0/3678	0.62	1/4955 (0.0%)
1	D	0.35	0/3678	0.59	1/4955 (0.0%)
All	All	0.35	0/14712	0.60	4/19820 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ASP	N-CA-C	-7.04	91.98	111.00
1	C	214	ASP	N-CA-C	-6.72	92.86	111.00
1	D	214	ASP	N-CA-C	-6.03	94.71	111.00
1	B	214	ASP	N-CA-C	-5.65	95.75	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	223	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3605	0	3548	67	0
1	B	3605	0	3548	87	1
1	C	3605	0	3548	70	0
1	D	3605	0	3548	73	0
2	A	15	0	6	1	0
2	B	15	0	6	1	0
2	C	15	0	6	2	0
2	D	15	0	6	1	0
3	A	103	0	0	2	0
3	B	77	0	0	3	0
3	C	158	0	0	3	0
3	D	87	0	0	0	0
All	All	14905	0	14216	289	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 289 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:28:MET:HE3	1:C:368:ALA:HA	1.43	1.01
1:C:97:HIS:HA	1:C:294:LEU:HD13	1.59	0.83
1:B:362:PRO:HG2	1:B:401:GLU:HG2	1.61	0.83
1:A:374:THR:OG1	1:A:376:VAL:HG13	1.79	0.82
1:B:374:THR:OG1	1:B:376:VAL:HG13	1.79	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:22:ASP:OD1	1:B:22:ASP:OD1[2_575]	1.80	0.40

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/456 (100%)	437 (96%)	17 (4%)	0	100	100
1	B	454/456 (100%)	433 (95%)	20 (4%)	1 (0%)	51	52
1	C	454/456 (100%)	438 (96%)	16 (4%)	0	100	100
1	D	454/456 (100%)	433 (95%)	21 (5%)	0	100	100
All	All	1816/1824 (100%)	1741 (96%)	74 (4%)	1 (0%)	55	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	391	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/380 (100%)	334 (88%)	46 (12%)	6	3
1	B	380/380 (100%)	333 (88%)	47 (12%)	5	3
1	C	380/380 (100%)	339 (89%)	41 (11%)	7	4
1	D	380/380 (100%)	338 (89%)	42 (11%)	7	4
All	All	1520/1520 (100%)	1344 (88%)	176 (12%)	6	3

5 of 176 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	388	ARG
1	C	48	LEU
1	D	318	ARG
1	B	391	GLU
1	B	446	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	161	ASN
1	B	130	GLN
1	C	130	GLN
1	D	130	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	1000	1	15,15,16	3.68	4 (26%)	20,22,23	4.06	14 (70%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	B	1000	1	15,15,16	3.71	5 (33%)	20,22,23	4.05	14 (70%)
2	PLP	C	1000	1	15,15,16	3.66	5 (33%)	20,22,23	3.98	14 (70%)
2	PLP	D	1000	1	15,15,16	3.51	5 (33%)	20,22,23	3.96	14 (70%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	1000	1	-	0/6/6/8	0/1/1/1
2	PLP	B	1000	1	-	0/6/6/8	0/1/1/1
2	PLP	C	1000	1	-	0/6/6/8	0/1/1/1
2	PLP	D	1000	1	-	0/6/6/8	0/1/1/1

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1000	PLP	C4A-C4	-2.40	1.46	1.51
2	C	1000	PLP	C4A-C4	-2.32	1.46	1.51
2	C	1000	PLP	P-O3P	-2.13	1.46	1.54
2	A	1000	PLP	C4A-C4	-2.08	1.47	1.51
2	B	1000	PLP	C4A-C4	-2.02	1.47	1.51

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1000	PLP	O3P-P-O4P	-9.54	81.34	106.73
2	B	1000	PLP	O3P-P-O4P	-9.50	81.45	106.73
2	C	1000	PLP	O3P-P-O4P	-9.31	81.96	106.73
2	D	1000	PLP	O3P-P-O4P	-9.23	82.16	106.73
2	D	1000	PLP	O3P-P-O2P	-6.23	82.49	107.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1000	PLP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1000	PLP	1	0
2	C	1000	PLP	2	0
2	D	1000	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.