



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 26, 2017 – 04:19 PM EDT

PDB ID : 3C72  
Title : Engineered RabGGTase in complex with a peptidomimetic inhibitor  
Authors : Guo, Z.; Wu, Y.W.; Tan, K.T.; Bon, R.S.; Guiu-Rozas, E.; Delon, C.; Nguyen, U.T.; Wetzel, S.; Arndt, S.; Goody, R.S.; Blankenfeldt, W.; Alexandrov, K.; Waldmann, H.  
Deposited on : unknown  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	334	<div> <div>8%</div> <div>74%</div> <div>16%</div> <div>• 8%</div> </div>
2	B	331	<div> <div>3%</div> <div>76%</div> <div>18%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CX1	B	334	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5255 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranylgeranyl transferase type-2 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	3	0
			2505	1598	430	465	12			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP Q08602
A	-2	HIS	-	EXPRESSION TAG	UNP Q08602
A	-1	GLY	-	EXPRESSION TAG	UNP Q08602
A	0	HIS	-	EXPRESSION TAG	UNP Q08602
A	237	GLY	-	LINKER	UNP Q08602
A	238	ALA	-	LINKER	UNP Q08602
A	239	GLY	-	LINKER	UNP Q08602
A	240	SER	-	LINKER	UNP Q08602
A	241	GLY	-	LINKER	UNP Q08602

- Molecule 2 is a protein called Geranylgeranyl transferase type-2 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	324	Total	C	N	O	S	0	1	0
			2512	1606	416	470	20			


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0

- Molecule 5 is N-[(benzyloxy)carbonyl]-L-histidyl-N-methyl-L-phenylalanyl-L-tyrosine (three-letter code: CX1) (formula: C<sub>33</sub>H<sub>35</sub>N<sub>5</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C N O 45 33 5 7	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	78	Total O 78 78	0	0
6	B	113	Total O 113 113	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Geranylgeranyl transferase type-2 subunit alpha



- Molecule 2: Geranylgeranyl transferase type-2 subunit beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.19 Å 91.67 Å 115.44 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.88 – 2.30 19.88 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.88-2.30) 99.9 (19.88-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	5.20	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.04 (at 2.30 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.189 , 0.256 0.194 , 0.263	Depositor DCC
$R_{free}$ test set	1587 reflections (5.25%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.145	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 43.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5255	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CX1, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.86	1/2570 (0.0%)	0.82	5/3488 (0.1%)
2	B	0.95	2/2574 (0.1%)	0.87	5/3494 (0.1%)
All	All	0.91	3/5144 (0.1%)	0.85	10/6982 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	303	VAL	CB-CG2	-7.37	1.37	1.52
2	B	43	GLU	CG-CD	5.50	1.60	1.51
2	B	183	CYS	CB-SG	-5.03	1.73	1.81

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	232	ARG	NE-CZ-NH1	6.09	123.34	120.30
1	A	150	ARG	NE-CZ-NH2	5.79	123.19	120.30
2	B	136	ASP	CB-CG-OD1	5.70	123.43	118.30
1	A	69	ARG	NE-CZ-NH2	-5.65	117.47	120.30
2	B	232	ARG	NE-CZ-NH2	-5.58	117.51	120.30
1	A	101	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	A	127	ARG	NE-CZ-NH2	-5.50	117.55	120.30
2	B	130	ASP	CB-CG-OD1	5.45	123.21	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	184	ARG	CG-CD-NE	-5.36	100.54	111.80
1	A	139	ASP	CB-CG-OD1	5.07	122.86	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	242	ARG	Peptide
1	A	243	CYS	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2505	0	2431	29	0
2	B	2512	0	2434	55	0
3	B	1	0	0	0	0
4	B	1	0	0	0	0
5	B	45	0	34	5	0
6	A	78	0	0	2	0
6	B	113	0	0	3	0
All	All	5255	0	4899	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:323:VAL:HG23	2:B:325:VAL:HG13	1.57	0.86
2:B:112[A]:HIS:ND1	6:B:447:HOH:O	2.26	0.68
1:A:188:LEU:HD11	1:A:211:GLU:CG	2.24	0.68
2:B:195:TYR:HE2	5:B:334:CX1:H3	1.58	0.67
1:A:242:ARG:N	1:A:243:CYS:HA	2.10	0.66
1:A:188:LEU:HD11	1:A:211:GLU:HG3	1.77	0.66

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:195:TYR:CE2	5:B:334:CX1:H3	2.31	0.66
1:A:116:LEU:O	1:A:120:LEU:HD13	1.96	0.65
1:A:25:LEU:O	1:A:29:GLN:HG3	1.98	0.64
1:A:192:HIS:CE1	1:A:207:VAL:HG21	2.32	0.64
1:A:174:ASN:OD1	6:A:408:HOH:O	2.16	0.63
1:A:203:LEU:HD13	1:A:238:ALA:HB1	1.80	0.62
2:B:14:ALA:HB1	2:B:15:PRO:HD2	1.82	0.62
2:B:255:LEU:HD12	2:B:255:LEU:O	2.00	0.61
1:A:28:TYR:CE1	1:A:55:ILE:HG23	2.35	0.61
5:B:334:CX1:H31	6:B:359:HOH:O	2.01	0.61
2:B:8:VAL:HG22	2:B:9:THR:N	2.17	0.60
1:A:51:LEU:O	1:A:55:ILE:HD12	2.02	0.60
1:A:188:LEU:CD1	1:A:211:GLU:HG3	2.32	0.59
2:B:111:ILE:CD1	2:B:112[B]:HIS:CD2	2.86	0.59
2:B:111:ILE:HD11	2:B:112[B]:HIS:CD2	2.40	0.57
1:A:225:ASP:O	1:A:225:ASP:OD1	2.24	0.56
2:B:246:LEU:HD11	2:B:255:LEU:CD1	2.36	0.56
2:B:325:VAL:HG22	2:B:325:VAL:O	2.06	0.55
1:A:45:ASP:OD2	1:A:47:SER:OG	2.25	0.55
1:A:300:LEU:HD13	1:A:311:LEU:HD22	1.89	0.55
2:B:45:LEU:C	2:B:45:LEU:HD12	2.28	0.54
2:B:21:GLU:HG2	2:B:22:LYS:H	1.72	0.54
1:A:49:LEU:HD21	1:A:72:VAL:HG22	1.90	0.54
1:A:225:ASP:O	1:A:225:ASP:CG	2.46	0.53
2:B:277:GLY:HA3	2:B:286:VAL:HG21	1.90	0.53
2:B:277:GLY:C	2:B:286:VAL:CG2	2.77	0.53
1:A:300:LEU:HD13	1:A:311:LEU:CD2	2.39	0.52
2:B:222:ARG:HD3	2:B:230:ASN:OD1	2.10	0.52
2:B:246:LEU:HD11	2:B:255:LEU:HD11	1.91	0.51
2:B:246:LEU:CD1	2:B:255:LEU:HD13	2.40	0.51
2:B:95:LEU:HD23	2:B:145:PHE:HB3	1.93	0.51
1:A:31:ALA:HB1	1:A:55:ILE:HD11	1.92	0.50
1:A:220:PHE:CZ	1:A:259:SER:HB3	2.47	0.49
5:B:334:CX1:C43	5:B:334:CX1:H13	2.42	0.49
2:B:62:GLN:HB3	2:B:65:ARG:HD3	1.94	0.49
2:B:323:VAL:HG23	2:B:325:VAL:CG1	2.35	0.49
2:B:8:VAL:CG2	2:B:9:THR:N	2.76	0.49
2:B:238:ASP:OD1	2:B:238:ASP:C	2.51	0.49
2:B:258:ILE:HG22	2:B:260:ARG:HG3	1.94	0.48
2:B:224:LEU:CD1	2:B:237:PRO:HD3	2.42	0.48
2:B:164:VAL:HG23	6:B:410:HOH:O	2.12	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:69:GLU:O	2:B:73:VAL:HG23	2.14	0.47
2:B:111:ILE:HD12	2:B:112[B]:HIS:CD2	2.49	0.47
2:B:232:ARG:HB2	2:B:233:PRO:HD2	1.96	0.47
1:A:215:VAL:CG1	1:A:216:GLN:N	2.77	0.47
2:B:54:LEU:HD12	2:B:66:MET:HG3	1.96	0.46
2:B:18:LEU:HD22	2:B:303:GLU:HG2	1.97	0.46
1:A:68:CYS:O	1:A:72:VAL:HG13	2.15	0.46
1:A:177:ASN:ND2	6:A:408:HOH:O	2.46	0.46
1:A:214:LEU:O	1:A:217:ASN:HB2	2.16	0.46
2:B:220:CYS:HA	2:B:258:ILE:HD11	1.97	0.45
2:B:277:GLY:CA	2:B:286:VAL:HG21	2.47	0.45
2:B:100:SER:HA	2:B:103:GLN:HE21	1.82	0.44
2:B:111:ILE:HD12	2:B:112[B]:HIS:CG	2.52	0.44
2:B:8:VAL:CG2	2:B:226:SER:HB3	2.47	0.44
1:A:18:ARG:NE	1:A:22:GLU:HG3	2.33	0.43
2:B:220:CYS:HA	2:B:258:ILE:CD1	2.48	0.43
2:B:303:GLU:HB3	2:B:306:ILE:HD12	2.00	0.43
2:B:278:PHE:C	2:B:286:VAL:HG23	2.39	0.43
2:B:278:PHE:HB2	2:B:291:THR:HA	2.01	0.43
2:B:246:LEU:HD12	2:B:255:LEU:HD13	2.00	0.43
1:A:64:THR:O	1:A:67:ASN:HB2	2.19	0.43
2:B:95:LEU:HD13	2:B:126:LEU:CD1	2.49	0.43
2:B:52:TRP:HA	2:B:292:LEU:HD22	2.01	0.43
2:B:232:ARG:HB2	2:B:233:PRO:CD	2.49	0.42
2:B:115:ASN:C	2:B:115:ASN:HD22	2.23	0.42
2:B:325:VAL:O	2:B:326:GLN:HG3	2.20	0.42
1:A:212:LEU:O	1:A:215:VAL:HG12	2.20	0.42
2:B:277:GLY:C	2:B:286:VAL:HG21	2.40	0.42
2:B:281:ARG:HB2	2:B:282:PRO:CD	2.50	0.42
1:A:86:ALA:O	1:A:89:VAL:HG22	2.20	0.41
2:B:300:LEU:HA	2:B:300:LEU:HD23	1.90	0.41
5:B:334:CX1:H17	5:B:334:CX1:H12	1.40	0.41
2:B:330:VAL:HG12	2:B:331:SER:N	2.36	0.41
2:B:54:LEU:HD23	2:B:104:ILE:HG23	2.01	0.41
1:A:141:ARG:HD3	1:A:176:SER:OG	2.21	0.40
2:B:246:LEU:CD1	2:B:255:LEU:CD1	2.98	0.40
2:B:45:LEU:O	2:B:45:LEU:HD12	2.20	0.40
2:B:67:ASN:HD22	2:B:67:ASN:C	2.25	0.40
1:A:49:LEU:HG	1:A:72:VAL:HG21	2.03	0.40
2:B:278:PHE:N	2:B:286:VAL:CG2	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/334 (91%)	291 (96%)	11 (4%)	1 (0%)	44	55
2	B	321/331 (97%)	305 (95%)	15 (5%)	1 (0%)	44	55
All	All	624/665 (94%)	596 (96%)	26 (4%)	2 (0%)	44	55

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	21	GLU
1	A	329	TYR

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/294 (91%)	249 (93%)	19 (7%)	17	22
2	B	272/284 (96%)	251 (92%)	21 (8%)	15	18
All	All	540/578 (93%)	500 (93%)	40 (7%)	16	20

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	GLN
1	A	53	SER
1	A	62	PHE
1	A	72	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	119	ARG
1	A	120	LEU
1	A	127	ARG
1	A	169	SER
1	A	215	VAL
1	A	225	ASP
1	A	227	SER
1	A	236	LEU
1	A	242	ARG
1	A	244	GLU
1	A	247	VAL
1	A	271	LYS
1	A	273	CYS
1	A	275	LEU
1	A	300	LEU
2	B	11	LYS
2	B	13	ASP
2	B	17	THR
2	B	32	SER
2	B	46	ARG
2	B	54	LEU
2	B	67	ASN
2	B	95	LEU
2	B	110	SER
2	B	115	ASN
2	B	156	LEU
2	B	159	LEU
2	B	222	ARG
2	B	234	GLU
2	B	236	LEU
2	B	238	ASP
2	B	240	CYS
2	B	255	LEU
2	B	258	ILE
2	B	300	LEU
2	B	325	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	54	GLN
1	A	192	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	264	GLN
2	B	67	ASN
2	B	91	HIS
2	B	103	GLN
2	B	115	ASN
2	B	193	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	CX1	B	334	-	41,48,48	1.29	2 (4%)	52,64,64	1.30	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	CX1	B	334	-	-	0/41/45/45	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	334	CX1	C7-C5	-5.28	1.38	1.50
5	B	334	CX1	C19-C33	2.58	1.57	1.51

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	334	CX1	O8-C9-O11	-2.90	118.40	124.22
5	B	334	CX1	C24-C23-N22	2.78	114.48	108.94
5	B	334	CX1	C17-N16-C18	3.14	123.43	117.80
5	B	334	CX1	O8-C9-N10	3.96	119.05	110.54
5	B	334	CX1	C7-O8-C9	4.09	125.56	115.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	334	CX1	5	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	306/334 (91%)	0.25	27 (8%) 11 15	23, 37, 61, 84	0
2	B	324/331 (97%)	0.05	11 (3%) 46 53	23, 34, 51, 66	0
All	All	630/665 (94%)	0.15	38 (6%) 23 29	23, 35, 57, 84	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	243	CYS	6.2
1	A	19	LEU	5.9
1	A	16	ALA	5.6
1	A	327	MET	5.4
1	A	21	ARG	5.1
2	B	13	ASP	4.6
1	A	242	ARG	4.5
1	A	18	ARG	4.4
1	A	175	PHE	4.4
1	A	176	SER	4.2
1	A	195	PRO	4.0
1	A	194	GLN	3.7
2	B	294	GLY	3.6
1	A	192	HIS	3.4
2	B	35	ASP	3.4
1	A	330	ALA	3.4
2	B	12	SER	3.3
1	A	121	PRO	3.2
2	B	331	SER	3.2
1	A	220	PHE	3.2
1	A	23	GLN	3.2
1	A	17	LYS	3.1
1	A	20	GLU	3.1
2	B	7	ASP	2.9

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	283	GLY	2.9
2	B	295	ILE	2.7
1	A	173	ARG	2.6
2	B	296	ALA	2.6
2	B	14	ALA	2.5
2	B	304	GLU	2.5
1	A	275	LEU	2.4
1	A	244	GLU	2.4
1	A	193	PRO	2.3
1	A	274	LEU	2.3
1	A	329	TYR	2.2
1	A	266	LEU	2.1
1	A	174	ASN	2.1
1	A	42	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	CX1	B	334	45/45	0.73	0.22	3.50	64,70,83,83	0
4	CA	B	333	1/1	0.96	0.04	-2.06	67,67,67,67	0
3	ZN	B	332	1/1	0.92	0.06	-	72,72,72,72	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.