



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2017 – 07:16 AM EDT

PDB ID : 3C7Q
Title : Structure of VEGFR2 kinase domain in complex with BIBF1120
Authors : Hilberg, F.; Roth, G.J.; Krssak, M.; Kautschitsch, S.; Sommergruber, W.;
Tontsch-Grunt, U.; Garin-Chesa, P.; Bader, G.; Zoephel, A.; Quant, J.; Heckel,
A.; Rettig, W.J.
Deposited on : unknown
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>7%</div> <div>72%</div> <div>19%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	3	-	-	-	X
2	SO4	A	6	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vascular endothelial growth factor receptor 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	291	Total	C	N	O	P	S	27	0	0
			2362	1508	407	426	2	19			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	804	GLY	-	EXPRESSION TAG	UNP P35968
A	805	SER	-	EXPRESSION TAG	UNP P35968
A	?	-	THR	DELETION	UNP P35968
A	?	-	LYS	DELETION	UNP P35968
A	?	-	GLY	DELETION	UNP P35968
A	?	-	ALA	DELETION	UNP P35968
A	?	-	ARG	DELETION	UNP P35968
A	?	-	PHE	DELETION	UNP P35968
A	?	-	ARG	DELETION	UNP P35968
A	?	-	GLN	DELETION	UNP P35968
A	?	-	GLY	DELETION	UNP P35968
A	?	-	LYS	DELETION	UNP P35968
A	?	-	ASP	DELETION	UNP P35968
A	?	-	TYR	DELETION	UNP P35968
A	?	-	VAL	DELETION	UNP P35968
A	?	-	GLY	DELETION	UNP P35968
A	?	-	ALA	DELETION	UNP P35968
A	?	-	ILE	DELETION	UNP P35968
A	?	-	PRO	DELETION	UNP P35968
A	?	-	VAL	DELETION	UNP P35968
A	?	-	ASP	DELETION	UNP P35968
A	?	-	LEU	DELETION	UNP P35968
A	?	-	LYS	DELETION	UNP P35968
A	?	-	ARG	DELETION	UNP P35968
A	?	-	ARG	DELETION	UNP P35968
A	?	-	LEU	DELETION	UNP P35968
A	?	-	ASP	DELETION	UNP P35968

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	DELETION	UNP P35968
A	?	-	ILE	DELETION	UNP P35968
A	?	-	THR	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	GLN	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	ALA	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	GLY	DELETION	UNP P35968
A	?	-	PHE	DELETION	UNP P35968
A	?	-	VAL	DELETION	UNP P35968
A	?	-	GLU	DELETION	UNP P35968
A	?	-	GLU	DELETION	UNP P35968
A	?	-	LYS	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	LEU	DELETION	UNP P35968
A	?	-	SER	DELETION	UNP P35968
A	?	-	ASP	DELETION	UNP P35968
A	?	-	VAL	DELETION	UNP P35968
A	?	-	GLU	DELETION	UNP P35968
A	?	-	GLU	DELETION	UNP P35968
A	?	-	GLU	DELETION	UNP P35968
A	?	-	GLU	DELETION	UNP P35968
A	?	-	ALA	DELETION	UNP P35968

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is methyl (3Z)-3-{[(4-{methyl[(4-methylpiperazin-1-yl)acetyl]amino}phenyl)amino](phenyl)methylidene}-2-oxo-2,3-dihydro-1H-indole-6-carboxylate (three-letter code: XIN) (formula: C₃₁H₃₃N₅O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			40	31	5	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	206	Total	O	0	0
			206	206		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vascular endothelial growth factor receptor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	38.25Å 94.44Å 96.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.54 – 2.10 28.54 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.54-2.10) 99.6 (28.54-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.91 (at 2.10Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.217 , 0.279 0.218 , 0.285	Depositor DCC
R_{free} test set	1050 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	26.9	Xtriage
Anisotropy	0.619	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.027 for -h,l,k	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2633	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CME, SO4, XIN, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/2349 (0.0%)	0.76	1/3162 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	907	LYS	CE-NZ	-7.07	1.31	1.49

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1058	ASP	CB-CG-OD1	6.11	123.80	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1053	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2362	0	2349	42	0
2	A	25	0	0	0	1
3	A	40	0	33	0	0
4	A	206	0	0	5	1
All	All	2633	0	2382	42	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1096:TRP:CE2	1:A:1125:MET:HE2	1.87	1.08
1:A:1096:TRP:CE2	1:A:1125:MET:CE	2.47	0.97
1:A:1072:MET:HE3	1:A:1076:THR:HG22	1.64	0.77
1:A:1096:TRP:NE1	1:A:1125:MET:HE2	2.02	0.74
1:A:1096:TRP:CZ2	1:A:1125:MET:CE	2.73	0.71
1:A:1096:TRP:CD2	1:A:1125:MET:HE1	2.29	0.68
1:A:1096:TRP:CE2	1:A:1125:MET:HE1	2.31	0.65
1:A:1096:TRP:CD2	1:A:1125:MET:CE	2.81	0.63
1:A:1064:ASP:O	4:A:1376:HOH:O	2.16	0.63
1:A:1072:MET:CE	1:A:1076:THR:HG22	2.27	0.63
1:A:1034:ILE:HG22	1:A:1042:VAL:HG22	1.85	0.57
1:A:849:ILE:HG22	1:A:850:GLU:O	2.04	0.57
1:A:854:PHE:HB2	1:A:862:CME:HE2	1.88	0.55
1:A:1171:ASP:C	1:A:1171:ASP:OD1	2.45	0.55
1:A:865:VAL:HG21	1:A:915:ILE:CG2	2.39	0.52
1:A:1153:PHE:O	1:A:1157:VAL:HG23	2.11	0.51
1:A:1031:ALA:HB2	1:A:1094:LEU:CD1	2.41	0.50
1:A:1052:ASP:HB3	4:A:1340:HOH:O	2.11	0.49
1:A:1059:PTR:HE2	1:A:1059:PTR:O2P	2.12	0.48
1:A:822:TYR:HB2	1:A:887:LYS:HE2	1.94	0.48
1:A:1096:TRP:CE3	1:A:1125:MET:HE1	2.49	0.48
1:A:1065:ALA:HA	4:A:1376:HOH:O	2.14	0.47
1:A:876:HIS:NE2	4:A:1274:HOH:O	2.34	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:883:MET:O	1:A:887:LYS:HG2	2.15	0.46
1:A:892:ILE:HD11	1:A:1024:CME:SD	2.55	0.46
1:A:1072:MET:HB3	1:A:1072:MET:HE2	1.77	0.45
1:A:1008:TYR:OH	1:A:1040:ASN:ND2	2.50	0.44
1:A:865:VAL:HG21	1:A:915:ILE:HG23	1.98	0.44
1:A:892:ILE:N	1:A:892:ILE:HD13	2.32	0.43
1:A:1077:ILE:HG21	1:A:1116:CYS:SG	2.59	0.43
1:A:892:ILE:HD12	1:A:1024:CME:SG	2.58	0.43
1:A:1038:GLU:HG2	1:A:1038:GLU:H	1.62	0.42
1:A:1107:PRO:HD3	1:A:1125:MET:HE3	2.01	0.42
1:A:854:PHE:HB2	1:A:862:CME:CE	2.48	0.42
1:A:867:VAL:CG1	1:A:913:MET:HB3	2.49	0.42
1:A:998:ASP:HA	4:A:1346:HOH:O	2.20	0.41
1:A:1127:ALA:HB1	1:A:1131:THR:HG21	2.03	0.41
1:A:1096:TRP:CZ2	1:A:1125:MET:HE2	2.41	0.41
1:A:843:GLY:O	1:A:871:LYS:NZ	2.46	0.41
1:A:1106:TYR:CE1	1:A:1125:MET:HG3	2.57	0.40
1:A:871:LYS:O	1:A:874:ALA:HB2	2.22	0.40
1:A:882:LEU:HD23	1:A:912:LEU:HD23	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1:SO4:O1	4:A:1275:HOH:O[1_455]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/316 (89%)	265 (95%)	13 (5%)	2 (1%)	25	20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1053	ILE
1	A	873	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	248/270 (92%)	232 (94%)	16 (6%)	20	16

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	835	LYS
1	A	852	ASP
1	A	863	ARG
1	A	998	ASP
1	A	1014	LYS
1	A	1035	LEU
1	A	1037	SER
1	A	1042	VAL
1	A	1047	PHE
1	A	1055	LYS
1	A	1061	ARG
1	A	1062	LYS
1	A	1079	ASP
1	A	1090	SER
1	A	1113	GLU
1	A	1170	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1040	ASN
1	A	1085	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CME	A	1007	1	9,9,10	0.68	0	6,9,11	1.54	1 (16%)
1	CME	A	1024	1	9,9,10	0.83	0	6,9,11	1.20	1 (16%)
1	PTR	A	1054	1	15,16,17	2.21	2 (13%)	19,22,24	1.23	3 (15%)
1	PTR	A	1059	1	15,16,17	1.97	2 (13%)	19,22,24	0.97	2 (10%)
1	CME	A	862	1	9,9,10	0.99	1 (11%)	6,9,11	2.24	3 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	A	1007	1	-	0/5/8/10	0/0/0/0
1	CME	A	1024	1	-	0/5/8/10	0/0/0/0
1	PTR	A	1054	1	-	0/9/11/13	0/1/1/1
1	PTR	A	1059	1	-	0/9/11/13	0/1/1/1
1	CME	A	862	1	-	0/5/8/10	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1054	PTR	OH-CZ	-7.68	1.22	1.40
1	A	1059	PTR	OH-CZ	-7.08	1.24	1.40
1	A	862	CME	CB-SG	-2.34	1.73	1.81
1	A	1059	PTR	P-O1P	-2.17	1.43	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1054	PTR	CA-C	3.36	1.54	1.50

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1054	PTR	CB-CA-C	-3.35	104.96	111.41
1	A	862	CME	CE-SD-SG	-2.57	90.49	103.40
1	A	1054	PTR	CG-CB-CA	-2.53	109.19	114.29
1	A	1054	PTR	O-C-CA	-2.39	118.42	125.02
1	A	1059	PTR	O-C-CA	-2.10	119.21	125.02
1	A	862	CME	O-C-CA	-2.00	119.49	125.02
1	A	1059	PTR	O2P-P-OH	2.03	112.87	105.63
1	A	1024	CME	CB-SG-SD	2.37	108.45	103.83
1	A	1007	CME	CB-SG-SD	3.31	110.27	103.83
1	A	862	CME	CB-SG-SD	4.30	112.21	103.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1024	CME	2	0
1	A	1059	PTR	1	0
1	A	862	CME	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1	-	4,4,4	0.19	0	6,6,6	0.18	0
3	XIN	A	1172	-	44,44,44	1.45	8 (18%)	61,62,62	1.70	13 (21%)
2	SO4	A	2	-	4,4,4	0.17	0	6,6,6	0.12	0
2	SO4	A	3	-	4,4,4	0.19	0	6,6,6	0.50	0
2	SO4	A	4	-	4,4,4	0.22	0	6,6,6	0.14	0
2	SO4	A	6	-	4,4,4	0.20	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1	-	-	0/0/0/0	0/0/0/0
3	XIN	A	1172	-	-	0/30/52/52	0/5/5/5
2	SO4	A	2	-	-	0/0/0/0	0/0/0/0
2	SO4	A	3	-	-	0/0/0/0	0/0/0/0
2	SO4	A	4	-	-	0/0/0/0	0/0/0/0
2	SO4	A	6	-	-	0/0/0/0	0/0/0/0

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1172	XIN	O10-C4	-4.37	1.15	1.23
3	A	1172	XIN	C19-C15	-3.10	1.34	1.39
3	A	1172	XIN	C15-N8	-3.09	1.35	1.41
3	A	1172	XIN	C1-C3	-2.84	1.34	1.39
3	A	1172	XIN	O25-C28	-2.36	1.39	1.45
3	A	1172	XIN	C2-C5	-2.31	1.38	1.41
3	A	1172	XIN	C6-C2	-2.20	1.36	1.39
3	A	1172	XIN	O26-C21	-2.01	1.17	1.22

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1172	XIN	C19-C15-N8	-2.79	111.06	120.41
3	A	1172	XIN	C28-O25-C21	-2.33	111.27	115.85
3	A	1172	XIN	C13-C7-C3	-2.06	118.28	120.75
3	A	1172	XIN	C31-N29-C27	2.14	120.57	117.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1172	XIN	C19-C23-C27	2.18	123.30	120.34
3	A	1172	XIN	C24-C20-C15	2.26	122.94	120.30
3	A	1172	XIN	C38-N39-C37	2.36	112.62	109.47
3	A	1172	XIN	C36-C38-N39	2.43	113.58	110.79
3	A	1172	XIN	C32-N34-C36	2.83	115.64	111.09
3	A	1172	XIN	C15-N8-C3	3.12	137.86	127.79
3	A	1172	XIN	C20-C15-N8	3.62	132.55	120.41
3	A	1172	XIN	C24-C27-N29	4.89	126.09	120.14
3	A	1172	XIN	C1-C3-N8	5.42	123.29	118.20

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	SO4	0	1

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	286/316 (90%)	0.48	23 (8%) 13 17	14, 28, 46, 52	7 (2%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1108	GLY	5.4
1	A	1078	PHE	4.0
1	A	938	TYR	3.9
1	A	874	ALA	3.6
1	A	1030	ALA	3.1
1	A	1077	ILE	3.0
1	A	1098	ILE	2.9
1	A	933	ASN	2.8
1	A	1094	LEU	2.8
1	A	1031	ALA	2.8
1	A	1095	LEU	2.7
1	A	999	PHE	2.6
1	A	1080	ARG	2.6
1	A	1034	ILE	2.5
1	A	875	THR	2.5
1	A	880	ARG	2.5
1	A	1047	PHE	2.4
1	A	924	LEU	2.3
1	A	845	PHE	2.2
1	A	876	HIS	2.1
1	A	877	SER	2.1
1	A	908	PRO	2.1
1	A	1081	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CME	A	1007	10/11	0.95	0.12	-	19,21,27,28	0
1	PTR	A	1059	16/17	0.75	0.31	-	41,46,55,56	0
1	CME	A	1024	10/11	0.94	0.17	-	34,36,51,53	0
1	CME	A	862	10/11	0.84	0.19	-	32,35,47,49	0
1	PTR	A	1054	16/17	0.87	0.20	-	41,42,43,44	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	A	3	5/5	0.80	0.28	4.12	73,74,74,75	0
2	SO4	A	6	5/5	0.93	0.27	3.89	80,80,80,80	0
2	SO4	A	1	5/5	0.85	0.23	1.94	65,66,68,68	0
3	XIN	A	1172	40/40	0.93	0.13	0.10	20,25,43,45	0
2	SO4	A	2	5/5	0.87	0.17	-	90,90,91,91	0
2	SO4	A	4	5/5	0.96	0.19	-	72,72,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.