



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:56 pm GMT

PDB ID : 5C7O
Title : Structure of human testis-specific glyceraldehyde-3-phosphate dehydrogenase holo form with NAD+
Authors : Betts, L.; Machius, M.; Danshina, P.; O'Brien, D.
Deposited on : 2015-06-24
Resolution : 1.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4385 (1.74-1.70)
Clashscore	112137	4841 (1.74-1.70)
Ramachandran outliers	110173	4771 (1.74-1.70)
Sidechain outliers	110143	4771 (1.74-1.70)
RSRZ outliers	101464	4426 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	O	335	<div> <div>2%</div> <div> <div></div> <div>96%</div> <div>.</div> </div> </div>
1	P	335	<div> <div>3%</div> <div> <div></div> <div>96%</div> <div>.</div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 11174 atoms, of which 5229 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glyceraldehyde-3-phosphate dehydrogenase, testis-specific.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	O	335	Total	C	H	N	O	S	0	7	0
			5190	1649	2585	450	490	16			
1	P	335	Total	C	H	N	O	S	0	8	0
			5205	1653	2592	452	492	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	408	LYS	-	expression tag	UNP O14556
P	408	LYS	-	expression tag	UNP O14556

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	O	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	P	1	Total	C	H	N	O	P	0	0
			70	21	26	7	14	2		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	O	1	Total	O	S	0	0
			5	4	1		
3	P	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	O	332	Total	O	0	0
			332	332		
4	P	297	Total	O	0	0
			297	297		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase, testis-specific

Chain O: 



- Molecule 1: Glyceraldehyde-3-phosphate dehydrogenase, testis-specific

Chain P: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	144.44Å 71.74Å 80.84Å 90.00° 123.01° 90.00°	Depositor
Resolution (Å)	31.71 – 1.73 31.71 – 1.73	Depositor EDS
% Data completeness (in resolution range)	94.0 (31.71-1.73) 91.1 (31.71-1.73)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.73Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.137 , 0.166 0.140 , 0.166	Depositor DCC
R_{free} test set	2016 reflections (2.96%)	DCC
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11174	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OCS, SO4, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	O	0.26	0/2670	0.51	2/3625 (0.1%)
1	P	0.26	0/2678	0.49	0/3636
All	All	0.26	0/5348	0.50	2/7261 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	O	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	150[A]	CYS	C-N-CA	-5.31	108.42	121.70
1	O	150[B]	CYS	C-N-CA	-5.31	108.42	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	O	149	GLN	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	O	2605	2585	2581	10	0
1	P	2613	2592	2586	7	0
2	O	44	26	23	1	0
2	P	44	26	23	0	0
3	O	5	0	0	0	0
3	P	5	0	0	1	0
4	O	332	0	0	8	4
4	P	297	0	0	4	4
All	All	5945	5229	5213	18	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:74:ARG:N	4:P:601:HOH:O	2.00	0.93
1:O:332[A]:GLU:OE1	4:O:601:HOH:O	2.08	0.71
1:O:162:GLY:O	4:O:602:HOH:O	2.09	0.70
1:P:217[A]:ASN:OD1	4:P:602:HOH:O	2.10	0.68
3:P:502:SO4:O4	4:P:603:HOH:O	2.13	0.67
1:P:97:LYS:NZ	4:P:604:HOH:O	2.19	0.65
1:O:159:ARG:NH1	4:O:609:HOH:O	2.32	0.62
1:O:335:LYS:NZ	4:O:607:HOH:O	2.30	0.58
1:P:101:VAL:O	1:P:143[A]:HIS:NE2	2.37	0.54
1:O:101:VAL:O	1:O:143[A]:HIS:NE2	2.35	0.53
1:O:407:ASP:O	4:O:603:HOH:O	2.19	0.52
1:P:201:MET:HE1	1:P:288:ALA:HB1	1.90	0.52
1:O:332[B]:GLU:OE2	4:O:604:HOH:O	2.20	0.47
1:P:180:SER:HA	1:P:218:ILE:HD11	1.97	0.46
1:O:299:LYS:HE3	4:O:695:HOH:O	2.17	0.45
1:O:234:LYS:HD2	4:O:645:HOH:O	2.16	0.44
1:P:209[A]:ASN:OD1	1:P:209[A]:ASN:N	2.52	0.42
1:O:84:GLY:HA3	2:O:501:NAD:O5B	2.19	0.42

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:O:821:HOH:O	4:P:832:HOH:O[2_556]	1.98	0.22
4:O:884:HOH:O	4:P:653:HOH:O[2_556]	2.08	0.12
4:O:850:HOH:O	4:P:792:HOH:O[2_556]	2.16	0.04
4:O:806:HOH:O	4:P:832:HOH:O[2_556]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	O	339/335 (101%)	326 (96%)	12 (4%)	1 (0%)	44	25
1	P	340/335 (102%)	326 (96%)	13 (4%)	1 (0%)	44	25
All	All	679/670 (101%)	652 (96%)	25 (4%)	2 (0%)	44	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	O	312	VAL
1	P	312	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	O	281/276 (102%)	279 (99%)	2 (1%)	87	80
1	P	282/276 (102%)	279 (99%)	3 (1%)	78	67
All	All	563/552 (102%)	558 (99%)	5 (1%)	82	74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	O	265	ARG
1	O	386	TYR
1	P	265	ARG
1	P	386	TYR
1	P	390	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCS	O	224	1	4,7,9	1.32	0	3,8,13	2.87	2 (66%)
1	OCS	P	224	1	4,7,9	1.36	0	3,8,13	2.44	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	O	224	1	-	0/3/6/9	0/0/0/0
1	OCS	P	224	1	-	0/3/6/9	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	O	224	OCS	OD3-SG-CB	2.91	109.65	105.33
1	P	224	OCS	OD1-SG-CB	3.59	110.67	105.33
1	O	224	OCS	OD1-SG-CB	3.66	110.78	105.33

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAD	O	501	-	41,48,48	5.17	13 (31%)	43,73,73	2.34	5 (11%)
3	SO4	O	502	-	4,4,4	0.18	0	6,6,6	0.08	0
2	NAD	P	501	-	41,48,48	5.19	13 (31%)	43,73,73	2.38	5 (11%)
3	SO4	P	502	-	4,4,4	0.18	0	6,6,6	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	O	501	-	-	0/22/62/62	0/5/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	O	502	-	-	0/0/0/0	0/0/0/0
2	NAD	P	501	-	-	0/22/62/62	0/5/5/5
3	SO4	P	502	-	-	0/0/0/0	0/0/0/0

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	501	NAD	C2B-C1B	-15.70	1.28	1.53
2	O	501	NAD	C2B-C1B	-15.48	1.29	1.53
2	O	501	NAD	C2D-C1D	-14.46	1.30	1.53
2	P	501	NAD	C2D-C1D	-14.39	1.30	1.53
2	P	501	NAD	O4D-C4D	-6.59	1.30	1.45
2	O	501	NAD	O4D-C4D	-6.40	1.30	1.45
2	P	501	NAD	O4B-C4B	-5.61	1.32	1.45
2	O	501	NAD	O4B-C4B	-5.41	1.32	1.45
2	O	501	NAD	O3B-C3B	-3.37	1.35	1.43
2	P	501	NAD	O3B-C3B	-3.28	1.35	1.43
2	O	501	NAD	O3D-C3D	-2.90	1.36	1.43
2	P	501	NAD	O3D-C3D	-2.82	1.36	1.43
2	P	501	NAD	C5A-C4A	-2.21	1.35	1.40
2	O	501	NAD	C5A-C4A	-2.17	1.35	1.40
2	P	501	NAD	C3N-C7N	2.04	1.53	1.50
2	O	501	NAD	C3N-C7N	2.10	1.53	1.50
2	P	501	NAD	C2A-N3A	3.11	1.37	1.32
2	O	501	NAD	C2A-N3A	3.11	1.37	1.32
2	P	501	NAD	O2B-C2B	3.41	1.50	1.43
2	O	501	NAD	O2B-C2B	3.58	1.51	1.43
2	P	501	NAD	C7N-N7N	7.40	1.47	1.33
2	O	501	NAD	C7N-N7N	7.41	1.47	1.33
2	O	501	NAD	O4B-C1B	14.48	1.61	1.41
2	P	501	NAD	O4B-C1B	14.60	1.61	1.41
2	P	501	NAD	O4D-C1D	15.11	1.62	1.41
2	O	501	NAD	O4D-C1D	15.14	1.62	1.41

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	501	NAD	N3A-C2A-N1A	-10.55	119.67	128.86
2	O	501	NAD	N3A-C2A-N1A	-10.48	119.73	128.86
2	P	501	NAD	N6A-C6A-N1A	-6.99	104.92	118.77
2	O	501	NAD	N6A-C6A-N1A	-6.60	105.69	118.77
2	O	501	NAD	C4B-O4B-C1B	-3.08	106.49	109.77

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	501	NAD	C4B-O4B-C1B	-2.80	106.78	109.77
2	O	501	NAD	C4D-O4D-C1D	-2.14	107.49	109.77
2	P	501	NAD	C4D-O4D-C1D	-2.03	107.61	109.77
2	O	501	NAD	C5A-C6A-N6A	6.66	134.05	120.47
2	P	501	NAD	C5A-C6A-N6A	6.93	134.59	120.47

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	O	501	NAD	1	0
3	P	502	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	O	334/335 (99%)	-0.31	6 (1%) 69 74	13, 21, 38, 60	0
1	P	334/335 (99%)	-0.23	10 (2%) 51 57	13, 22, 42, 61	0
All	All	668/670 (99%)	-0.27	16 (2%) 59 65	13, 21, 40, 61	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	O	407	ASP	4.5
1	O	142	ASN	3.1
1	O	150[A]	CYS	2.8
1	P	265	ARG	2.7
1	O	339	LYS	2.7
1	P	266	LYS	2.7
1	O	133	PHE	2.5
1	P	407	ASP	2.5
1	O	266	LYS	2.4
1	P	214	GLY	2.3
1	P	408	LYS	2.3
1	P	74	ARG	2.3
1	P	133	PHE	2.3
1	P	150[A]	CYS	2.2
1	P	215	SER	2.0
1	P	177	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy

less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	OCS	O	224	8/10	0.99	0.11	-	12,17,21,23	0
1	OCS	P	224	8/10	0.98	0.09	-	16,19,23,26	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	NAD	O	501	44/44	0.95	0.09	-0.04	15,25,32,38	70
2	NAD	P	501	44/44	0.94	0.09	-0.07	16,27,37,39	70
3	SO4	O	502	5/5	0.94	0.15	-	34,37,41,42	5
3	SO4	P	502	5/5	0.90	0.16	-	36,40,41,42	5

6.5 Other polymers [i](#)

There are no such residues in this entry.