



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 12:27 am GMT

PDB ID : 3C8N
Title : Crystal structure of apo-FGD1 from Mycobacterium tuberculosis
Authors : Bashiri, G.; Squire, C.J.; Moreland, N.J.; Baker, E.N.
Deposited on : 2008-02-12
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>12%</div> <div>• •</div> </div> </div>
1	B	356	<div> <div>%</div> <div> <div></div> <div>81%</div> <div>12%</div> <div>• 7%</div> </div> </div>
1	C	356	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>11%</div> <div>• 7%</div> </div> </div>
1	D	356	<div> <div>3%</div> <div> <div></div> <div>78%</div> <div>14%</div> <div>• 7%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 10741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable F420-dependent glucose-6-phosphate dehydrogenase FGD1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	343	Total	C	N	O	S	Se	0	1	0
			2667	1693	472	491	2	9			
1	B	332	Total	C	N	O	S	Se	0	2	0
			2577	1637	449	481	2	8			
1	C	332	Total	C	N	O	S	Se	0	2	0
			2567	1632	452	473	2	8			
1	D	330	Total	C	N	O	S	Se	0	2	0
			2559	1630	451	468	2	8			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	EXPRESSION TAG	UNP P96253
A	-18	GLY	-	EXPRESSION TAG	UNP P96253
A	-17	SER	-	EXPRESSION TAG	UNP P96253
A	-16	SER	-	EXPRESSION TAG	UNP P96253
A	-15	HIS	-	EXPRESSION TAG	UNP P96253
A	-14	HIS	-	EXPRESSION TAG	UNP P96253
A	-13	HIS	-	EXPRESSION TAG	UNP P96253
A	-12	HIS	-	EXPRESSION TAG	UNP P96253
A	-11	HIS	-	EXPRESSION TAG	UNP P96253
A	-10	HIS	-	EXPRESSION TAG	UNP P96253
A	-9	SER	-	EXPRESSION TAG	UNP P96253
A	-8	SER	-	EXPRESSION TAG	UNP P96253
A	-7	GLY	-	EXPRESSION TAG	UNP P96253
A	-6	LEU	-	EXPRESSION TAG	UNP P96253
A	-5	VAL	-	EXPRESSION TAG	UNP P96253
A	-4	PRO	-	EXPRESSION TAG	UNP P96253
A	-3	ARG	-	EXPRESSION TAG	UNP P96253
A	-2	GLY	-	EXPRESSION TAG	UNP P96253
A	-1	SER	-	EXPRESSION TAG	UNP P96253
A	0	HIS	-	EXPRESSION TAG	UNP P96253

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	243	MSE	LEU	ENGINEERED	UNP P96253
B	-19	MSE	-	EXPRESSION TAG	UNP P96253
B	-18	GLY	-	EXPRESSION TAG	UNP P96253
B	-17	SER	-	EXPRESSION TAG	UNP P96253
B	-16	SER	-	EXPRESSION TAG	UNP P96253
B	-15	HIS	-	EXPRESSION TAG	UNP P96253
B	-14	HIS	-	EXPRESSION TAG	UNP P96253
B	-13	HIS	-	EXPRESSION TAG	UNP P96253
B	-12	HIS	-	EXPRESSION TAG	UNP P96253
B	-11	HIS	-	EXPRESSION TAG	UNP P96253
B	-10	HIS	-	EXPRESSION TAG	UNP P96253
B	-9	SER	-	EXPRESSION TAG	UNP P96253
B	-8	SER	-	EXPRESSION TAG	UNP P96253
B	-7	GLY	-	EXPRESSION TAG	UNP P96253
B	-6	LEU	-	EXPRESSION TAG	UNP P96253
B	-5	VAL	-	EXPRESSION TAG	UNP P96253
B	-4	PRO	-	EXPRESSION TAG	UNP P96253
B	-3	ARG	-	EXPRESSION TAG	UNP P96253
B	-2	GLY	-	EXPRESSION TAG	UNP P96253
B	-1	SER	-	EXPRESSION TAG	UNP P96253
B	0	HIS	-	EXPRESSION TAG	UNP P96253
B	243	MSE	LEU	ENGINEERED	UNP P96253
C	-19	MSE	-	EXPRESSION TAG	UNP P96253
C	-18	GLY	-	EXPRESSION TAG	UNP P96253
C	-17	SER	-	EXPRESSION TAG	UNP P96253
C	-16	SER	-	EXPRESSION TAG	UNP P96253
C	-15	HIS	-	EXPRESSION TAG	UNP P96253
C	-14	HIS	-	EXPRESSION TAG	UNP P96253
C	-13	HIS	-	EXPRESSION TAG	UNP P96253
C	-12	HIS	-	EXPRESSION TAG	UNP P96253
C	-11	HIS	-	EXPRESSION TAG	UNP P96253
C	-10	HIS	-	EXPRESSION TAG	UNP P96253
C	-9	SER	-	EXPRESSION TAG	UNP P96253
C	-8	SER	-	EXPRESSION TAG	UNP P96253
C	-7	GLY	-	EXPRESSION TAG	UNP P96253
C	-6	LEU	-	EXPRESSION TAG	UNP P96253
C	-5	VAL	-	EXPRESSION TAG	UNP P96253
C	-4	PRO	-	EXPRESSION TAG	UNP P96253
C	-3	ARG	-	EXPRESSION TAG	UNP P96253
C	-2	GLY	-	EXPRESSION TAG	UNP P96253
C	-1	SER	-	EXPRESSION TAG	UNP P96253
C	0	HIS	-	EXPRESSION TAG	UNP P96253

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	243	MSE	LEU	ENGINEERED	UNP P96253
D	-19	MSE	-	EXPRESSION TAG	UNP P96253
D	-18	GLY	-	EXPRESSION TAG	UNP P96253
D	-17	SER	-	EXPRESSION TAG	UNP P96253
D	-16	SER	-	EXPRESSION TAG	UNP P96253
D	-15	HIS	-	EXPRESSION TAG	UNP P96253
D	-14	HIS	-	EXPRESSION TAG	UNP P96253
D	-13	HIS	-	EXPRESSION TAG	UNP P96253
D	-12	HIS	-	EXPRESSION TAG	UNP P96253
D	-11	HIS	-	EXPRESSION TAG	UNP P96253
D	-10	HIS	-	EXPRESSION TAG	UNP P96253
D	-9	SER	-	EXPRESSION TAG	UNP P96253
D	-8	SER	-	EXPRESSION TAG	UNP P96253
D	-7	GLY	-	EXPRESSION TAG	UNP P96253
D	-6	LEU	-	EXPRESSION TAG	UNP P96253
D	-5	VAL	-	EXPRESSION TAG	UNP P96253
D	-4	PRO	-	EXPRESSION TAG	UNP P96253
D	-3	ARG	-	EXPRESSION TAG	UNP P96253
D	-2	GLY	-	EXPRESSION TAG	UNP P96253
D	-1	SER	-	EXPRESSION TAG	UNP P96253
D	0	HIS	-	EXPRESSION TAG	UNP P96253
D	243	MSE	LEU	ENGINEERED	UNP P96253


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	124	Total O 124 124	0	0
2	B	111	Total O 111 111	0	0
2	C	78	Total O 78 78	0	0
2	D	58	Total O 58 58	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Probable F420-dependent glucose-6-phosphate dehydrogenase FGD1




- Molecule 1: Probable F420-dependent glucose-6-phosphate dehydrogenase FGD1




- Molecule 1: Probable F420-dependent glucose-6-phosphate dehydrogenase FGD1



- Molecule 1: Probable F420-dependent glucose-6-phosphate dehydrogenase FGD1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	80.95Å 89.13Å 90.86Å 90.00° 91.64° 90.00°	Depositor
Resolution (Å)	40.10 – 1.90 40.09 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.10-1.90) 91.7 (40.09-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.3.0008	Depositor
R, R_{free}	0.212 , 0.255 0.213 , 0.209	Depositor DCC
R_{free} test set	5067 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 27.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.030 for -h,l,k 0.035 for -h,-l,-k 0.099 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10741	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/2730	0.81	1/3689 (0.0%)
1	B	0.80	0/2639	0.79	1/3569 (0.0%)
1	C	0.72	0/2633	0.77	3/3567 (0.1%)
1	D	0.68	0/2623	0.72	2/3548 (0.1%)
All	All	0.75	0/10625	0.78	7/14373 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	140	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	C	140	ARG	NE-CZ-NH1	6.23	123.41	120.30
1	C	100	ARG	NE-CZ-NH2	-6.23	117.19	120.30
1	D	100	ARG	NE-CZ-NH1	5.59	123.10	120.30
1	A	100	ARG	NE-CZ-NH2	-5.38	117.61	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2667	0	2582	35	0
1	B	2577	0	2507	34	0
1	C	2567	0	2470	26	0
1	D	2559	0	2477	30	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	124	0	0	4	0
2	B	111	0	0	10	0
2	C	78	0	0	4	0
2	D	58	0	0	5	0
All	All	10741	0	10036	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:150:PHE:O	1:A:156:ARG:HG3	1.51	1.09
1:D:55:LEU:HD13	2:D:378:HOH:O	1.68	0.90
1:B:285:ILE:HG23	1:B:296:LYS:HD3	1.54	0.89
1:B:144:SER:HB3	2:B:358:HOH:O	1.74	0.88
1:B:269:GLU:HB3	2:B:364:HOH:O	1.76	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/356 (95%)	329 (98%)	8 (2%)	0	100	100
1	B	330/356 (93%)	324 (98%)	6 (2%)	0	100	100
1	C	332/356 (93%)	323 (97%)	9 (3%)	0	100	100
1	D	326/356 (92%)	320 (98%)	6 (2%)	0	100	100
All	All	1325/1424 (93%)	1296 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/270 (99%)	263 (98%)	5 (2%)	62	57
1	B	260/270 (96%)	256 (98%)	4 (2%)	70	67
1	C	255/270 (94%)	251 (98%)	4 (2%)	68	65
1	D	255/270 (94%)	246 (96%)	9 (4%)	41	30
All	All	1038/1080 (96%)	1016 (98%)	22 (2%)	59	53

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	208	MSE
1	C	334	ARG
1	D	264	ASP
1	C	221	SER
1	C	283	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-12	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/356 (93%)	0.23	9 (2%) 55 59	12, 23, 35, 43	0
1	B	323/356 (90%)	0.01	5 (1%) 74 77	12, 23, 37, 48	0
1	C	324/356 (91%)	0.42	11 (3%) 46 49	18, 26, 39, 45	0
1	D	322/356 (90%)	0.46	10 (3%) 49 53	19, 28, 37, 42	0
All	All	1303/1424 (91%)	0.28	35 (2%) 55 59	12, 25, 37, 48	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	144	SER	6.8
1	B	46	HIS	4.1
1	C	256	ALA	4.0
1	C	260	HIS	3.9
1	D	44	TRP	3.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.