



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:34 pm GMT

PDB ID : 1C9D
Title : CRYSTAL STRUCTURE OF THE COMPLEX OF BACTERIAL TRYPTOPHAN SYNTHASE WITH THE TRANSITION STATE ANALOGUE INHIBITOR 4-(2-HYDROXY-4-FLUOROPHENYLTHIO)-BUTYLPHOSPHONIC ACID
Authors : Lolis, E.; Sachpatzidis, A.
Deposited on : 1999-08-02
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	268	<div> <div>2%</div> <div> <div></div> <div>72%</div> <div>22%</div> <div>5%</div> </div> </div>
2	B	397	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>22%</div> <div></div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PLP	B	900	-	X	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPTOPHAN SYNTHASE (ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	264	Total	C	N	O	S	0	0	0
			1992	1264	348	373	7			


- Molecule 2 is a protein called TRYPTOPHAN SYNTHASE (BETA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	387	Total	C	N	O	S	0	0	0
			2934	1843	516	556	19			

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		

- Molecule 4 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 5 is 4-(2-HYDROXY-4-FLUOROPHENYLTHIO)-BUTYLPHOSPHONIC ACID (three-letter code: HF1) (formula: C₁₀H₁₂FO₄PS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	F	O	P	S	0	0
			17	10	1	4	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	75	Total	O	0	0
			75	75		
6	B	113	Total	O	0	0
			113	113		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRYPTOPHAN SYNTHASE (ALPHA CHAIN)



• Molecule 2: TRYPTOPHAN SYNTHASE (BETA CHAIN)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	184.22Å 60.52Å 67.79Å 90.00° 94.43° 90.00°	Depositor
Resolution (Å)	30.00 – 2.30 39.42 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.9 (30.00-2.30) 94.6 (39.42-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 2.29Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.211 , 0.269 0.204 , 0.260	Depositor DCC
R_{free} test set	3162 reflections (11.13%)	DCC
Wilson B-factor (Å ²)	14.6	Xtriage
Anisotropy	0.705	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5147	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HF1, NA, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2031	0.68	0/2757
2	B	0.45	0/2992	0.77	2/4042 (0.0%)
All	All	0.44	0/5023	0.73	2/6799 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	49	LEU	CA-CB-CG	5.65	128.29	115.30
2	B	207	MET	CB-CG-SD	-5.15	96.94	112.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	298	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1992	0	1995	41	0
2	B	2934	0	2902	52	0
3	B	1	0	0	0	0
4	B	15	0	7	0	0
5	A	17	0	10	1	0
6	A	75	0	0	1	0
6	B	113	0	0	5	0
All	All	5147	0	4914	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:VAL:HG11	2:B:175:ARG:HG2	1.53	0.91
2:B:202:ARG:HD3	2:B:312:GLN:HE22	1.50	0.76
1:A:133:VAL:H	2:B:19:GLN:HE22	1.36	0.74
2:B:112:ALA:O	2:B:142:GLN:HG2	1.87	0.74
2:B:89:ASN:HB2	2:B:205:GLN:HE22	1.53	0.72
1:A:141:GLN:HB3	1:A:145:ARG:NH1	2.04	0.71
2:B:327:ILE:HG23	2:B:331:GLU:HB3	1.73	0.71
2:B:142:GLN:HG3	6:B:1075:HOH:O	1.90	0.70
1:A:89:ARG:HH21	1:A:89:ARG:HG3	1.56	0.69
2:B:89:ASN:HD22	2:B:205:GLN:NE2	1.93	0.66
1:A:211:GLY:O	1:A:212:PHE:HB2	1.96	0.65
1:A:89:ARG:HD2	1:A:122:GLY:HA3	1.77	0.64
2:B:77:ARG:HG2	2:B:80:LEU:HD22	1.79	0.64
1:A:85:LEU:HB3	1:A:121:VAL:CG1	2.28	0.64
2:B:4:LEU:HD11	2:B:30:GLU:HG2	1.81	0.62
1:A:140:ARG:HD2	1:A:169:TYR:HB3	1.82	0.62
2:B:77:ARG:CG	2:B:80:LEU:HD22	2.30	0.62
2:B:216:ILE:HG21	2:B:224:PRO:HD3	1.83	0.60
1:A:53:PRO:HA	1:A:68:ASN:HD22	1.65	0.60
1:A:85:LEU:HB3	1:A:121:VAL:HG11	1.82	0.60
1:A:247:SER:HB2	1:A:250:GLN:HB2	1.83	0.60
2:B:185:HIS:HD2	6:B:1093:HOH:O	1.84	0.60
2:B:288:GLN:HE21	2:B:292:GLY:HA2	1.67	0.60
2:B:89:ASN:HB2	2:B:205:GLN:NE2	2.17	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:16:TYR:O	2:B:281:GLY:HA2	2.03	0.58
2:B:202:ARG:HH21	2:B:312:GLN:HE22	1.50	0.58
2:B:134:MET:O	2:B:158:PRO:HA	2.04	0.57
2:B:108:ALA:HB3	2:B:132:ILE:HG12	1.86	0.57
2:B:76:LYS:HE3	2:B:215:GLN:HE22	1.70	0.56
1:A:234:GLY:HA3	5:A:270:HF1:H2	1.88	0.56
2:B:382:LYS:NZ	2:B:382:LYS:HB3	2.20	0.56
1:A:104:ASN:HD21	2:B:288:GLN:NE2	2.04	0.56
2:B:195:HIS:HD2	6:B:1006:HOH:O	1.90	0.55
2:B:89:ASN:HD22	2:B:205:GLN:HE22	1.53	0.55
1:A:141:GLN:HB3	1:A:145:ARG:HH11	1.72	0.55
2:B:360:LYS:O	2:B:364:GLU:HB2	2.06	0.54
1:A:26:GLY:HA3	1:A:76:VAL:HG21	1.90	0.53
2:B:106:ILE:HD11	2:B:128:LEU:HD13	1.90	0.53
1:A:19:PHE:HE2	1:A:259:VAL:HB	1.75	0.52
1:A:135:GLU:HG2	2:B:16:TYR:CE2	2.45	0.52
1:A:162:LEU:HD22	1:A:166:VAL:HG23	1.92	0.52
2:B:202:ARG:HH21	2:B:312:GLN:NE2	2.08	0.52
1:A:255:LEU:O	1:A:259:VAL:HG13	2.10	0.51
1:A:89:ARG:NH2	1:A:89:ARG:HG3	2.21	0.51
2:B:87:LYS:HE2	2:B:114:GLN:HB3	1.93	0.51
2:B:135:GLY:HA2	2:B:159:VAL:HG22	1.93	0.50
2:B:271:LEU:HB2	2:B:309:VAL:HG11	1.93	0.49
2:B:129:LYS:HE3	2:B:129:LYS:HB2	1.67	0.48
1:A:19:PHE:CE1	1:A:262:MET:HG2	2.49	0.48
2:B:106:ILE:CD1	2:B:128:LEU:HD13	2.43	0.48
2:B:329:ASP:O	2:B:333:LEU:HD13	2.14	0.48
1:A:179:ARG:HG3	1:A:189:GLY:HA2	1.95	0.47
2:B:21:LEU:HD21	2:B:178:SER:HA	1.97	0.47
2:B:76:LYS:HD2	2:B:76:LYS:HA	1.67	0.47
1:A:145:ARG:HD2	6:A:1063:HOH:O	2.13	0.47
2:B:81:LEU:HD13	2:B:88:THR:HB	1.97	0.47
1:A:249:LYS:HE3	1:A:249:LYS:HB2	1.68	0.46
1:A:163:LEU:HD21	1:A:196:LEU:HD22	1.97	0.46
2:B:370:GLN:HG2	6:B:1144:HOH:O	2.17	0.45
1:A:46:ASP:OD1	1:A:263:LYS:HE3	2.17	0.44
1:A:137:ALA:HB3	1:A:138:PRO:CD	2.47	0.44
1:A:55:SER:HB3	2:B:293:GLN:HG3	1.98	0.44
1:A:89:ARG:NH1	1:A:95:ILE:O	2.46	0.44
2:B:211:GLU:O	2:B:215:GLN:HB2	2.18	0.44
1:A:263:LYS:HE2	1:A:266:SER:OG	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:MET:O	1:A:255:LEU:HB2	2.18	0.43
2:B:234:GLY:O	2:B:313:HIS:HE1	2.02	0.43
2:B:106:ILE:HD13	2:B:128:LEU:HB3	1.99	0.43
1:A:185:ALA:HB1	1:A:215:SER:HB3	2.00	0.43
1:A:11:LEU:HD12	1:A:11:LEU:HA	1.90	0.43
2:B:115:HIS:CE1	2:B:189:GLY:HA2	2.54	0.43
2:B:87:LYS:HG2	6:B:1048:HOH:O	2.19	0.42
2:B:34:ARG:HG3	2:B:34:ARG:HH11	1.85	0.42
1:A:44:GLY:O	1:A:263:LYS:HD3	2.19	0.42
2:B:76:LYS:HD3	2:B:212:THR:OG1	2.19	0.42
1:A:89:ARG:NH2	1:A:92:HIS:O	2.52	0.41
1:A:91:LYS:HB3	1:A:92:HIS:CD2	2.56	0.41
2:B:205:GLN:HA	2:B:205:GLN:HE21	1.86	0.41
1:A:20:VAL:HG22	1:A:47:ALA:HB3	2.02	0.41
1:A:52:VAL:HG21	1:A:101:MET:CE	2.50	0.41
2:B:147:PHE:O	2:B:151:LEU:HB2	2.21	0.41
1:A:105:LEU:HA	1:A:105:LEU:HD12	1.92	0.41
2:B:70:ARG:NH1	2:B:367:GLU:HG3	2.36	0.41
1:A:132:PRO:HD3	2:B:17:VAL:O	2.21	0.40
1:A:177:LEU:HA	1:A:177:LEU:HD12	1.99	0.40
2:B:195:HIS:CD2	2:B:196:PRO:HA	2.57	0.40
2:B:201:VAL:O	2:B:205:GLN:HB2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/268 (97%)	249 (96%)	9 (4%)	2 (1%)	22	26
2	B	385/397 (97%)	371 (96%)	13 (3%)	1 (0%)	44	55
All	All	645/665 (97%)	620 (96%)	22 (3%)	3 (0%)	32	39

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	PHE
1	A	188	ARG
2	B	181	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/208 (99%)	176 (85%)	30 (15%)	3	3
2	B	303/311 (97%)	259 (86%)	44 (14%)	4	3
All	All	509/519 (98%)	435 (86%)	74 (14%)	4	3

All (74) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	10	GLN
1	A	11	LEU
1	A	32	GLN
1	A	40	LEU
1	A	48	LEU
1	A	50	LEU
1	A	80	GLN
1	A	85	LEU
1	A	87	LEU
1	A	99	LEU
1	A	101	MET
1	A	105	LEU
1	A	106	VAL
1	A	112	ASP
1	A	117	ARG
1	A	120	GLN
1	A	141	GLN
1	A	144	LEU
1	A	162	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	165	GLN
1	A	187	ASN
1	A	209	LEU
1	A	220	VAL
1	A	249	LYS
1	A	250	GLN
1	A	255	LEU
1	A	259	VAL
1	A	262	MET
1	A	263	LYS
2	B	15	MET
2	B	30	GLU
2	B	36	GLN
2	B	40	GLU
2	B	42	GLN
2	B	49	LEU
2	B	50	LYS
2	B	55	ARG
2	B	61	LYS
2	B	63	GLN
2	B	65	ILE
2	B	77	ARG
2	B	78	GLU
2	B	80	LEU
2	B	81	LEU
2	B	90	GLN
2	B	101	MET
2	B	103	LYS
2	B	105	GLU
2	B	129	LYS
2	B	141	ARG
2	B	143	SER
2	B	152	MET
2	B	166	LEU
2	B	167	LYS
2	B	203	GLU
2	B	205	GLN
2	B	207	MET
2	B	236	ASN
2	B	266	GLU
2	B	272	LYS
2	B	280	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	287	MET
2	B	293	GLN
2	B	295	GLU
2	B	305	ASP
2	B	341	ARG
2	B	347	PRO
2	B	355	LEU
2	B	359	LEU
2	B	367	GLU
2	B	368	LYS
2	B	370	GLN
2	B	372	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	68	ASN
1	A	80	GLN
1	A	120	GLN
1	A	165	GLN
1	A	187	ASN
1	A	244	ASN
2	B	19	GLN
2	B	27	GLN
2	B	64	ASN
2	B	82	HIS
2	B	90	GLN
2	B	94	GLN
2	B	171	ASN
2	B	185	HIS
2	B	195	HIS
2	B	205	GLN
2	B	215	GLN
2	B	236	ASN
2	B	288	GLN
2	B	312	GLN
2	B	313	HIS
2	B	365	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	HF1	A	270	-	16,17,17	4.48	8 (50%)	18,23,23	1.70	4 (22%)
4	PLP	B	900	2	15,15,16	2.00	6 (40%)	20,22,23	4.22	9 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	HF1	A	270	-	-	0/5/9/9	0/1/1/1
4	PLP	B	900	2	-	0/6/6/8	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	270	HF1	P1-O2	-9.66	1.40	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	270	HF1	P1-O1	-9.49	1.40	1.54
4	B	900	PLP	C4A-C4	-4.36	1.42	1.51
5	A	270	HF1	P1-O3	-2.97	1.42	1.48
4	B	900	PLP	P-O4P	-2.75	1.51	1.60
4	B	900	PLP	C3-C2	-2.56	1.39	1.40
4	B	900	PLP	O4P-C5A	-2.43	1.35	1.44
4	B	900	PLP	P-O3P	-2.19	1.45	1.54
5	A	270	HF1	C5-C6	2.09	1.42	1.38
4	B	900	PLP	C5A-C5	2.28	1.57	1.50
5	A	270	HF1	OH-C7	3.22	1.42	1.36
5	A	270	HF1	C5-C10	3.29	1.43	1.37
5	A	270	HF1	P1-C1	6.40	1.85	1.76
5	A	270	HF1	C7-C8	7.21	1.45	1.40

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	900	PLP	C4A-C4-C3	-8.76	105.41	120.54
5	A	270	HF1	C9-C8-S1	-3.79	116.93	122.77
4	B	900	PLP	C5A-C5-C6	-3.31	113.65	119.33
5	A	270	HF1	C4-C2-C1	-3.17	119.15	125.77
4	B	900	PLP	O2P-P-O1P	-3.11	98.31	110.50
4	B	900	PLP	C3-C2-N1	-2.92	116.92	120.75
4	B	900	PLP	O4P-C5A-C5	-2.83	103.63	109.32
4	B	900	PLP	O3P-P-O4P	-2.57	99.89	106.73
5	A	270	HF1	C5-C10-C9	-2.42	120.12	123.29
4	B	900	PLP	O3P-P-O1P	2.35	119.70	110.50
5	A	270	HF1	C8-C9-C10	3.10	120.44	116.59
4	B	900	PLP	O4P-P-O1P	6.43	124.51	106.47
4	B	900	PLP	C4A-C4-C5	13.37	134.36	120.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	270	HF1	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	264/268 (98%)	-0.12	6 (2%) 61 67	4, 16, 44, 75	0
2	B	387/397 (97%)	-0.29	3 (0%) 86 89	2, 10, 28, 51	0
All	All	651/665 (97%)	-0.22	9 (1%) 75 80	2, 13, 36, 75	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	186	GLU	2.8
1	A	109	ASN	2.7
1	A	185	ALA	2.5
1	A	267	ARG	2.4
2	B	280	PHE	2.2
1	A	189	GLY	2.2
2	B	147	PHE	2.2
1	A	249	LYS	2.1
2	B	141	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	PLP	B	900	15/16	0.95	0.17	1.51	7,13,18,18	0
5	HF1	A	270	17/17	0.95	0.13	-0.17	9,11,15,20	0
3	NA	B	1191	1/1	0.91	0.08	-1.64	7,7,7,7	0

6.5 Other polymers [i](#)

There are no such residues in this entry.