



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 19, 2017 – 09:37 PM EDT

PDB ID : 3C96  
Title : Crystal structure of the flavin-containing monooxygenase phzS from *Pseudomonas aeruginosa*. Northeast Structural Genomics Consortium target PaR240  
Authors : Vorobiev, S.M.; Chen, Y.; Seetharaman, J.; Wang, D.; Mao, L.; Xiao, R.; Acton, T.B.; Montelione, G.T.; Tong, L.; Hunt, J.F.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : unknown  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	<div> <div>2%</div> <div>76%</div> <div>16%</div> <div>7%</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flavin-containing monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	381	2888	1813	537	529	9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	403	LEU	-	EXPRESSION TAG	UNP Q9HWG9
A	404	GLU	-	EXPRESSION TAG	UNP Q9HWG9
A	405	HIS	-	EXPRESSION TAG	UNP Q9HWG9
A	406	HIS	-	EXPRESSION TAG	UNP Q9HWG9
A	407	HIS	-	EXPRESSION TAG	UNP Q9HWG9
A	408	HIS	-	EXPRESSION TAG	UNP Q9HWG9
A	409	HIS	-	EXPRESSION TAG	UNP Q9HWG9
A	410	HIS	-	EXPRESSION TAG	UNP Q9HWG9

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	278	Total	O	0	0
			278	278		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Flavin-containing monooxygenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.15Å 65.26Å 88.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.65 – 1.90 40.65 – 1.89	Depositor EDS
% Data completeness (in resolution range)	94.7 (40.65-1.90) 96.8 (40.65-1.89)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.77 (at 1.89Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.183 , 0.197 0.190 , 0.208	Depositor DCC
$R_{free}$ test set	1440 reflections (4.93%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.9	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 52.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.016 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3219	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/2948	0.62	1/4007 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	166	GLY	N-CA-C	5.13	125.92	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2888	0	2854	46	0
2	A	53	0	30	1	0
3	A	278	0	0	5	0
All	All	3219	0	2884	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:HIS:HD2	1:A:186:GLY:H	1.23	0.85
1:A:81:GLN:O	1:A:402:ARG:HD2	1.90	0.72
1:A:378:ARG:NH1	1:A:378:ARG:HB3	2.04	0.72
1:A:270:ASP:O	1:A:402:ARG:HD3	1.93	0.69
1:A:278:ASP:OD1	1:A:282:ARG:HD3	1.93	0.68
1:A:184:HIS:CD2	1:A:186:GLY:H	2.11	0.66
1:A:8:LEU:HD11	1:A:34:LEU:HG	1.79	0.65
1:A:76:LEU:HD13	1:A:91:ARG:HH21	1.64	0.62
1:A:378:ARG:HB3	1:A:378:ARG:HH11	1.64	0.61
1:A:46:GLY:H	1:A:234:ASN:ND2	1.99	0.60
1:A:367:GLU:O	1:A:371:GLU:HG3	2.01	0.60
1:A:296:PRO:HG2	1:A:363:LEU:HD13	1.85	0.59
1:A:181:PRO:HD2	3:A:736:HOH:O	2.05	0.56
1:A:55:GLU:O	1:A:59:GLU:HG3	2.05	0.56
1:A:4:PRO:HB2	1:A:29:GLY:HA2	1.87	0.56
1:A:216:LEU:HD22	1:A:276:ILE:HD12	1.89	0.53
1:A:77:ARG:HB2	1:A:206:ILE:HD13	1.91	0.52
1:A:368:ARG:HD3	3:A:765:HOH:O	2.10	0.51
1:A:321:ASN:HD21	1:A:365:ASN:ND2	2.10	0.49
1:A:283:ASN:HD22	1:A:285:LEU:N	2.11	0.48
1:A:34:LEU:HD11	1:A:156:LEU:HD12	1.95	0.48
1:A:119:GLU:HG3	3:A:749:HOH:O	2.14	0.47
1:A:252:ASP:HB3	1:A:255:ARG:HB2	1.96	0.47
1:A:189:MET:HE1	1:A:317:PRO:HG2	1.95	0.47
1:A:355:ARG:HB3	1:A:356:PRO:HD3	1.97	0.47
1:A:133:GLU:HG2	1:A:134:ARG:HG3	1.97	0.47
1:A:206:ILE:HA	1:A:206:ILE:HD13	1.80	0.46
1:A:47:ILE:HD13	1:A:106:ARG:HG3	1.99	0.45
1:A:138:ARG:NH1	1:A:139:ASP:OD2	2.49	0.45
1:A:402:ARG:HG3	3:A:640:HOH:O	2.15	0.45
1:A:74:HIS:CE1	1:A:201:ASP:OD2	2.71	0.43
1:A:378:ARG:HA	1:A:379:PRO:HD3	1.78	0.43
1:A:402:ARG:HG3	1:A:402:ARG:H	1.61	0.43
1:A:278:ASP:OD1	1:A:282:ARG:CD	2.64	0.43
1:A:165:ASP:OD1	1:A:165:ASP:N	2.52	0.43
1:A:291:MET:HG2	1:A:316:TYR:CD1	2.53	0.42
1:A:114:LEU:HD22	1:A:118:ARG:HD2	2.00	0.42
1:A:76:LEU:HD23	1:A:76:LEU:C	2.39	0.42
1:A:258:ARG:NH2	3:A:806:HOH:O	2.52	0.42
1:A:82:SER:HA	1:A:402:ARG:HD2	2.02	0.41
1:A:268:ASP:O	1:A:403:LEU:HB2	2.21	0.41
1:A:43:LEU:HB2	2:A:501:FAD:HM72	2.02	0.41

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:TYR:HA	1:A:220:PRO:HD3	1.92	0.41
1:A:184:HIS:HD2	1:A:186:GLY:N	2.04	0.41
1:A:50:GLN:HE21	1:A:328:LEU:HD11	1.84	0.41
1:A:8:LEU:HD21	1:A:34:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	377/410 (92%)	366 (97%)	10 (3%)	1 (0%)	44 34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	283/315 (90%)	271 (96%)	12 (4%)	34 23

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	GLU
1	A	60	LEU
1	A	74	HIS
1	A	114	LEU
1	A	143	LEU
1	A	180	ARG
1	A	205	MET
1	A	206	ILE
1	A	287	LEU
1	A	350	TYR
1	A	363	LEU
1	A	378	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
1	A	74	HIS
1	A	81	GLN
1	A	101	GLN
1	A	168	HIS
1	A	184	HIS
1	A	234	ASN
1	A	254	ASN
1	A	283	ASN
1	A	359	ASN
1	A	365	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	A	501	-	51,58,58	1.63	11 (21%)	54,89,89	2.05	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	501	-	-	0/28/50/50	0/6/6/6

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	FAD	O2B-C2B	-4.33	1.33	1.43
2	A	501	FAD	C5X-N5	2.30	1.38	1.35
2	A	501	FAD	C2A-N1A	2.46	1.38	1.33
2	A	501	FAD	O4B-C1B	2.84	1.45	1.41
2	A	501	FAD	C2B-C1B	2.86	1.58	1.53
2	A	501	FAD	C4-N3	2.90	1.38	1.33
2	A	501	FAD	C1'-N10	3.02	1.51	1.48
2	A	501	FAD	C5'-C4'	3.41	1.56	1.51
2	A	501	FAD	C4X-N5	3.58	1.38	1.33
2	A	501	FAD	C10-N1	3.67	1.38	1.33
2	A	501	FAD	C2A-N3A	4.23	1.39	1.32

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	N3A-C2A-N1A	-10.32	119.87	128.86
2	A	501	FAD	C4X-C4-N3	-2.64	119.73	123.48
2	A	501	FAD	C1B-N9A-C4A	-2.42	122.45	126.64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	FAD	C5X-C9A-N10	2.46	119.49	117.66
2	A	501	FAD	O2B-C2B-C3B	3.04	121.58	111.83
2	A	501	FAD	C1'-N10-C9A	3.30	121.37	118.35
2	A	501	FAD	C4X-N5-C5X	3.33	120.28	116.76
2	A	501	FAD	C4-N3-C2	6.01	120.42	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	FAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	381/410 (92%)	0.00	10 (2%) 56 60	7, 15, 26, 31	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	401	PRO	3.1
1	A	400	ARG	3.0
1	A	134	ARG	3.0
1	A	247	LEU	2.6
1	A	149	GLY	2.4
1	A	403	LEU	2.3
1	A	250	GLU	2.3
1	A	138	ARG	2.1
1	A	402	ARG	2.1
1	A	123	GLN	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FAD	A	501	53/53	0.96	0.12	1.23	13,16,19,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.