



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2018 – 06:12 PM EST

PDB ID : 6CBP
Title : Crystal structure of the single chain variable fragment of the DH270.6 bnAb
in complex with the Man9-V3 glycopeptide
Authors : Fera, D.; Harrison, S.C.
Deposited on : 2018-02-03
Resolution : 2.17 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030736
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030736


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5526 (2.20-2.16)
Clashscore	112137	6386 (2.20-2.16)
Ramachandran outliers	110173	6282 (2.20-2.16)
Sidechain outliers	110143	6282 (2.20-2.16)
RSRZ outliers	101464	5562 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	P	30	<div> <div>7%</div> <div>73%</div> <div>23%</div> <div>.</div> </div>
2	A	256	<div> <div>%</div> <div>76%</div> <div>15%</div> <div>8%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MAN	A	305	-	-	-	X
5	MAN	A	311	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Man9-V3 glycopeptide.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	P	30	Total	C	N	O	S	0	0	0
			233	136	50	45	2			


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
P	305	PRO	-	linker	UNP Q74448

- Molecule 2 is a protein called DH270.6 single chain variable fragment.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	235	Total	C	N	O	S	0	0	0
			1786	1119	303	352	12			

- Molecule 3 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	P	1	Total	C	N	O	0	0
			14	8	1	5		
3	P	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is BETA-D-MANNOSE (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	P	1	Total	C	O	0	0
			11	6	5		
4	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	P	14	Total	O	0	0
			14	14		
6	A	81	Total	O	0	0
			81	81		

4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	68.00Å 68.00Å 128.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	60.14 – 2.17 60.14 – 2.17	Depositor EDS
% Data completeness (in resolution range)	95.1 (60.14-2.17) 95.1 (60.14-2.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 2.16Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.221 , 0.278 0.226 , 0.273	Depositor DCC
R_{free} test set	812 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.290	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 42.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2280	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	P	0.28	0/235	0.54	0/317
2	A	0.27	0/1832	0.48	0/2492
All	All	0.27	0/2067	0.49	0/2809

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	P	233	0	223	5	0
2	A	1786	0	1685	18	0
3	A	28	0	24	1	0
3	P	28	0	24	0	0
4	A	11	0	9	1	0
4	P	11	0	10	1	0
5	A	88	0	75	5	0
6	A	81	0	0	2	0
6	P	14	0	0	1	0
All	All	2280	0	2050	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:307:MAN:O6	5:A:310:MAN:C1	1.99	1.09
2:A:69:THR:OG1	2:A:84:ARG:NH1	2.03	0.91
2:A:205:ARG:NH2	2:A:226:ASP:OD2	2.11	0.84
4:A:303:BMA:O6	5:A:307:MAN:H5	1.91	0.70
1:P:330:HIS:CE1	1:P:332:ASN:HD21	2.10	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	28/30 (93%)	27 (96%)	1 (4%)	0	100	100
2	A	231/256 (90%)	220 (95%)	9 (4%)	2 (1%)	20	17
All	All	259/286 (91%)	247 (95%)	10 (4%)	2 (1%)	22	19

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	65	GLN
2	A	174	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	P	26/26 (100%)	23 (88%)	3 (12%)	6	5
2	A	193/204 (95%)	184 (95%)	9 (5%)	30	35
All	All	219/230 (95%)	207 (94%)	12 (6%)	25	27

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	67	ARG
2	A	72	ARG
2	A	170	ASP
2	A	63	ASN
2	A	123	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	P	330	HIS
1	P	332	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	301	3	14,14,15	0.39	0	15,19,21	0.58	0
3	NAG	A	302	3,4	14,14,15	0.34	0	15,19,21	0.77	0
4	BMA	A	303	3,5	11,11,12	0.33	0	13,15,17	0.81	1 (7%)
5	MAN	A	304	5,4	11,11,12	0.23	0	13,15,17	0.69	0
5	MAN	A	305	5	11,11,12	0.22	0	13,15,17	0.89	0
5	MAN	A	306	5	11,11,12	0.25	0	13,15,17	0.60	0
5	MAN	A	307	5	11,11,12	0.26	0	13,15,17	0.57	0
5	MAN	A	308	5	11,11,12	0.23	0	13,15,17	0.81	0
5	MAN	A	309	5	11,11,12	0.26	0	13,15,17	1.05	1 (7%)
5	MAN	A	310	5	11,11,12	0.47	0	13,15,17	1.42	2 (15%)
5	MAN	A	311	5	11,11,12	0.46	0	13,15,17	1.60	3 (23%)
3	NAG	P	401	1,3	14,14,15	0.74	1 (7%)	15,19,21	0.69	0
3	NAG	P	402	3,4	14,14,15	0.34	0	15,19,21	0.77	0
4	BMA	P	403	3	11,11,12	0.25	0	13,15,17	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	301	3	-	0/6/23/26	0/1/1/1
3	NAG	A	302	3,4	-	0/6/23/26	0/1/1/1
4	BMA	A	303	3,5	-	0/2/19/22	0/1/1/1
5	MAN	A	304	5,4	-	0/2/19/22	0/1/1/1
5	MAN	A	305	5	-	0/2/19/22	0/1/1/1
5	MAN	A	306	5	-	0/2/19/22	0/1/1/1
5	MAN	A	307	5	-	0/2/19/22	0/1/1/1
5	MAN	A	308	5	-	0/2/19/22	0/1/1/1
5	MAN	A	309	5	-	0/2/19/22	0/1/1/1
5	MAN	A	310	5	-	0/2/19/22	0/1/1/1
5	MAN	A	311	5	-	0/2/19/22	0/1/1/1
3	NAG	P	401	1,3	-	0/6/23/26	0/1/1/1
3	NAG	P	402	3,4	-	0/6/23/26	0/1/1/1
4	BMA	P	403	3	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	P	401	NAG	C1-C2	2.58	1.56	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	309	MAN	O5-C1-C2	-2.68	106.58	110.79
5	A	311	MAN	C1-O5-C5	-2.60	108.58	112.17
5	A	310	MAN	C3-C4-C5	2.09	113.89	110.22
5	A	311	MAN	C3-C4-C5	2.35	114.36	110.22
4	A	303	BMA	C1-C2-C3	2.41	112.70	109.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	302	NAG	1	0
4	A	303	BMA	1	0
5	A	307	MAN	4	0
5	A	308	MAN	1	0
5	A	310	MAN	2	0
5	A	311	MAN	1	0
4	P	403	BMA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	P	30/30 (100%)	0.52	2 (6%) 19 19	26, 36, 75, 96	0
2	A	235/256 (91%)	0.17	3 (1%) 77 77	23, 36, 54, 78	0
All	All	265/286 (92%)	0.21	5 (1%) 67 67	23, 36, 56, 96	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	P	303	THR	5.7
2	A	126	SER	2.6
1	P	305	PRO	2.3
2	A	42	GLY	2.2
2	A	65	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	MAN	A	311	11/12	0.65	0.47	16.58	30,30,30,30	0
5	MAN	A	305	11/12	0.86	0.16	3.23	30,30,30,30	0
3	NAG	A	302	14/15	0.89	0.14	0.37	30,30,30,30	0
3	NAG	P	401	14/15	0.84	0.15	-0.20	30,30,30,30	0
3	NAG	A	301	14/15	0.91	0.13	-0.64	30,30,30,30	0
5	MAN	A	306	11/12	0.93	0.11	-0.69	30,30,30,30	0
5	MAN	A	309	11/12	0.39	0.54	-	30,30,30,30	0
5	MAN	A	308	11/12	0.64	0.40	-	30,30,30,30	0
5	MAN	A	307	11/12	0.73	0.36	-	30,30,30,30	0
3	NAG	P	402	14/15	0.92	0.19	-	30,30,30,30	0
4	BMA	P	403	11/12	0.86	0.27	-	30,30,30,30	0
4	BMA	A	303	11/12	0.85	0.13	-	30,30,30,30	0
5	MAN	A	310	11/12	0.78	0.36	-	30,30,30,30	0
5	MAN	A	304	11/12	0.92	0.12	-	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.