



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2017 – 03:01 AM EST

PDB ID : 1CCE
Title : CONSTRUCTION OF A BIS-AQUO HEME ENZYME AND REPLACEMENT WITH EXOGENOUS LIGAND
Authors : Mcree, D.E.; Jensen, G.M.; Fitzgerald, M.M.; Siegel, H.A.; Goodin, D.B.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	291	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2976 atoms, of which 505 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	291	Total	C	H	N	O	S	0	0	0
			2847	1497	505	388	451	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	THR	CONFLICT	UNP P00431
A	152	GLY	ASP	CONFLICT	UNP P00431
A	175	GLY	HIS	CONFLICT	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	86	Total	O	0	0
			86	86		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CYTOCHROME C PEROXIDASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.20 Å 74.30 Å 45.40 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (7.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.190 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2976	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/2407	1.75	47/3258 (1.4%)

There are no bond length outliers.

All (47) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	ARG	NE-CZ-NH1	15.69	128.14	120.30
1	A	130	ARG	NE-CZ-NH2	-15.14	112.73	120.30
1	A	155	ARG	NE-CZ-NH2	-15.03	112.78	120.30
1	A	130	ARG	NE-CZ-NH1	14.47	127.53	120.30
1	A	14	ARG	NE-CZ-NH1	13.59	127.10	120.30
1	A	160	ARG	NE-CZ-NH1	12.78	126.69	120.30
1	A	251	TYR	CB-CG-CD2	-10.80	114.52	121.00
1	A	160	ARG	NE-CZ-NH2	-9.63	115.49	120.30
1	A	172	MET	CG-SD-CE	-8.71	86.27	100.20
1	A	101	TRP	CD1-CG-CD2	8.56	113.15	106.30
1	A	101	TRP	CE2-CD2-CG	-7.93	100.95	107.30
1	A	51	TRP	CD1-CG-CD2	7.93	112.64	106.30
1	A	191	TRP	CD1-CG-CD2	7.76	112.51	106.30
1	A	51	TRP	CE2-CD2-CG	-7.54	101.27	107.30
1	A	278	LYS	CA-CB-CG	-7.13	97.71	113.40
1	A	166	ARG	NE-CZ-NH1	7.07	123.84	120.30
1	A	57	TRP	CE2-CD2-CG	-7.02	101.69	107.30
1	A	251	TYR	CB-CG-CD1	6.95	125.17	121.00
1	A	14	ARG	NE-CZ-NH2	-6.86	116.87	120.30
1	A	163	MET	CG-SD-CE	-6.85	89.25	100.20
1	A	229	TYR	CB-CG-CD2	-6.84	116.90	121.00
1	A	211	TRP	CD1-CG-CD2	6.82	111.75	106.30
1	A	16	TYR	CB-CG-CD2	-6.77	116.94	121.00
1	A	57	TRP	CD1-CG-CD2	6.72	111.68	106.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	101	TRP	CB-CG-CD1	-6.58	118.44	127.00
1	A	223	TRP	CD1-CG-CD2	6.50	111.50	106.30
1	A	223	TRP	CE2-CD2-CG	-6.45	102.14	107.30
1	A	126	TRP	CD1-CG-CD2	6.45	111.46	106.30
1	A	72	ARG	NE-CZ-NH1	6.33	123.47	120.30
1	A	191	TRP	CE2-CD2-CG	-6.25	102.30	107.30
1	A	211	TRP	CE2-CD2-CG	-6.06	102.45	107.30
1	A	244	TYR	CB-CG-CD2	-6.02	117.39	121.00
1	A	101	TRP	CG-CD2-CE3	5.98	139.28	133.90
1	A	126	TRP	CE2-CD2-CG	-5.95	102.54	107.30
1	A	101	TRP	CG-CD1-NE1	-5.68	104.42	110.10
1	A	191	TRP	CG-CD1-NE1	-5.51	104.59	110.10
1	A	23	TYR	CB-CG-CD2	-5.40	117.76	121.00
1	A	57	TRP	CG-CD2-CE3	5.37	138.74	133.90
1	A	14	ARG	CB-CG-CD	5.28	125.33	111.60
1	A	175	GLY	O-C-N	-5.24	114.32	122.70
1	A	195	ASN	N-CA-C	5.23	125.13	111.00
1	A	61	ASP	CB-CG-OD1	5.14	122.93	118.30
1	A	162	ASN	CA-CB-CG	5.13	124.70	113.40
1	A	162	ASN	CA-C-N	-5.11	105.97	117.20
1	A	48	ARG	CB-CG-CD	5.06	124.76	111.60
1	A	160	ARG	CB-CG-CD	5.03	124.69	111.60
1	A	48	ARG	NE-CZ-NH1	5.02	122.81	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2342	505	2214	18	0
2	A	43	0	30	1	0
3	A	86	0	0	0	0
All	All	2471	505	2244	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:GLN:HE22	1:A:287:LYS:H	1.37	0.72
1:A:125:PRO:HG3	1:A:285:ILE:HD11	1.80	0.63
1:A:200:ASN:H	1:A:255:GLN:HE21	1.45	0.62
1:A:74:LYS:H	1:A:74:LYS:HD2	1.67	0.60
1:A:20:GLN:HE22	1:A:287:LYS:N	2.06	0.53
1:A:168:VAL:O	1:A:172:MET:HB2	2.12	0.50
1:A:51:TRP:CD1	1:A:52:HIS:HD2	2.31	0.48
1:A:167:GLU:O	1:A:171:LEU:HB2	2.15	0.46
1:A:127:ARG:O	1:A:130:ARG:NH2	2.49	0.46
1:A:200:ASN:HB3	1:A:255:GLN:HG3	1.98	0.46
1:A:14:ARG:HG3	1:A:101:TRP:CZ3	2.53	0.44
1:A:48:ARG:HD3	2:A:1:HEM:C4D	2.53	0.43
1:A:164:ASN:O	1:A:168:VAL:HG23	2.19	0.42
1:A:218:ALA:O	1:A:219:ASN:HB2	2.19	0.42
1:A:236:TYR:O	1:A:239:ILE:HG12	2.20	0.42
1:A:14:ARG:HG3	1:A:101:TRP:CE3	2.56	0.41
1:A:155:ARG:HH22	1:A:241:ASP:CG	2.24	0.40
1:A:151:ALA:HB2	1:A:237:SER:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/291 (99%)	279 (96%)	9 (3%)	1 (0%)	44 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	246/248 (99%)	236 (96%)	10 (4%)	35 48

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	35	GLU
1	A	47	VAL
1	A	51	TRP
1	A	74	LYS
1	A	81	SER
1	A	143	ARG
1	A	156	THR
1	A	195	ASN
1	A	279	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	195	ASN
1	A	220	ASN
1	A	240	GLN
1	A	255	GLN
1	A	292	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	1	3	28,50,50	1.64	7 (25%)	17,82,82	1.99	6 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	1	3	-	0/6/54/54	0/0/8/8

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	HEM	C3B-CAB	-3.62	1.40	1.47
2	A	1	HEM	C4C-NC	-3.45	1.32	1.36
2	A	1	HEM	C3C-CAC	-3.35	1.41	1.47
2	A	1	HEM	C4D-ND	-2.56	1.33	1.36
2	A	1	HEM	C1B-NB	-2.55	1.33	1.36
2	A	1	HEM	CBC-CAC	2.35	1.45	1.28
2	A	1	HEM	CBB-CAB	2.40	1.45	1.28

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	HEM	C1D-C2D-C3D	-3.49	104.57	107.00
2	A	1	HEM	C4A-C3A-C2A	-2.59	105.19	107.00
2	A	1	HEM	CAD-CBD-CGD	2.48	116.90	112.66
2	A	1	HEM	CAA-CBA-CGA	2.55	117.02	112.66
2	A	1	HEM	CMB-C2B-C3B	2.75	129.99	124.89
2	A	1	HEM	CMC-C2C-C3C	2.77	130.04	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.