



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 07:00 pm GMT

PDB ID : 2CCI
Title : CRYSTAL STRUCTURE OF PHOSPHO-CDK2 CYCLIN A IN COMPLEX WITH A PEPTIDE CONTAINING BOTH THE SUBSTRATE AND RECRUITMENT SITES OF CDC6
Authors : Cheng, K.Y.; Noble, M.E.M.; Skamnaki, V.; Brown, N.R.; Lowe, E.D.; Kontogiannis, L.; Shen, K.; Cole, P.A.; Siligardi, G.; Johnson, L.N.
Deposited on : 2006-01-16
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	<div> <div>5%</div> <div>79%</div> <div>18%</div> <div>..</div> </div>
1	C	299	<div> <div>18%</div> <div>80%</div> <div>17%</div> <div>..</div> </div>
2	B	258	<div> <div>%</div> <div>85%</div> <div>14%</div> <div>.</div> </div>
2	D	258	<div> <div>13%</div> <div>80%</div> <div>18%</div> <div>.</div> </div>
3	F	30	<div> <div>23%</div> <div>50%</div> <div>10%</div> <div>•</div> <div>37%</div> </div>
3	I	30	<div> <div>17%</div> <div>30%</div> <div>13%</div> <div>•</div> <div>53%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9391 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	P	S	0	0	0
			2388	1550	404	425	1	8			
1	C	297	Total	C	N	O	P	S	0	0	0
			2388	1550	404	425	1	8			

- Molecule 2 is a protein called CYCLIN A2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	258	Total	C	N	O	S		0	0	0
			2083	1350	339	383	11				
2	D	258	Total	C	N	O	S		0	0	0
			2083	1350	339	383	11				



- Molecule 3 is a protein called CELL DIVISION CONTROL PROTEIN 6 HOMOLOG.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	F	19	Total	C	N	O	0	0	0
			160	99	37	24			
3	I	14	Total	C	N	O	0	0	0
			114	73	25	16			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	67	HIS	PRO	ENGINEERED MUTATION	UNP Q99741
F	68	HIS	PRO	ENGINEERED MUTATION	UNP Q99741
F	69	ALA	CYS	ENGINEERED MUTATION	UNP Q99741
F	72	ARG	PRO	ENGINEERED MUTATION	UNP Q99741
I	69	ALA	CYS	ENGINEERED MUTATION	UNP Q99741
I	72	ARG	PRO	ENGINEERED MUTATION	UNP Q99741

- Molecule 4 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
4	C	1	Total 31	C 10	N 5	O 13	P 3	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0
5	C	1	Total Mg 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	44	Total O 44 44	0	0
6	B	23	Total O 23 23	0	0
6	C	22	Total O 22 22	0	0
6	D	12	Total O 12 12	0	0
6	F	7	Total O 7 7	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	I	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CELL DIVISION PROTEIN KINASE 2



• Molecule 1: CELL DIVISION PROTEIN KINASE 2




• Molecule 2: CYCLIN A2




• Molecule 2: CYCLIN A2






• Molecule 3: CELL DIVISION CONTROL PROTEIN 6 HOMOLOG



• Molecule 3: CELL DIVISION CONTROL PROTEIN 6 HOMOLOG



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.42Å 114.39Å 170.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	94.92 – 2.70 37.02 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.0 (94.92-2.70) 97.1 (37.02-2.70)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.261 , 0.321 0.261 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	42.6	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 49.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	9391	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/2438	0.59	1/3308 (0.0%)
1	C	0.39	0/2438	0.55	0/3308
2	B	0.43	0/2133	0.56	0/2897
2	D	0.37	0/2133	0.53	0/2897
3	F	0.37	0/163	0.50	0/215
3	I	0.43	0/114	0.60	0/148
All	All	0.41	0/9419	0.56	1/12773 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	LEU	CA-CB-CG	5.01	126.83	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2430	43	0
1	C	2388	0	2430	31	0
2	B	2083	0	2107	24	0
2	D	2083	0	2107	28	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	160	0	161	2	0
3	I	114	0	130	2	0
4	A	31	0	12	0	0
4	C	31	0	12	1	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	44	0	0	3	0
6	B	23	0	0	0	0
6	C	22	0	0	0	0
6	D	12	0	0	0	0
6	F	7	0	0	1	0
6	I	3	0	0	1	0
All	All	9391	0	9389	117	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (117) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:203:GLN:HE22	2:D:247:SER:HA	1.22	1.05
1:A:154:VAL:O	2:B:316:THR:HG22	1.62	0.97
1:C:1:MET:HE1	1:C:70:ILE:HD13	1.51	0.90
1:C:155:PRO:HD2	2:D:316:THR:HG22	1.54	0.89
1:A:5:GLN:HB2	1:A:24:LYS:HE3	1.66	0.77
2:B:366:THR:HG23	2:B:427:PRO:HD3	1.68	0.75
1:A:177:CYS:HB2	6:A:2023:HOH:O	1.87	0.75
1:A:95:ALA:HA	1:A:199:ARG:HH11	1.54	0.72
1:A:42:GLU:OE1	2:B:275:VAL:HG23	1.89	0.71
3:F:67:HIS:O	3:F:68:HIS:HB2	1.90	0.71
1:A:154:VAL:O	2:B:316:THR:CG2	2.39	0.71
1:C:1:MET:HE2	1:C:1:MET:HA	1.71	0.71
2:D:219:VAL:HG22	2:D:232:LEU:HD11	1.72	0.71
1:A:155:PRO:HD2	2:B:316:THR:HG23	1.73	0.71
1:A:51:GLU:O	1:A:55:LEU:HB2	1.91	0.70
2:D:203:GLN:NE2	2:D:247:SER:HA	2.03	0.67
1:C:230:VAL:HA	1:C:233:MET:HE2	1.76	0.67
1:C:1:MET:CE	1:C:1:MET:HA	2.24	0.66
1:C:73:GLU:HG2	2:D:293:ARG:HH22	1.64	0.62
3:F:73:LYS:HA	6:F:2004:HOH:O	2.00	0.62
1:A:175:LEU:HD13	1:A:233:MET:HE3	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:51:GLU:O	1:C:55:LEU:HB2	2.01	0.61
1:A:230:VAL:HG23	1:A:233:MET:HE1	1.82	0.61
2:B:248:VAL:HG12	2:B:249:LEU:O	2.03	0.58
1:A:252:VAL:HG12	1:A:254:PRO:HD2	1.86	0.57
2:B:219:VAL:HG22	2:B:232:LEU:HD11	1.85	0.57
1:C:64:VAL:HG23	1:C:143:LEU:O	2.04	0.57
1:A:177:CYS:SG	1:A:233:MET:SD	3.02	0.57
1:A:57:GLU:OE2	2:B:307:ALA:HB3	2.05	0.57
1:A:71:HIS:NE2	2:B:296:HIS:CD2	2.74	0.56
4:C:1297:ATP:O1G	4:C:1297:ATP:O1A	2.22	0.56
1:A:39:THR:HG22	1:A:40:GLU:HG2	1.87	0.56
2:D:347:TYR:OH	2:D:394:LEU:HA	2.06	0.56
2:B:344:ALA:HB1	2:B:348:LEU:HD22	1.87	0.55
1:A:1:MET:HA	1:A:1:MET:HE2	1.88	0.55
1:C:5:GLN:HB2	1:C:24:LYS:HE3	1.89	0.54
1:C:72:THR:HB	1:C:75:LYS:H	1.72	0.54
1:A:9:LYS:HD3	1:A:17:VAL:HG11	1.90	0.54
1:C:13:GLY:O	1:C:15:TYR:N	2.40	0.54
1:A:218:THR:HA	1:A:246:GLN:NE2	2.23	0.54
1:A:163:VAL:HG13	1:A:164:VAL:HG23	1.88	0.54
1:A:95:ALA:HA	1:A:199:ARG:HD2	1.89	0.54
2:D:255:LEU:HB2	2:D:286:TYR:CE1	2.44	0.52
1:C:64:VAL:HG21	1:C:144:ALA:HB2	1.92	0.52
2:D:344:ALA:HB1	2:D:348:LEU:HD22	1.92	0.52
2:D:407:GLN:OE1	2:D:410:ARG:HD3	2.10	0.52
1:A:115:LEU:HD11	1:A:185:ASP:HB3	1.92	0.52
2:D:215:VAL:HG12	2:D:342:ILE:HD13	1.91	0.52
2:B:303:THR:O	2:B:304:PHE:HB2	2.10	0.51
2:B:388:LYS:HB3	2:B:389:PRO:HD3	1.93	0.51
1:C:130:PRO:HD3	1:C:192:ILE:HG12	1.94	0.50
2:D:222:GLY:HA2	2:D:227:LEU:HD12	1.93	0.50
1:A:60:HIS:HD2	1:A:62:ASN:H	1.60	0.49
1:A:119:HIS:HE1	1:A:185:ASP:OD2	1.95	0.49
2:D:230:GLU:OE2	2:D:313:GLN:NE2	2.40	0.49
2:B:416:SER:HB3	1:C:8:GLU:OE2	2.13	0.49
1:A:60:HIS:CD2	1:A:62:ASN:H	2.31	0.48
1:A:71:HIS:CE1	2:B:296:HIS:HD2	2.32	0.48
1:A:249:SER:HA	1:A:260:ARG:HD3	1.94	0.47
2:B:262:LEU:HD11	2:B:266:LYS:HE3	1.97	0.47
2:D:329:VAL:HG11	2:D:364:LEU:HD12	1.96	0.47
1:A:252:VAL:HB	1:A:255:LEU:HD22	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:162:GLU:CD	1:C:162:GLU:H	2.18	0.47
1:C:164:VAL:HB	1:C:169:ARG:HG3	1.97	0.47
1:A:162:GLU:H	1:A:162:GLU:CD	2.18	0.47
1:A:65:LYS:HE2	6:A:2009:HOH:O	2.14	0.46
2:D:335:PHE:CZ	2:D:339:LEU:HD11	2.50	0.46
2:D:183:HIS:HB2	2:D:317:GLN:HE22	1.82	0.45
1:A:71:HIS:NE2	2:B:296:HIS:HD2	2.12	0.45
2:B:347:TYR:OH	2:B:394:LEU:HA	2.16	0.45
2:D:277:GLU:O	2:D:281:ILE:HG13	2.16	0.45
1:A:60:HIS:HB3	1:A:63:ILE:HD12	1.98	0.45
2:D:213:ILE:HG22	3:I:92:LEU:HD22	1.99	0.45
1:A:126:ARG:NH2	1:A:150:ARG:HB3	2.31	0.44
1:C:220:GLY:HA3	1:C:245:ARG:HG2	1.99	0.44
1:C:181:SER:O	1:C:184:VAL:HG22	2.17	0.44
1:A:52:ILE:HD11	1:A:78:LEU:HD21	2.00	0.43
1:A:230:VAL:HA	1:A:233:MET:HE2	2.01	0.43
2:B:388:LYS:O	2:B:392:MET:HG2	2.18	0.43
2:D:250:ARG:HD3	6:I:2003:HOH:O	2.18	0.43
1:A:71:HIS:HA	1:A:76:LEU:HD12	1.99	0.43
1:C:88:LYS:HA	1:C:91:MET:HE2	2.00	0.43
1:C:154:VAL:HA	1:C:155:PRO:HA	1.86	0.43
1:C:5:GLN:HG2	1:C:22:ARG:NH1	2.34	0.43
2:D:190:GLU:N	2:D:309:PRO:HG2	2.33	0.43
1:C:133:LEU:HD11	1:C:192:ILE:HD13	2.01	0.43
1:C:38:ASP:HB2	1:C:39:THR:H	1.59	0.43
2:D:254:GLN:HE21	2:D:282:THR:HG22	1.83	0.43
1:A:41:THR:HB	1:A:42:GLU:H	1.73	0.43
1:C:17:VAL:HB	1:C:19:TYR:CE2	2.53	0.43
1:C:172:GLU:O	1:C:177:CYS:HB2	2.19	0.42
1:A:230:VAL:HG23	1:A:233:MET:CE	2.50	0.42
1:A:167:TRP:CD1	1:A:204:PRO:HA	2.54	0.42
2:D:425:ASN:HA	2:D:426:PRO:HD2	1.96	0.42
1:C:108:LEU:HD22	1:C:193:PHE:CD1	2.54	0.42
2:D:278:PHE:HA	2:D:281:ILE:HD11	2.00	0.42
2:B:410:ARG:O	2:B:414:LYS:HG3	2.18	0.42
2:D:210:MET:HE1	2:D:250:ARG:HB3	2.01	0.42
1:C:155:PRO:HD2	2:D:316:THR:CG2	2.37	0.42
1:C:53:SER:HB3	2:D:304:PHE:O	2.20	0.42
2:D:334:MET:O	2:D:338:GLU:HB2	2.20	0.41
2:B:398:TYR:CE2	2:B:426:PRO:HB3	2.55	0.41
2:D:361:HIS:HB2	2:D:372:TRP:HB2	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:GLY:HA2	1:A:199:ARG:NE	2.36	0.41
1:A:208:GLU:HG3	6:A:2029:HOH:O	2.19	0.41
2:B:404:HIS:O	2:B:407:GLN:NE2	2.54	0.41
1:A:195:GLU:O	1:A:199:ARG:HA	2.21	0.41
2:B:332:LEU:HD23	2:B:363:ALA:HA	2.02	0.41
1:C:249:SER:HA	1:C:260:ARG:HD2	2.02	0.41
1:C:122:ARG:O	1:C:122:ARG:HD2	2.21	0.41
1:C:71:HIS:HD2	1:C:76:LEU:HD13	1.86	0.40
1:A:162:GLU:CD	1:A:162:GLU:N	2.75	0.40
1:A:248:PHE:CZ	1:A:264:SER:HA	2.56	0.40
2:D:236:VAL:HA	2:D:239:ILE:HD12	2.03	0.40
3:I:91:ARG:O	3:I:91:ARG:HG2	2.21	0.40
2:B:345:ASP:HA	2:B:346:PRO:HA	1.88	0.40
2:B:254:GLN:HB3	2:B:286:TYR:OH	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/299 (98%)	270 (92%)	18 (6%)	6 (2%)	9	22
1	C	294/299 (98%)	266 (90%)	26 (9%)	2 (1%)	25	53
2	B	256/258 (99%)	246 (96%)	9 (4%)	1 (0%)	38	66
2	D	256/258 (99%)	239 (93%)	16 (6%)	1 (0%)	38	66
3	F	15/30 (50%)	13 (87%)	1 (7%)	1 (7%)	1	2
3	I	10/30 (33%)	6 (60%)	3 (30%)	1 (10%)	1	0
All	All	1125/1174 (96%)	1040 (92%)	73 (6%)	12 (1%)	17	40

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	12	GLU
2	B	177	ASP
1	C	14	THR
1	A	11	GLY
1	A	164	VAL
1	C	164	VAL
3	I	92	LEU
1	A	14	THR
3	F	68	HIS
1	A	38	ASP
1	A	145	ASP
2	D	281	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	261/263 (99%)	250 (96%)	11 (4%)	34	65
1	C	261/263 (99%)	245 (94%)	16 (6%)	22	47
2	B	232/232 (100%)	226 (97%)	6 (3%)	51	81
2	D	232/232 (100%)	223 (96%)	9 (4%)	37	68
3	F	17/26 (65%)	16 (94%)	1 (6%)	23	49
3	I	12/26 (46%)	9 (75%)	3 (25%)	1	2
All	All	1015/1042 (97%)	969 (96%)	46 (4%)	32	62

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	THR
1	A	17	VAL
1	A	55	LEU
1	A	74	ASN
1	A	121	HIS
1	A	122	ARG
1	A	131	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	150	ARG
1	A	163	VAL
1	A	226	VAL
1	A	255	LEU
2	B	175	VAL
2	B	232	LEU
2	B	274	GLU
2	B	292	LEU
2	B	348	LEU
2	B	429	THR
1	C	1	MET
1	C	22	ARG
1	C	38	ASP
1	C	42	GLU
1	C	78	LEU
1	C	120	SER
1	C	122	ARG
1	C	150	ARG
1	C	177	CYS
1	C	206	ASP
1	C	248	PHE
1	C	250	LYS
1	C	252	VAL
1	C	255	LEU
1	C	268	HIS
1	C	272	ASN
2	D	247	SER
2	D	284	ASP
2	D	292	LEU
2	D	293	ARG
2	D	348	LEU
2	D	368	THR
2	D	428	GLU
2	D	429	THR
2	D	431	ASN
3	F	72	ARG
3	I	70	SER
3	I	72	ARG
3	I	93	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (24) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	A	60	HIS
1	A	62	ASN
1	A	84	HIS
1	A	119	HIS
1	A	246	GLN
1	A	265	GLN
1	A	287	GLN
2	B	183	HIS
2	B	296	HIS
2	B	317	GLN
2	B	395	HIS
2	B	396	GLN
2	B	403	GLN
2	B	425	ASN
2	B	431	ASN
1	C	60	HIS
1	C	71	HIS
1	C	84	HIS
1	C	85	GLN
2	D	233	HIS
2	D	317	GLN
2	D	425	ASN
3	F	85	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	160	1	9,10,11	2.67	3 (33%)	10,14,16	0.82	0
1	TPO	C	160	1	9,10,11	2.80	4 (44%)	10,14,16	0.89	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/8/11/13	0/0/0/0
1	TPO	C	160	1	-	0/8/11/13	0/0/0/0

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	160	TPO	CA-C	2.52	1.53	1.50
1	A	160	TPO	P-O2P	3.62	1.69	1.54
1	A	160	TPO	P-O3P	3.69	1.70	1.54
1	C	160	TPO	P-O3P	3.72	1.70	1.54
1	C	160	TPO	P-O2P	3.77	1.70	1.54
1	A	160	TPO	P-O1P	5.60	1.69	1.50
1	C	160	TPO	P-O1P	5.81	1.70	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	160	TPO	O-C-CA	-2.06	120.34	125.15

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ATP	A	1297	5	27,33,33	1.07	1 (3%)	25,52,52	1.54	2 (8%)
4	ATP	C	1297	5	27,33,33	1.02	2 (7%)	25,52,52	1.43	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ATP	A	1297	5	-	0/18/38/38	0/3/3/3
4	ATP	C	1297	5	-	0/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	1297	ATP	PG-O3B	2.12	1.63	1.60
4	A	1297	ATP	C5-C4	3.29	1.47	1.40
4	C	1297	ATP	C5-C4	3.32	1.48	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1297	ATP	N3-C2-N1	-5.97	123.65	128.86
4	C	1297	ATP	N3-C2-N1	-5.28	124.25	128.86
4	C	1297	ATP	C4-C5-N7	-2.55	106.95	109.41
4	A	1297	ATP	C4-C5-N7	-2.39	107.10	109.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	1297	ATP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/299 (98%)	0.63	16 (5%) 26 25	30, 35, 46, 51	0
1	C	296/299 (98%)	1.17	54 (18%) 1 1	30, 38, 46, 48	0
2	B	258/258 (100%)	0.32	2 (0%) 86 86	31, 36, 41, 48	0
2	D	258/258 (100%)	0.98	33 (12%) 4 3	30, 37, 44, 49	0
3	F	19/30 (63%)	1.56	7 (36%) 0 0	42, 45, 56, 57	0
3	I	14/30 (46%)	1.86	5 (35%) 0 0	53, 61, 66, 66	0
All	All	1141/1174 (97%)	0.81	117 (10%) 7 5	30, 37, 46, 66	0

All (117) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	THR	7.7
1	C	39	THR	7.2
1	A	40	GLU	6.5
1	C	248	PHE	5.8
2	D	175	VAL	5.5
1	A	73	GLU	5.1
1	A	71	HIS	5.0
1	C	245	ARG	4.9
3	I	86	THR	4.9
1	A	74	ASN	4.8
2	D	323	GLN	4.8
1	C	247	ASP	4.8
1	C	40	GLU	4.7
1	C	137	THR	4.5
1	C	293	VAL	4.5
1	C	213	PHE	4.4
1	C	241	PRO	4.3
1	C	288	ASP	4.3
1	C	73	GLU	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	95	ALA	4.2
1	C	74	ASN	4.2
2	D	367	VAL	4.2
1	C	0	SER	4.0
2	D	432	LEU	3.9
1	C	296	LEU	3.7
1	C	295	HIS	3.6
1	A	295	HIS	3.6
1	C	287	GLN	3.5
1	C	13	GLY	3.5
2	B	175	VAL	3.4
2	D	178	TYR	3.4
1	C	246	GLN	3.4
1	C	243	TRP	3.3
1	C	238	PRO	3.3
1	C	217	ARG	3.2
1	C	101	LEU	3.2
2	D	371	SER	3.1
1	C	16	GLY	3.1
2	D	180	GLU	3.1
2	D	176	PRO	3.1
1	C	251	VAL	3.0
1	C	100	PRO	3.0
3	F	72	ARG	3.0
3	F	85	HIS	3.0
1	C	235	ASP	3.0
1	C	242	LYS	2.9
3	I	93	VAL	2.9
1	C	294	PRO	2.9
2	D	425	ASN	2.9
1	C	221	THR	2.9
2	D	372	TRP	2.9
1	C	207	SER	2.9
1	C	289	VAL	2.9
2	D	402	PRO	2.8
2	D	324	PRO	2.8
3	F	67	HIS	2.8
1	C	240	PHE	2.8
1	A	96	LEU	2.8
2	D	429	THR	2.7
2	D	383	THR	2.7
1	C	239	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	375	SER	2.7
2	D	395	HIS	2.7
2	D	197	VAL	2.6
1	C	96	LEU	2.6
2	D	401	ALA	2.6
1	C	223	ASP	2.6
2	D	420	GLY	2.6
2	D	281	ILE	2.5
1	C	209	ILE	2.5
1	C	290	THR	2.5
1	A	38	ASP	2.5
3	I	69	ALA	2.5
1	A	14	THR	2.5
2	D	421	VAL	2.5
3	F	95	ASP	2.5
1	C	225	VAL	2.4
3	F	96	ASN	2.4
1	C	268	HIS	2.4
1	A	36	ARG	2.4
1	C	226	VAL	2.4
1	C	227	TRP	2.4
1	A	97	THR	2.4
1	C	14	THR	2.4
1	A	77	TYR	2.3
2	D	403	GLN	2.3
1	C	253	PRO	2.3
1	C	249	SER	2.3
1	C	237	LYS	2.3
2	D	368	THR	2.3
1	A	75	LYS	2.3
1	C	222	PRO	2.3
1	C	2	GLU	2.3
2	B	177	ASP	2.3
1	A	72	THR	2.2
1	C	180	TYR	2.2
1	A	296	LEU	2.2
2	D	380	THR	2.2
1	C	38	ASP	2.2
2	D	296	HIS	2.1
2	D	369	GLY	2.1
1	A	95	ALA	2.1
2	D	423	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	I	87	LEU	2.1
1	C	176	GLY	2.1
2	D	415	ASN	2.1
2	D	399	LEU	2.1
1	C	257	GLU	2.1
1	C	206	ASP	2.1
3	I	92	LEU	2.1
3	F	73	LYS	2.1
1	C	35	ILE	2.1
2	D	329	VAL	2.1
2	D	365	TYR	2.1
3	F	86	THR	2.1
2	D	325	ALA	2.0
2	D	390	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	C	160	11/12	0.91	0.17	-	35,36,36,36	0
1	TPO	A	160	11/12	0.92	0.22	-	31,33,33,33	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ATP	A	1297	31/31	0.90	0.19	-0.49	35,37,58,58	0
4	ATP	C	1297	31/31	0.91	0.18	-0.67	32,36,55,56	0
5	MG	C	1298	1/1	0.88	0.16	-1.80	43,43,43,43	0
5	MG	A	1298	1/1	0.89	0.08	-1.87	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.