



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 16, 2017 – 05:38 PM EDT

PDB ID : 3CC9
Title : Crystal structure of Plasmodium vivax putative polyprenyl pyrophosphate synthase in complex with geranylgeranyl diphosphate
Authors : Wernimont, A.K.; Dunford, J.; Lew, J.; Zhao, Y.; Kozieradzki, I.; Cossar, D.; Schapira, M.; Bochkarev, A.; Arrowsmith, C.H.; Bountra, C.; Weigelt, J.; Edwards, A.M.; Hui, R.; Artz, J.D.; Structural Genomics Consortium (SGC)
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	<div> <div>13%</div> <div> <div></div> <div>74%</div> <div>11%</div> <div>•</div> <div>13%</div> </div> </div>
1	B	396	<div> <div>13%</div> <div> <div></div> <div>84%</div> <div>7%</div> <div></div> <div>10%</div> </div> </div>
1	C	396	<div> <div>25%</div> <div> <div></div> <div>76%</div> <div>8%</div> <div></div> <div>16%</div> </div> </div>
1	D	396	<div> <div>15%</div> <div> <div></div> <div>79%</div> <div>6%</div> <div>•</div> <div>15%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative farnesyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	344	Total	C	N	O	S	0	1	0
			2764	1793	439	517	15			
1	B	358	Total	C	N	O	S	0	2	0
			2885	1873	462	535	15			
1	C	332	Total	C	N	O	S	0	0	0
			2578	1663	420	481	14			
1	D	337	Total	C	N	O	S	0	0	0
			2639	1702	423	499	15			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	UNP A5K4U6
A	2	GLY	-	EXPRESSION TAG	UNP A5K4U6
A	3	SER	-	EXPRESSION TAG	UNP A5K4U6
A	4	SER	-	EXPRESSION TAG	UNP A5K4U6
A	5	HIS	-	EXPRESSION TAG	UNP A5K4U6
A	6	HIS	-	EXPRESSION TAG	UNP A5K4U6
A	7	HIS	-	EXPRESSION TAG	UNP A5K4U6
A	8	HIS	-	EXPRESSION TAG	UNP A5K4U6
A	9	HIS	-	EXPRESSION TAG	UNP A5K4U6
A	10	HIS	-	EXPRESSION TAG	UNP A5K4U6
A	11	SER	-	EXPRESSION TAG	UNP A5K4U6
A	12	SER	-	EXPRESSION TAG	UNP A5K4U6
A	13	GLY	-	EXPRESSION TAG	UNP A5K4U6
A	14	ARG	-	EXPRESSION TAG	UNP A5K4U6
A	15	GLU	-	EXPRESSION TAG	UNP A5K4U6
A	16	ASN	-	EXPRESSION TAG	UNP A5K4U6
A	17	LEU	-	EXPRESSION TAG	UNP A5K4U6
A	18	TYR	-	EXPRESSION TAG	UNP A5K4U6
A	19	PHE	-	EXPRESSION TAG	UNP A5K4U6
A	20	GLN	-	EXPRESSION TAG	UNP A5K4U6
A	21	GLY	-	EXPRESSION TAG	UNP A5K4U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	134	MET	THR	SEE REMARK 999	UNP A5K4U6
A	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6
B	1	MET	-	EXPRESSION TAG	UNP A5K4U6
B	2	GLY	-	EXPRESSION TAG	UNP A5K4U6
B	3	SER	-	EXPRESSION TAG	UNP A5K4U6
B	4	SER	-	EXPRESSION TAG	UNP A5K4U6
B	5	HIS	-	EXPRESSION TAG	UNP A5K4U6
B	6	HIS	-	EXPRESSION TAG	UNP A5K4U6
B	7	HIS	-	EXPRESSION TAG	UNP A5K4U6
B	8	HIS	-	EXPRESSION TAG	UNP A5K4U6
B	9	HIS	-	EXPRESSION TAG	UNP A5K4U6
B	10	HIS	-	EXPRESSION TAG	UNP A5K4U6
B	11	SER	-	EXPRESSION TAG	UNP A5K4U6
B	12	SER	-	EXPRESSION TAG	UNP A5K4U6
B	13	GLY	-	EXPRESSION TAG	UNP A5K4U6
B	14	ARG	-	EXPRESSION TAG	UNP A5K4U6
B	15	GLU	-	EXPRESSION TAG	UNP A5K4U6
B	16	ASN	-	EXPRESSION TAG	UNP A5K4U6
B	17	LEU	-	EXPRESSION TAG	UNP A5K4U6
B	18	TYR	-	EXPRESSION TAG	UNP A5K4U6
B	19	PHE	-	EXPRESSION TAG	UNP A5K4U6
B	20	GLN	-	EXPRESSION TAG	UNP A5K4U6
B	21	GLY	-	EXPRESSION TAG	UNP A5K4U6
B	134	MET	THR	SEE REMARK 999	UNP A5K4U6
B	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6
C	1	MET	-	EXPRESSION TAG	UNP A5K4U6
C	2	GLY	-	EXPRESSION TAG	UNP A5K4U6
C	3	SER	-	EXPRESSION TAG	UNP A5K4U6
C	4	SER	-	EXPRESSION TAG	UNP A5K4U6
C	5	HIS	-	EXPRESSION TAG	UNP A5K4U6
C	6	HIS	-	EXPRESSION TAG	UNP A5K4U6
C	7	HIS	-	EXPRESSION TAG	UNP A5K4U6
C	8	HIS	-	EXPRESSION TAG	UNP A5K4U6
C	9	HIS	-	EXPRESSION TAG	UNP A5K4U6
C	10	HIS	-	EXPRESSION TAG	UNP A5K4U6
C	11	SER	-	EXPRESSION TAG	UNP A5K4U6
C	12	SER	-	EXPRESSION TAG	UNP A5K4U6
C	13	GLY	-	EXPRESSION TAG	UNP A5K4U6
C	14	ARG	-	EXPRESSION TAG	UNP A5K4U6
C	15	GLU	-	EXPRESSION TAG	UNP A5K4U6
C	16	ASN	-	EXPRESSION TAG	UNP A5K4U6
C	17	LEU	-	EXPRESSION TAG	UNP A5K4U6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	18	TYR	-	EXPRESSION TAG	UNP A5K4U6
C	19	PHE	-	EXPRESSION TAG	UNP A5K4U6
C	20	GLN	-	EXPRESSION TAG	UNP A5K4U6
C	21	GLY	-	EXPRESSION TAG	UNP A5K4U6
C	134	MET	THR	SEE REMARK 999	UNP A5K4U6
C	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6
D	1	MET	-	EXPRESSION TAG	UNP A5K4U6
D	2	GLY	-	EXPRESSION TAG	UNP A5K4U6
D	3	SER	-	EXPRESSION TAG	UNP A5K4U6
D	4	SER	-	EXPRESSION TAG	UNP A5K4U6
D	5	HIS	-	EXPRESSION TAG	UNP A5K4U6
D	6	HIS	-	EXPRESSION TAG	UNP A5K4U6
D	7	HIS	-	EXPRESSION TAG	UNP A5K4U6
D	8	HIS	-	EXPRESSION TAG	UNP A5K4U6
D	9	HIS	-	EXPRESSION TAG	UNP A5K4U6
D	10	HIS	-	EXPRESSION TAG	UNP A5K4U6
D	11	SER	-	EXPRESSION TAG	UNP A5K4U6
D	12	SER	-	EXPRESSION TAG	UNP A5K4U6
D	13	GLY	-	EXPRESSION TAG	UNP A5K4U6
D	14	ARG	-	EXPRESSION TAG	UNP A5K4U6
D	15	GLU	-	EXPRESSION TAG	UNP A5K4U6
D	16	ASN	-	EXPRESSION TAG	UNP A5K4U6
D	17	LEU	-	EXPRESSION TAG	UNP A5K4U6
D	18	TYR	-	EXPRESSION TAG	UNP A5K4U6
D	19	PHE	-	EXPRESSION TAG	UNP A5K4U6
D	20	GLN	-	EXPRESSION TAG	UNP A5K4U6
D	21	GLY	-	EXPRESSION TAG	UNP A5K4U6
D	134	MET	THR	SEE REMARK 999	UNP A5K4U6
D	227	ASP	ASN	SEE REMARK 999	UNP A5K4U6

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0

- Molecule 3 is GERANYLGERANYL DIPHOSPHATE (three-letter code: GRG) (formula: C₂₀H₃₆O₇P₂).



- Molecule 4 is water.



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	107.09Å 108.98Å 141.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.61 – 2.30 34.40 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (34.61-2.30) 99.7 (34.40-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.65 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.241 , 0.296 0.273 , 0.310	Depositor DCC
R_{free} test set	3721 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	53.5	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.007 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	11221	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GRG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.36	0/2821	0.54	3/3829 (0.1%)
1	B	0.38	0/2952	0.53	2/3999 (0.1%)
1	C	0.33	0/2626	0.46	0/3569
1	D	0.34	0/2687	0.49	0/3644
All	All	0.36	0/11086	0.51	5/15041 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	157	LEU	CA-CB-CG	-5.83	101.88	115.30
1	A	84	ARG	NE-CZ-NH1	-5.60	117.50	120.30
1	A	123	LEU	CA-CB-CG	5.59	128.15	115.30
1	A	157	LEU	CA-CB-CG	-5.41	102.85	115.30
1	B	123	LEU	CA-CB-CG	5.29	127.46	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2764	0	2595	31	0
1	B	2885	0	2749	18	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2578	0	2319	17	0
1	D	2639	0	2417	14	0
2	A	1	0	0	0	0
3	A	29	0	21	2	0
3	B	29	0	21	2	0
3	C	27	0	15	0	0
3	D	29	0	23	1	0
4	A	75	0	0	0	0
4	B	83	0	0	1	0
4	C	47	0	0	1	0
4	D	35	0	0	0	0
All	All	11221	0	10160	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:ASN:HB3	1:B:189:LEU:HD11	1.71	0.73
1:A:161:ASN:HB3	1:B:189:LEU:CD1	2.19	0.72
1:A:157:LEU:HD22	3:B:501:GRG:H203	1.73	0.71
1:D:127:ASP:OD2	1:D:135:ARG:HG3	1.92	0.69
1:A:84:ARG:NH1	3:A:501:GRG:O2A	2.25	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	335/396 (85%)	329 (98%)	6 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	352/396 (89%)	341 (97%)	11 (3%)	0	100	100
1	C	316/396 (80%)	308 (98%)	8 (2%)	0	100	100
1	D	319/396 (81%)	308 (97%)	11 (3%)	0	100	100
All	All	1322/1584 (84%)	1286 (97%)	36 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/357 (78%)	266 (96%)	11 (4%)	36	50
1	B	294/357 (82%)	290 (99%)	4 (1%)	71	85
1	C	235/357 (66%)	228 (97%)	7 (3%)	46	63
1	D	255/357 (71%)	248 (97%)	7 (3%)	50	67
All	All	1061/1428 (74%)	1032 (97%)	29 (3%)	51	67

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	262	ILE
1	C	64	HIS
1	D	157	LEU
1	B	307	GLN
1	C	135	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	82	ASN
1	C	64	HIS
1	C	82	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	173	ASN
1	D	195	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GRG	A	501	-	27,28,28	2.87	7 (25%)	30,37,37	7.31	12 (40%)
3	GRG	B	501	-	27,28,28	2.83	8 (29%)	30,37,37	6.87	11 (36%)
3	GRG	C	501	-	25,26,28	2.96	7 (28%)	27,34,37	8.00	11 (40%)
3	GRG	D	501	-	27,28,28	2.82	8 (29%)	30,37,37	6.51	9 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GRG	A	501	-	-	1/31/31/31	0/0/0/0
3	GRG	B	501	-	-	2/31/31/31	0/0/0/0
3	GRG	C	501	-	-	0/29/29/31	0/0/0/0
3	GRG	D	501	-	-	0/31/31/31	0/0/0/0

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	501	GRG	C1-C2	-7.54	1.25	1.49
3	A	501	GRG	C1-C2	-7.41	1.26	1.49
3	C	501	GRG	C11-C12	-7.18	1.26	1.50
3	C	501	GRG	C6-C7	-7.14	1.26	1.50
3	D	501	GRG	C11-C12	-7.13	1.26	1.50

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	501	GRG	C4-C3-C2	-2.28	117.61	123.69
3	B	501	GRG	C10-C8-C7	-2.27	117.63	123.69
3	A	501	GRG	C10-C8-C7	-2.06	118.19	123.69
3	A	501	GRG	C14-C13-C12	-2.05	118.23	123.69
3	C	501	GRG	O1-C1-C2	2.06	118.88	109.73

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	GRG	C1-C2-C3-C4
3	B	501	GRG	C1-C2-C3-C4
3	B	501	GRG	O1-C1-C2-C3

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	GRG	2	0
3	B	501	GRG	2	0
3	D	501	GRG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	344/396 (86%)	0.74	50 (14%) 3 4	45, 57, 70, 76	1 (0%)
1	B	358/396 (90%)	0.75	52 (14%) 3 4	45, 57, 70, 80	0
1	C	332/396 (83%)	1.33	98 (29%) 1 0	49, 65, 77, 91	0
1	D	337/396 (85%)	0.91	61 (18%) 1 2	50, 61, 77, 110	0
All	All	1371/1584 (86%)	0.93	261 (19%) 1 2	45, 59, 75, 110	1 (0%)

The worst 5 of 261 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	349	TYR	7.1
1	D	206	TYR	6.8
1	C	330	ILE	6.3
1	C	354	ILE	5.7
1	C	325	PRO	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GRG	D	501	29/29	0.93	0.37	1.56	38,49,65,65	20
3	GRG	C	501	27/29	0.94	0.22	-0.16	67,70,74,75	0
3	GRG	B	501	29/29	0.96	0.21	-0.79	43,49,51,52	0
3	GRG	A	501	29/29	0.96	0.17	-1.01	43,48,54,55	0
2	NA	A	502	1/1	0.94	0.15	-1.79	54,54,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.