



Full wwPDB X-ray Structure Validation Report i

Feb 14, 2017 – 01:51 am GMT

PDB ID : 1CD9
Title : 2:2 COMPLEX OF G-CSF WITH ITS RECEPTOR
Authors : Aritomi, M.; Kunishima, N.; Okamoto, T.; Kuroki, R.; Ota, Y.; Morikawa, K.
Deposited on : 1999-03-08
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6080 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (GRANULOCYTE COLONY-STIMULATING FACTOR).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	171	Total	C	N	O	S	0	0	0
			1283	820	218	237	8			

1	C	169	Total	C	N	O	S	0	0	0
			1254	800	212	234	8			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	ALA	CLONING ARTIFACT	UNP P09919
C	1	MET	ALA	CLONING ARTIFACT	UNP P09919

- Molecule 2 is a protein called PROTEIN (G-CSF RECEPTOR).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	206	Total	C	N	O	S	0	0	0
			1638	1050	274	296	18			

2	D	204	Total	C	N	O	S	0	0	0
			1617	1036	270	293	18			

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C N O 14 8 1 5	0	0
3	D	1	Total C N O 14 8 1 5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	46	Total O 46 46	0	0
4	B	108	Total O 108 108	0	0
4	C	37	Total O 37 37	0	0
4	D	69	Total O 69 69	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROTEIN (GRANULOCYTE COLONY-STIMULATING FACTOR)




- Molecule 1: PROTEIN (GRANULOCYTE COLONY-STIMULATING FACTOR)



- Molecule 2: PROTEIN (G-CSF RECEPTOR)



- Molecule 2: PROTEIN (G-CSF RECEPTOR)





4 Data and refinement statistics i

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	125.47Å 125.47Å 372.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.80 44.14 – 2.80	Depositor EDS
% Data completeness (in resolution range)	95.7 (15.00-2.80) 78.3 (44.14-2.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	5.33 (at 2.81Å)	Xtriage
Refinement program	REFMAC	Depositor
R , R_{free}	0.237 , 0.319 0.198 , 0.249	Depositor DCC
R_{free} test set	425 reflections (1.48%)	DCC
Wilson B-factor (Å ²)	44.2	Xtriage
Anisotropy	0.119	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 73.6	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6080	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/1312	0.73	0/1784
1	C	0.66	0/1279	0.76	0/1738
2	B	0.65	0/1690	0.84	2/2306 (0.1%)
2	D	0.58	0/1669	0.81	2/2279 (0.1%)
All	All	0.62	0/5950	0.79	4/8107 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	151	ARG	NE-CZ-NH2	6.00	123.30	120.30
2	B	151	ARG	NE-CZ-NH2	5.52	123.06	120.30
2	B	102	ASP	N-CA-C	-5.50	96.17	111.00
2	D	102	ASP	N-CA-C	-5.13	97.15	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1283	0	1283	20	0
1	C	1254	0	1254	25	0
2	B	1638	0	1593	44	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1617	0	1557	39	0
3	B	14	0	13	2	0
3	D	14	0	13	0	0
4	A	46	0	0	0	0
4	B	108	0	0	2	0
4	C	37	0	0	0	0
4	D	69	0	0	0	0
All	All	6080	0	5713	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (124) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:23:ARG:HA	1:C:26:GLN:HE21	1.43	0.84
2:D:3:TYR:HB2	2:D:91:LEU:HD12	1.58	0.84
2:D:114:LEU:HD13	2:D:189:MET:HE3	1.61	0.81
2:B:5:PRO:CG	2:B:89:ASN:HB2	2.12	0.79
2:B:114:LEU:HD13	2:B:189:MET:HE3	1.67	0.75
1:A:23:ARG:HA	1:A:26:GLN:HE21	1.56	0.70
2:B:111:PRO:HD3	2:B:202:PRO:HD2	1.73	0.70
2:B:79:GLN:HE21	2:B:80:TYR:H	1.40	0.69
1:A:80:HIS:HE1	1:A:133:PRO:HG3	1.59	0.68
1:A:144:ALA:O	1:A:148:ARG:HG3	1.97	0.64
2:B:39:ILE:HG12	2:B:60:VAL:HG12	1.81	0.63
2:B:5:PRO:HG3	2:B:89:ASN:HB2	1.81	0.63
2:B:31:GLU:HB2	4:B:374:HOH:O	1.99	0.63
2:D:113:MET:O	2:D:135:TRP:HA	1.99	0.63
2:B:151:ARG:HG2	2:B:203:TRP:CZ3	2.34	0.62
2:B:15:MET:HB3	2:B:107:VAL:HA	1.84	0.58
2:B:207:LEU:HD13	2:B:209:LEU:HD11	1.87	0.56
1:C:57:ILE:HD11	1:C:141:PHE:HE1	1.70	0.56
1:C:125:LEU:HD12	1:C:127:MET:SD	2.45	0.56
2:D:190:ARG:HD2	2:D:203:TRP:CE2	2.40	0.56
2:B:190:ARG:HG3	2:B:200:TRP:CE3	2.41	0.56
2:B:190:ARG:HD2	2:B:203:TRP:CE2	2.42	0.55
1:C:144:ALA:O	1:C:148:ARG:HG3	2.06	0.55
2:D:184:VAL:HG22	2:D:210:ARG:HG2	1.88	0.55
1:C:10:LEU:HD23	1:C:125:LEU:HD11	1.88	0.55
2:D:139:LYS:O	2:D:142:GLU:HG2	2.06	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:113:ASP:OD2	2:B:72:ARG:NH2	2.41	0.54
2:D:127:GLN:O	2:D:212:THR:HB	2.08	0.54
2:D:3:TYR:H	2:D:4:PRO:HD3	1.72	0.54
1:A:80:HIS:CE1	1:A:133:PRO:HG3	2.42	0.54
2:B:90:MET:O	2:B:90:MET:HG2	2.07	0.54
2:B:116:ALA:HB2	2:B:209:LEU:HD21	1.88	0.53
2:B:151:ARG:HG2	2:B:203:TRP:CH2	2.43	0.53
1:C:71:GLN:O	1:C:72:LEU:HB2	2.09	0.53
1:A:123:GLU:HG2	1:A:128:ALA:HB2	1.91	0.52
1:C:64:SER:C	1:C:66:PRO:HD3	2.29	0.52
2:D:39:ILE:HG12	2:D:60:VAL:HG12	1.91	0.52
2:B:79:GLN:NE2	2:B:80:TYR:H	2.04	0.52
1:A:58:PRO:HD2	1:A:89:LEU:HD21	1.92	0.51
2:B:25:GLN:OE1	3:B:323:NAG:C6	2.58	0.51
2:D:190:ARG:HD2	2:D:203:TRP:CZ2	2.46	0.51
1:A:70:LEU:O	1:A:70:LEU:HD23	2.10	0.51
2:D:150:LEU:HD13	2:D:189:MET:HE2	1.93	0.51
2:B:182:ALA:HB1	2:B:183:PRO:HD2	1.92	0.50
1:C:39:THR:HG22	1:C:40:TYR:CE1	2.47	0.50
2:D:156:LEU:HD11	2:D:181:GLN:O	2.12	0.50
2:D:207:LEU:HD13	2:D:209:LEU:HD11	1.93	0.50
2:B:155:GLN:O	2:B:156:LEU:HD23	2.12	0.49
2:D:3:TYR:N	2:D:4:PRO:HD3	2.27	0.49
2:B:32:THR:O	2:B:33:HIS:HB2	2.12	0.49
2:D:190:ARG:HG3	2:D:200:TRP:CE3	2.48	0.49
2:D:40:LEU:HB3	2:D:59:CYS:HB3	1.95	0.49
1:A:46:GLU:HA	1:A:49:VAL:HG23	1.93	0.49
2:B:107:VAL:HB	4:B:334:HOH:O	2.13	0.48
1:C:142:ALA:HB3	1:C:146:GLN:OE1	2.13	0.48
2:B:190:ARG:HD2	2:B:203:TRP:CZ2	2.48	0.48
2:D:79:GLN:HE21	2:D:80:TYR:H	1.62	0.48
1:A:59:TRP:CZ2	1:A:61:PRO:HB3	2.49	0.48
1:A:55:LEU:HB2	1:A:57:ILE:HD13	1.95	0.48
2:B:149:GLU:HG3	2:B:200:TRP:CH2	2.47	0.48
1:C:68:GLN:C	1:C:70:LEU:N	2.67	0.48
2:D:150:LEU:HD13	2:D:189:MET:CE	2.44	0.47
1:C:23:ARG:HD2	2:D:143:TYR:O	2.13	0.47
1:C:74:GLY:O	1:C:77:SER:HB3	2.14	0.47
2:D:111:PRO:HD3	2:D:202:PRO:HD2	1.96	0.47
2:B:173:GLN:O	2:B:173:GLN:HG3	2.15	0.47
1:A:6:PRO:HA	1:A:175:PRO:OXT	2.16	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:25:GLN:OE1	3:B:323:NAG:H62	2.15	0.46
1:A:24:LYS:HE3	1:A:110:ASP:OD1	2.15	0.46
1:A:56:GLY:O	1:A:58:PRO:HD3	2.14	0.46
2:B:150:LEU:HD12	2:B:188:GLN:O	2.15	0.46
1:C:171:HIS:O	1:C:174:GLN:HG2	2.15	0.46
2:B:116:ALA:HB2	2:B:209:LEU:CD2	2.45	0.46
1:A:58:PRO:HB2	1:A:85:LEU:HD21	1.98	0.46
1:C:13:SER:O	1:C:17:LYS:HG3	2.15	0.46
2:D:39:ILE:CG2	2:D:58:ASP:HB3	2.46	0.45
1:C:46:GLU:HA	1:C:49:VAL:HG23	1.99	0.44
2:D:186:THR:C	2:D:187:LEU:HD12	2.38	0.44
2:D:20:ASN:HA	2:D:20:ASN:HD22	1.71	0.44
2:B:112:PRO:HG3	2:B:191:CYS:SG	2.56	0.44
2:D:110:GLU:HB2	2:D:138:TRP:CD2	2.52	0.44
2:B:139:LYS:O	2:B:142:GLU:HG2	2.18	0.44
2:B:73:LYS:HG3	2:B:74:ASN:ND2	2.31	0.44
2:D:117:LEU:HD22	2:D:173:GLN:CB	2.47	0.44
2:D:90:MET:O	2:D:90:MET:HG2	2.18	0.44
1:A:142:ALA:N	1:A:146:GLN:OE1	2.48	0.44
2:B:5:PRO:HD3	2:B:89:ASN:CB	2.48	0.44
2:D:184:VAL:CG2	2:D:210:ARG:HG2	2.48	0.44
1:C:68:GLN:C	1:C:70:LEU:H	2.20	0.44
1:C:138:MET:HA	1:C:139:PRO:HD2	1.92	0.43
2:D:151:ARG:HD2	2:D:161:TRP:CD2	2.53	0.43
2:B:45:SER:OG	2:B:49:CYS:HA	2.17	0.43
2:D:79:GLN:NE2	2:D:80:TYR:H	2.17	0.43
1:A:120:GLN:OE1	2:B:79:GLN:HG2	2.19	0.43
1:A:68:GLN:HB2	1:A:71:GLN:O	2.18	0.43
2:B:187:LEU:N	2:B:187:LEU:HD12	2.33	0.43
2:B:117:LEU:HA	2:B:117:LEU:HD23	1.74	0.43
2:D:183:PRO:O	2:D:211:PRO:HD2	2.19	0.43
2:B:192:ILE:HB	2:B:200:TRP:CE3	2.54	0.43
2:B:148:CYS:O	2:B:167:LEU:HB2	2.18	0.43
2:B:40:LEU:HB3	2:B:59:CYS:HB3	1.99	0.42
1:C:65:CYS:N	1:C:66:PRO:HD3	2.33	0.42
2:D:3:TYR:N	2:D:4:PRO:CD	2.83	0.42
2:B:5:PRO:HD3	2:B:89:ASN:HB3	2.02	0.42
1:C:150:GLY:O	1:C:154:VAL:HG23	2.20	0.42
1:C:43:CYS:HB2	1:C:44:HIS:CE1	2.55	0.42
2:B:153:GLN:HB3	2:B:161:TRP:CE3	2.55	0.42
2:D:127:GLN:HA	2:D:128:PRO:HD3	1.83	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:72:ARG:HA	2:D:75:LEU:HD12	2.02	0.41
1:A:172:LEU:O	1:A:175:PRO:HD3	2.20	0.41
2:B:151:ARG:NH1	2:B:188:GLN:OE1	2.51	0.41
1:C:10:LEU:HD23	1:C:125:LEU:CD1	2.50	0.41
1:C:57:ILE:HD11	1:C:141:PHE:CE1	2.51	0.41
1:A:22:VAL:O	1:A:26:GLN:HG3	2.20	0.41
2:D:207:LEU:HA	2:D:207:LEU:HD23	1.89	0.41
2:D:20:ASN:O	2:D:72:ARG:HB3	2.20	0.41
2:D:187:LEU:N	2:D:187:LEU:HD12	2.35	0.41
1:C:113:ASP:CG	2:D:76:LEU:HD12	2.41	0.41
2:B:5:PRO:HG2	2:B:89:ASN:HB2	1.95	0.41
2:D:112:PRO:HG3	2:D:191:CYS:SG	2.60	0.41
1:C:132:GLN:H	1:C:132:GLN:HG2	1.73	0.40
2:B:186:THR:HA	2:B:207:LEU:O	2.21	0.40
1:C:21:GLN:OE1	1:C:117:THR:HG21	2.22	0.40
2:D:210:ARG:HA	2:D:211:PRO:HD3	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	169/175 (97%)	155 (92%)	13 (8%)	1 (1%)	28  62 
1	C	167/175 (95%)	150 (90%)	11 (7%)	6 (4%)	4  13 
2	B	202/215 (94%)	191 (95%)	11 (5%)	0	100  100 
2	D	200/215 (93%)	185 (92%)	11 (6%)	4 (2%)	9  28 
All	All	738/780 (95%)	681 (92%)	46 (6%)	11 (2%)	12  37 

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	D	51	TYR
1	C	41	LYS
1	C	63	SER
2	D	159	ALA
1	A	41	LYS
1	C	64	SER
1	C	65	CYS
1	C	72	LEU
1	C	94	GLU
2	D	35	PRO
2	D	3	TYR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	136/142 (96%)	130 (96%)	6 (4%)	33 67
1	C	132/142 (93%)	126 (96%)	6 (4%)	32 66
2	B	186/195 (95%)	178 (96%)	8 (4%)	33 67
2	D	183/195 (94%)	170 (93%)	13 (7%)	17 44
All	All	637/674 (94%)	604 (95%)	33 (5%)	27 60

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	HIS
1	A	59	TRP
1	A	72	LEU
1	A	79	LEU
1	A	98	PRO
1	A	108	GLN
2	B	8	PRO
2	B	20	ASN
2	B	52	GLN
2	B	72	ARG
2	B	109	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	151	ARG
2	B	173	GLN
2	B	190	ARG
1	C	9	SER
1	C	44	HIS
1	C	51	LEU
1	C	79	LEU
1	C	108	GLN
1	C	174	GLN
2	D	7	SER
2	D	20	ASN
2	D	46	ARG
2	D	51	TYR
2	D	72	ARG
2	D	109	LEU
2	D	130	CYS
2	D	173	GLN
2	D	177	CYS
2	D	180	HIS
2	D	190	ARG
2	D	207	LEU
2	D	212	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	33	GLN
1	A	53	HIS
1	A	80	HIS
1	A	108	GLN
1	A	121	GLN
2	B	20	ASN
2	B	52	GLN
2	B	74	ASN
2	B	79	GLN
2	B	127	GLN
2	B	153	GLN
1	C	26	GLN
1	C	121	GLN
1	C	174	GLN
2	D	20	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	74	ASN
2	D	79	GLN
2	D	153	GLN
2	D	155	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	323	2	14,14,15	1.29	1 (7%)	15,19,21	1.31	1 (6%)
3	NAG	D	923	2	14,14,15	0.93	1 (7%)	15,19,21	0.75	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	323	2	-	0/6/23/26	0/1/1/1
3	NAG	D	923	2	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	923	NAG	C1-C2	2.66	1.56	1.52
3	B	323	NAG	C1-C2	4.02	1.58	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	323	NAG	O7-C7-C8	-2.55	117.42	122.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	323	NAG	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	171/175 (97%)	-0.25	10 (5%) 24 15	3, 20, 115, 192	0
1	C	169/175 (96%)	-0.26	3 (1%) 69 60	5, 24, 114, 146	0
2	B	206/215 (95%)	-0.52	1 (0%) 90 88	3, 16, 57, 95	0
2	D	204/215 (94%)	0.06	14 (6%) 18 10	4, 34, 97, 199	0
All	All	750/780 (96%)	-0.24	28 (3%) 42 31	3, 23, 99, 199	0

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	GLY	7.0
1	A	175	PRO	5.9
2	D	33	HIS	4.7
1	A	130	ALA	4.3
1	C	69	ALA	4.3
1	C	64	SER	3.9
2	B	118	ASP	3.8
1	A	134	THR	3.8
2	D	213	MET	3.3
2	D	118	ASP	2.9
2	D	29	GLY	2.9
2	D	159	ALA	2.8
2	D	2	GLY	2.7
1	A	6	PRO	2.7
2	D	3	TYR	2.6
1	A	136	GLY	2.6
2	D	132	TRP	2.4
1	C	66	PRO	2.4
1	A	129	PRO	2.3
1	A	131	LEU	2.2
2	D	39	ILE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	132	GLN	2.1
2	D	4	PRO	2.1
2	D	65	GLN	2.1
2	D	212	THR	2.1
1	A	71	GLN	2.0
2	D	117	LEU	2.0
2	D	173	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAG	B	323	14/15	0.80	0.14	-	63,87,127,128	0
3	NAG	D	923	14/15	0.51	0.27	-	70,103,108,112	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.