



# wwPDB X-ray Structure Validation Summary Report i

Feb 15, 2017 – 08:26 am GMT

PDB ID : 5CDI  
Title : Chloroplast chaperonin 60b1 of Chlamydomonas  
Authors : Zhang, S.; Zhou, H.; Yu, F.; Gao, F.; He, J.; Liu, C.  
Deposited on : 2015-07-04  
Resolution : 3.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 3.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1019 (4.08-3.52)
Clashscore	112137	1030 (4.04-3.56)
Ramachandran outliers	110173	1011 (4.06-3.54)
Sidechain outliers	110143	1005 (4.06-3.54)
RSRZ outliers	101464	1032 (4.08-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...



## 2 Entry composition i

There is only 1 type of molecule in this entry. The entry contains 55706 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chaperonin 60B1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	N	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	B	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	C	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	D	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	E	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	F	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	G	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	H	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	I	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	J	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	K	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	L	528	Total	C 3979	N 2471	O 696	S 788	24	0	0
1	M	528	Total	C 3979	N 2471	O 696	S 788	24	0	0

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP A8JE91

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	138	VAL	-	see sequence details	UNP A8JE91
N	1	MET	-	expression tag	UNP A8JE91
N	138	VAL	-	see sequence details	UNP A8JE91
B	1	MET	-	expression tag	UNP A8JE91
B	138	VAL	-	see sequence details	UNP A8JE91
C	1	MET	-	expression tag	UNP A8JE91
C	138	VAL	-	see sequence details	UNP A8JE91
D	1	MET	-	expression tag	UNP A8JE91
D	138	VAL	-	see sequence details	UNP A8JE91
E	1	MET	-	expression tag	UNP A8JE91
E	138	VAL	-	see sequence details	UNP A8JE91
F	1	MET	-	expression tag	UNP A8JE91
F	138	VAL	-	see sequence details	UNP A8JE91
G	1	MET	-	expression tag	UNP A8JE91
G	138	VAL	-	see sequence details	UNP A8JE91
H	1	MET	-	expression tag	UNP A8JE91
H	138	VAL	-	see sequence details	UNP A8JE91
I	1	MET	-	expression tag	UNP A8JE91
I	138	VAL	-	see sequence details	UNP A8JE91
J	1	MET	-	expression tag	UNP A8JE91
J	138	VAL	-	see sequence details	UNP A8JE91
K	1	MET	-	expression tag	UNP A8JE91
K	138	VAL	-	see sequence details	UNP A8JE91
L	1	MET	-	expression tag	UNP A8JE91
L	138	VAL	-	see sequence details	UNP A8JE91
M	1	MET	-	expression tag	UNP A8JE91
M	138	VAL	-	see sequence details	UNP A8JE91


### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Chaperonin 60B1




- Molecule 1: Chaperonin 60B1







- Molecule 1: Chaperonin 60B1




- Molecule 1: Chaperonin 60B1





- Molecule 1: Chaperonin 60B1









- Molecule 1: Chaperonin 60B1




- Molecule 1: Chaperonin 60B1




- Molecule 1: Chaperonin 60B1






- Molecule 1: Chaperonin 60B1



- Molecule 1: Chaperonin 60B1



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	143.34Å    174.39Å    213.68Å 90.00°    94.69°    90.00°	Depositor
Resolution (Å)	48.66 – 3.81 48.66 – 3.81	Depositor EDS
% Data completeness (in resolution range)	96.7 (48.66-3.81) 85.1 (48.66-3.81)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.50 (at 3.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
$R$ , $R_{free}$	0.275 , 0.325 0.294 , 0.340	Depositor DCC
$R_{free}$ test set	2005 reflections (2.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	133.5	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 114.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	55706	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	186.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.21	0/4011	0.40	0/5417
1	B	0.25	1/4011 (0.0%)	0.43	1/5417 (0.0%)
1	C	0.25	0/4011	0.44	3/5417 (0.1%)
1	D	0.23	0/4011	0.41	0/5417
1	E	0.32	3/4011 (0.1%)	0.49	4/5417 (0.1%)
1	F	0.23	0/4011	0.41	0/5417
1	G	0.25	1/4011 (0.0%)	0.40	0/5417
1	H	0.23	0/4011	0.43	2/5417 (0.0%)
1	I	0.22	0/4011	0.41	0/5417
1	J	0.24	0/4011	0.40	0/5417
1	K	0.21	0/4011	0.43	1/5417 (0.0%)
1	L	0.22	0/4011	0.41	0/5417
1	M	0.24	0/4011	0.40	0/5417
1	N	0.24	1/4011 (0.0%)	0.41	0/5417
All	All	0.24	6/56154 (0.0%)	0.42	11/75838 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
1	E	0	1
1	L	0	1
All	All	0	4

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	377	VAL	C-N	10.25	1.57	1.34
1	N	276	ALA	C-N	-5.80	1.20	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	206	VAL	C-N	-5.68	1.21	1.34
1	E	209	PRO	N-CD	5.45	1.55	1.47
1	B	280	PRO	N-CD	5.40	1.55	1.47

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	276	ALA	O-C-N	-7.55	110.61	122.70
1	H	276	ALA	O-C-N	-6.88	111.69	122.70
1	B	277	ILE	N-CA-C	6.17	127.65	111.00
1	C	277	ILE	O-C-N	-5.72	113.55	122.70
1	K	376	GLY	O-C-N	-5.68	113.61	122.70

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	279	ALA	Peptide
1	C	277	ILE	Mainchain
1	E	202	SER	Peptide
1	L	277	ILE	Mainchain

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3979	0	4104	88	0
1	B	3979	0	4104	88	0
1	C	3979	0	4103	82	0
1	D	3979	0	4104	94	0
1	E	3979	0	4104	212	0
1	F	3979	0	4104	87	0
1	G	3979	0	4103	82	0
1	H	3979	0	4103	75	0
1	I	3979	0	4104	71	0
1	J	3979	0	4104	81	0
1	K	3979	0	4104	71	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	3979	0	4104	71	0
1	M	3979	0	4103	69	0
1	N	3979	0	4103	75	0
All	All	55706	0	57451	1242	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 1242 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:205:PHE:CE2	1:E:267:LYS:CB	1.85	1.55
1:E:205:PHE:CE2	1:E:267:LYS:HB3	0.97	1.47
1:E:59:ARG:HH22	1:E:212:MET:CE	1.27	1.46
1:E:251:MET:HA	1:E:277:ILE:CG2	1.46	1.46
1:E:277:ILE:HD12	1:E:278:LYS:N	1.27	1.43

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	526/552 (95%)	480 (91%)	40 (8%)	6 (1%)	17 60
1	B	526/552 (95%)	473 (90%)	46 (9%)	7 (1%)	14 56
1	C	526/552 (95%)	479 (91%)	40 (8%)	7 (1%)	14 56
1	D	526/552 (95%)	479 (91%)	42 (8%)	5 (1%)	18 61
1	E	526/552 (95%)	472 (90%)	40 (8%)	14 (3%)	6 43
1	F	526/552 (95%)	480 (91%)	42 (8%)	4 (1%)	22 65
1	G	526/552 (95%)	483 (92%)	37 (7%)	6 (1%)	17 60

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	H	526/552 (95%)	478 (91%)	43 (8%)	5 (1%)	18 61
1	I	526/552 (95%)	482 (92%)	39 (7%)	5 (1%)	18 61
1	J	526/552 (95%)	480 (91%)	42 (8%)	4 (1%)	22 65
1	K	526/552 (95%)	478 (91%)	43 (8%)	5 (1%)	18 61
1	L	526/552 (95%)	478 (91%)	43 (8%)	5 (1%)	18 61
1	M	526/552 (95%)	475 (90%)	47 (9%)	4 (1%)	22 65
1	N	526/552 (95%)	480 (91%)	42 (8%)	4 (1%)	22 65
All	All	7364/7728 (95%)	6697 (91%)	586 (8%)	81 (1%)	17 60

5 of 81 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	158	ASP
1	A	173	THR
1	A	245	ASN
1	N	158	ASP
1	N	173	THR

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	433/445 (97%)	422 (98%)	11 (2%)	53 79
1	B	433/445 (97%)	424 (98%)	9 (2%)	59 82
1	C	433/445 (97%)	424 (98%)	9 (2%)	59 82
1	D	433/445 (97%)	422 (98%)	11 (2%)	53 79
1	E	433/445 (97%)	410 (95%)	23 (5%)	26 64
1	F	433/445 (97%)	422 (98%)	11 (2%)	53 79
1	G	433/445 (97%)	421 (97%)	12 (3%)	49 76
1	H	433/445 (97%)	421 (97%)	12 (3%)	49 76
1	I	433/445 (97%)	420 (97%)	13 (3%)	46 75

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	433/445 (97%)	422 (98%)	11 (2%)	53	79
1	K	433/445 (97%)	424 (98%)	9 (2%)	59	82
1	L	433/445 (97%)	420 (97%)	13 (3%)	46	75
1	M	433/445 (97%)	421 (97%)	12 (3%)	49	76
1	N	433/445 (97%)	425 (98%)	8 (2%)	64	85
All	All	6062/6230 (97%)	5898 (97%)	164 (3%)	50	77

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	235	ILE
1	G	435	THR
1	L	521	LEU
1	F	277	ILE
1	G	235	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	528/552 (95%)	0.40	47 (8%) 10 8	64, 175, 274, 319	0
1	B	528/552 (95%)	0.39	36 (6%) 18 13	49, 165, 283, 325	0
1	C	528/552 (95%)	0.39	39 (7%) 15 12	50, 166, 284, 318	0
1	D	528/552 (95%)	0.30	34 (6%) 20 14	55, 153, 262, 292	0
1	E	528/552 (95%)	0.28	25 (4%) 32 25	26, 159, 234, 301	0
1	F	528/552 (95%)	0.40	34 (6%) 20 14	69, 184, 271, 303	0
1	G	528/552 (95%)	0.55	52 (9%) 8 7	55, 193, 280, 315	0
1	H	528/552 (95%)	0.43	43 (8%) 13 10	57, 182, 288, 319	0
1	I	528/552 (95%)	0.53	63 (11%) 5 5	57, 192, 299, 325	0
1	J	528/552 (95%)	0.51	58 (10%) 6 6	84, 203, 315, 369	0
1	K	528/552 (95%)	0.64	69 (13%) 4 4	71, 189, 307, 340	0
1	L	528/552 (95%)	0.70	72 (13%) 3 4	70, 185, 310, 357	0
1	M	528/552 (95%)	0.70	80 (15%) 2 3	81, 207, 326, 366	0
1	N	528/552 (95%)	0.64	59 (11%) 6 6	77, 194, 294, 335	0
All	All	7392/7728 (95%)	0.49	711 (9%) 9 7	26, 178, 296, 369	0

The worst 5 of 711 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	302	VAL	18.9
1	C	302	VAL	12.1
1	G	316	ALA	10.3
1	L	225	ASP	9.8
1	M	362	GLU	9.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.