



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:44 am GMT

PDB ID : 4CFE
Title : Structure of full length human AMPK in complex with a small molecule activator, a benzimidazole derivative (991)
Authors : Xiao, B.; Sanders, M.J.; Carmena, D.; Bright, N.J.; Haire, L.F.; Underwood, E.; Patel, B.R.; Heath, R.B.; Walker, P.A.; Hallen, S.; Giordanetto, F.; Martin, S.R.; Carling, D.; Gamblin, S.J.
Deposited on : 2013-11-14
Resolution : 3.02 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.02 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1924 (3.04-3.00)
Clashscore	112137	2279 (3.04-3.00)
Ramachandran outliers	110173	2207 (3.04-3.00)
Sidechain outliers	110143	2210 (3.04-3.00)
RSRZ outliers	101464	1948 (3.04-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	571	<div> <div>62%</div> <div>12%</div> <div>25%</div> </div>
1	C	571	<div> <div>60%</div> <div>10%</div> <div>29%</div> </div>
2	B	286	<div> <div>49%</div> <div>9%</div> <div>41%</div> </div>
2	D	286	<div> <div>46%</div> <div>12%</div> <div>42%</div> </div>
3	E	331	<div> <div>75%</div> <div>13%</div> <div>11%</div> </div>
3	F	331	<div> <div>74%</div> <div>13%</div> <div>13%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	AMP	E	1326	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 14322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-AMP-ACTIVATED PROTEIN KINASE CATALYTIC SUB-UNIT ALPHA-2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	426	Total	C	N	O	P	S	3	0	0
			3411	2191	593	604	1	22			
1	C	404	Total	C	N	O	P	S	7	0	0
			3268	2101	568	576	1	22			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MET	-	EXPRESSION TAG	UNP P54646
A	-17	SER	-	EXPRESSION TAG	UNP P54646
A	-16	HIS	-	EXPRESSION TAG	UNP P54646
A	-15	HIS	-	EXPRESSION TAG	UNP P54646
A	-14	HIS	-	EXPRESSION TAG	UNP P54646
A	-13	HIS	-	EXPRESSION TAG	UNP P54646
A	-12	HIS	-	EXPRESSION TAG	UNP P54646
A	-11	HIS	-	EXPRESSION TAG	UNP P54646
A	-10	SER	-	EXPRESSION TAG	UNP P54646
A	-9	SER	-	EXPRESSION TAG	UNP P54646
A	-8	GLY	-	EXPRESSION TAG	UNP P54646
A	-7	LEU	-	EXPRESSION TAG	UNP P54646
A	-6	GLU	-	EXPRESSION TAG	UNP P54646
A	-5	VAL	-	EXPRESSION TAG	UNP P54646
A	-4	LEU	-	EXPRESSION TAG	UNP P54646
A	-3	PHE	-	EXPRESSION TAG	UNP P54646
A	-2	GLN	-	EXPRESSION TAG	UNP P54646
A	-1	GLY	-	EXPRESSION TAG	UNP P54646
A	0	PRO	-	EXPRESSION TAG	UNP P54646
C	-18	MET	-	EXPRESSION TAG	UNP P54646
C	-17	SER	-	EXPRESSION TAG	UNP P54646
C	-16	HIS	-	EXPRESSION TAG	UNP P54646
C	-15	HIS	-	EXPRESSION TAG	UNP P54646
C	-14	HIS	-	EXPRESSION TAG	UNP P54646

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-13	HIS	-	EXPRESSION TAG	UNP P54646
C	-12	HIS	-	EXPRESSION TAG	UNP P54646
C	-11	HIS	-	EXPRESSION TAG	UNP P54646
C	-10	SER	-	EXPRESSION TAG	UNP P54646
C	-9	SER	-	EXPRESSION TAG	UNP P54646
C	-8	GLY	-	EXPRESSION TAG	UNP P54646
C	-7	LEU	-	EXPRESSION TAG	UNP P54646
C	-6	GLU	-	EXPRESSION TAG	UNP P54646
C	-5	VAL	-	EXPRESSION TAG	UNP P54646
C	-4	LEU	-	EXPRESSION TAG	UNP P54646
C	-3	PHE	-	EXPRESSION TAG	UNP P54646
C	-2	GLN	-	EXPRESSION TAG	UNP P54646
C	-1	GLY	-	EXPRESSION TAG	UNP P54646
C	0	PRO	-	EXPRESSION TAG	UNP P54646

- Molecule 2 is a protein called 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT BETA-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	168	Total	C	N	O	P	S	0	0	0
			1336	863	224	243	1	5			
2	D	167	Total	C	N	O	P	S	0	0	0
			1322	855	221	241	1	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	MET	-	EXPRESSION TAG	UNP Q9Y478
B	-14	GLY	-	EXPRESSION TAG	UNP Q9Y478
B	-13	LEU	-	EXPRESSION TAG	UNP Q9Y478
B	-12	ASN	-	EXPRESSION TAG	UNP Q9Y478
B	-11	ASP	-	EXPRESSION TAG	UNP Q9Y478
B	-10	ILE	-	EXPRESSION TAG	UNP Q9Y478
B	-9	PHE	-	EXPRESSION TAG	UNP Q9Y478
B	-8	GLU	-	EXPRESSION TAG	UNP Q9Y478
B	-7	ALA	-	EXPRESSION TAG	UNP Q9Y478
B	-6	GLN	-	EXPRESSION TAG	UNP Q9Y478
B	-5	LYS	-	EXPRESSION TAG	UNP Q9Y478
B	-4	ILE	-	EXPRESSION TAG	UNP Q9Y478
B	-3	GLU	-	EXPRESSION TAG	UNP Q9Y478
B	-2	TRP	-	EXPRESSION TAG	UNP Q9Y478
B	-1	HIS	-	EXPRESSION TAG	UNP Q9Y478

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLU	-	EXPRESSION TAG	UNP Q9Y478
D	-15	MET	-	EXPRESSION TAG	UNP Q9Y478
D	-14	GLY	-	EXPRESSION TAG	UNP Q9Y478
D	-13	LEU	-	EXPRESSION TAG	UNP Q9Y478
D	-12	ASN	-	EXPRESSION TAG	UNP Q9Y478
D	-11	ASP	-	EXPRESSION TAG	UNP Q9Y478
D	-10	ILE	-	EXPRESSION TAG	UNP Q9Y478
D	-9	PHE	-	EXPRESSION TAG	UNP Q9Y478
D	-8	GLU	-	EXPRESSION TAG	UNP Q9Y478
D	-7	ALA	-	EXPRESSION TAG	UNP Q9Y478
D	-6	GLN	-	EXPRESSION TAG	UNP Q9Y478
D	-5	LYS	-	EXPRESSION TAG	UNP Q9Y478
D	-4	ILE	-	EXPRESSION TAG	UNP Q9Y478
D	-3	GLU	-	EXPRESSION TAG	UNP Q9Y478
D	-2	TRP	-	EXPRESSION TAG	UNP Q9Y478
D	-1	HIS	-	EXPRESSION TAG	UNP Q9Y478
D	0	GLU	-	EXPRESSION TAG	UNP Q9Y478

- Molecule 3 is a protein called 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT GAMMA-1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	294	Total	C	N	O	S	8	0	0
			2369	1541	393	428	7			
3	F	288	Total	C	N	O	S	15	0	0
			2315	1504	385	419	7			

- Molecule 4 is STAUROSPORINE (three-letter code: STU) (formula: C₂₈H₂₆N₄O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			35	28	4	3		
4	C	1	Total	C	N	O	0	0
			35	28	4	3		

- Molecule 5 is 5-[[6-CHLORANYL-5-(1-METHYLINDOL-5-YL)-1H-BENZIMIDAZOL-2-YL]OXY]-2-METHYL-BENZOIC ACID (three-letter code: 992) (formula: C₂₄H₁₈ClN₃O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	Cl	N	O	0
			31	24	1	3	3	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	Cl	N	O	0	0
			31	24	1	3	3		

- Molecule 6 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: $C_{10}H_{14}N_5O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
6	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
6	E	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
6	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
6	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		
6	F	1	Total	C	N	O	P	0	0
			23	10	5	7	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	13	Total 13 O 13	0	0
7	B	6	Total 6 O 6	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	6	Total 6	O 6	0	0
7	D	1	Total 1	O 1	0	0
7	E	2	Total 2	O 2	0	0
7	F	3	Total 3	O 3	0	0

AS51
ARG



• Molecule 2: 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT BETA-1



• Molecule 2: 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT BETA-1




• Molecule 3: 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT GAMMA-1



• Molecule 3: 5'-AMP-ACTIVATED PROTEIN KINASE SUBUNIT GAMMA-1





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.03Å 134.14Å 140.56Å 90.00° 92.42° 90.00°	Depositor
Resolution (Å)	19.91 – 3.02 19.91 – 3.02	Depositor EDS
% Data completeness (in resolution range)	92.0 (19.91-3.02) 92.0 (19.91-3.02)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.21 (at 3.04Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.218 , 0.253 0.216 , 0.253	Depositor DCC
R_{free} test set	2547 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	97.0	Xtriage
Anisotropy	0.303	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 67.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	14322	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, STU, 992, AMP, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/3478	0.36	0/4698
1	C	0.22	0/3333	0.37	0/4497
2	B	0.22	0/1361	0.43	1/1851 (0.1%)
2	D	0.23	0/1347	0.45	1/1834 (0.1%)
3	E	0.21	0/2418	0.40	0/3282
3	F	0.22	0/2360	0.39	0/3202
All	All	0.22	0/14297	0.39	2/19364 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	191(A)	PRO	N-CA-CB	6.22	110.77	103.30
2	B	191(B)	PRO	N-CA-CB	5.89	110.37	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3411	0	3398	43	0
1	C	3268	0	3277	35	0
2	B	1336	0	1322	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	1322	0	1300	17	0
3	E	2369	0	2444	25	0
3	F	2315	0	2398	19	0
4	A	35	0	26	2	0
4	C	35	0	26	1	0
5	A	31	0	17	2	0
5	C	31	0	17	2	0
6	E	69	0	36	2	0
6	F	69	0	36	0	0
7	A	13	0	0	0	0
7	B	6	0	0	0	0
7	C	6	0	0	1	0
7	D	1	0	0	0	0
7	E	2	0	0	0	0
7	F	3	0	0	0	0
All	All	14322	0	14297	146	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 146 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:430:TRP:HB3	1:C:440:VAL:HA	1.71	0.72
1:C:179:TYR:HA	1:C:202:VAL:HG11	1.77	0.67
1:C:158:PHE:O	1:C:160:LEU:N	2.28	0.66
1:C:423:MET:HB3	1:C:428:PHE:HB2	1.78	0.65
2:D:122:GLU:HG2	2:D:154:GLN:HG2	1.79	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/571 (73%)	394 (95%)	21 (5%)	0	100	100
1	C	393/571 (69%)	369 (94%)	23 (6%)	1 (0%)	44	80
2	B	161/286 (56%)	155 (96%)	5 (3%)	1 (1%)	28	68
2	D	160/286 (56%)	150 (94%)	8 (5%)	2 (1%)	14	49
3	E	290/331 (88%)	279 (96%)	7 (2%)	4 (1%)	13	47
3	F	282/331 (85%)	271 (96%)	10 (4%)	1 (0%)	38	76
All	All	1701/2376 (72%)	1618 (95%)	74 (4%)	9 (0%)	32	72

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	159	GLY
2	D	193(A)	VAL
2	B	191(B)	PRO
3	E	185	LYS
2	D	221	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/503 (73%)	356 (98%)	9 (2%)	53	84
1	C	355/503 (71%)	345 (97%)	10 (3%)	49	81
2	B	149/253 (59%)	144 (97%)	5 (3%)	42	78
2	D	146/253 (58%)	139 (95%)	7 (5%)	30	67
3	E	270/304 (89%)	260 (96%)	10 (4%)	39	75
3	F	265/304 (87%)	252 (95%)	13 (5%)	29	67
All	All	1550/2120 (73%)	1496 (96%)	54 (4%)	41	77

5 of 54 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	98	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	234	VAL
3	F	269	ARG
2	D	113	VAL
2	D	146	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	ASN
2	B	145	GLN
3	F	162	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	172	1	9,10,11	0.84	0	10,14,16	1.11	1 (10%)
2	SEP	B	108	2	9,9,10	0.87	0	9,12,14	1.09	0
1	TPO	C	172	1	9,10,11	0.86	0	10,14,16	1.17	1 (10%)
2	SEP	D	108	2	9,9,10	0.85	0	9,12,14	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	172	1	-	0/8/11/13	0/0/0/0
2	SEP	B	108	2	-	0/5/8/10	0/0/0/0
1	TPO	C	172	1	-	0/8/11/13	0/0/0/0
2	SEP	D	108	2	-	0/5/8/10	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	172	TPO	C-CA-N	2.12	114.14	109.86
1	C	172	TPO	C-CA-N	2.19	114.27	109.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	108	SEP	2	0
2	D	108	SEP	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	STU	A	1552	-	29,42,42	0.87	0	27,68,68	1.83	6 (22%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	992	A	1553	-	31,35,35	1.30	3 (9%)	33,52,52	1.16	2 (6%)
4	STU	C	1552	-	29,42,42	0.89	1 (3%)	27,68,68	1.86	7 (25%)
5	992	C	1553	-	31,35,35	1.29	3 (9%)	33,52,52	1.11	2 (6%)
6	AMP	E	1326	-	22,25,25	0.94	1 (4%)	24,38,38	1.58	2 (8%)
6	AMP	E	1327	-	22,25,25	0.95	1 (4%)	24,38,38	1.53	2 (8%)
6	AMP	E	1328	-	22,25,25	0.94	1 (4%)	24,38,38	1.59	2 (8%)
6	AMP	F	1326	-	22,25,25	0.93	1 (4%)	24,38,38	1.60	2 (8%)
6	AMP	F	1327	-	22,25,25	0.93	1 (4%)	24,38,38	1.59	2 (8%)
6	AMP	F	1328	-	22,25,25	0.93	1 (4%)	24,38,38	1.62	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	STU	A	1552	-	-	0/4/42/42	0/0/8/8
5	992	A	1553	-	-	0/6/12/12	0/5/5/5
4	STU	C	1552	-	-	0/4/42/42	0/0/8/8
5	992	C	1553	-	-	0/6/12/12	0/5/5/5
6	AMP	E	1326	-	-	0/6/26/26	0/3/3/3
6	AMP	E	1327	-	-	0/6/26/26	0/3/3/3
6	AMP	E	1328	-	-	0/6/26/26	0/3/3/3
6	AMP	F	1326	-	-	0/6/26/26	0/3/3/3
6	AMP	F	1327	-	-	0/6/26/26	0/3/3/3
6	AMP	F	1328	-	-	0/6/26/26	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1553	992	C9-N7	-2.91	1.32	1.37
5	C	1553	992	C9-N7	-2.89	1.32	1.37
4	C	1552	STU	C10-C11	-2.02	1.39	1.42
5	C	1553	992	O25-C23	2.82	1.40	1.35
5	A	1553	992	O25-C23	2.98	1.41	1.35

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	1328	AMP	N3-C2-N1	-6.10	123.54	128.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	F	1326	AMP	N3-C2-N1	-6.03	123.60	128.86
6	E	1326	AMP	N3-C2-N1	-5.94	123.68	128.86
6	F	1327	AMP	N3-C2-N1	-5.94	123.69	128.86
6	E	1328	AMP	N3-C2-N1	-5.91	123.71	128.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1552	STU	2	0
5	A	1553	992	2	0
4	C	1552	STU	1	0
5	C	1553	992	2	0
6	E	1326	AMP	1	0
6	E	1327	AMP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/571 (74%)	-0.12	7 (1%) 72 43	39, 77, 122, 161	1 (0%)
1	C	403/571 (70%)	-0.00	11 (2%) 55 26	44, 75, 134, 171	2 (0%)
2	B	167/286 (58%)	-0.16	4 (2%) 59 29	50, 71, 122, 146	0
2	D	166/286 (58%)	0.12	9 (5%) 26 10	41, 91, 149, 173	0
3	E	294/331 (88%)	-0.25	7 (2%) 59 29	43, 66, 117, 161	2 (0%)
3	F	288/331 (87%)	-0.25	5 (1%) 70 41	39, 64, 117, 161	3 (1%)
All	All	1743/2376 (73%)	-0.12	43 (2%) 58 28	39, 72, 130, 173	8 (0%)

The worst 5 of 43 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	148	THR	5.8
2	D	119	PRO	4.6
3	E	236	ARG	4.5
1	C	396	ALA	4.1
3	E	233	GLU	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	A	172	11/12	0.97	0.10	-	68,70,72,73	0
2	SEP	B	108	10/11	0.94	0.18	-	73,84,88,92	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	TPO	C	172	11/12	0.97	0.12	-	55,62,67,68	0
2	SEP	D	108	10/11	0.94	0.19	-	80,88,111,113	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	AMP	E	1326	23/23	0.82	0.29	2.73	127,147,161,165	0
5	992	A	1553	31/31	0.91	0.24	1.98	61,67,84,88	0
5	992	C	1553	31/31	0.91	0.21	0.46	73,75,80,82	0
4	STU	A	1552	35/35	0.96	0.19	0.03	37,44,48,87	0
6	AMP	F	1326	23/23	0.91	0.20	0.01	120,134,147,151	0
6	AMP	E	1327	23/23	0.95	0.18	-0.48	49,57,71,79	0
4	STU	C	1552	35/35	0.96	0.18	-0.49	36,43,51,52	0
6	AMP	F	1328	23/23	0.96	0.16	-0.81	57,62,67,69	0
6	AMP	E	1328	23/23	0.97	0.14	-0.93	53,68,75,77	0
6	AMP	F	1327	23/23	0.97	0.14	-1.59	74,84,90,101	0

6.5 Other polymers [i](#)

There are no such residues in this entry.