



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2017 – 11:09 AM EDT

PDB ID : 3CG0
Title : Crystal structure of signal receiver domain of modulated diguanylate cyclase from *Desulfovibrio desulfuricans* G20, an example of alternate folding
Authors : Patskovsky, Y.; Bonanno, J.B.; Romero, R.; Gilmore, M.; Chang, S.; Groshong, C.; Koss, J.; Wasserman, S.R.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : unknown
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	<div> <div>6%</div> <div> <div></div> <div>80%</div> <div>9%</div> <div>10%</div> </div> </div>
1	B	140	<div> <div>%</div> <div> <div></div> <div>81%</div> <div>9%</div> <div>11%</div> </div> </div>
1	C	140	<div> <div>4%</div> <div> <div></div> <div>74%</div> <div>13%</div> <div>10%</div> </div> </div>
1	D	140	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>11%</div> <div>10%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Response regulator receiver modulated diguanylate cyclase with PAS/PAC sensor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	0	5	0
			983	623	167	187	6			
1	B	125	Total	C	N	O	S	0	7	0
			986	626	171	184	5			
1	C	126	Total	C	N	O	S	0	3	0
			970	613	168	184	5			
1	D	126	Total	C	N	O	S	0	2	0
			960	607	165	182	6			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	EXPRESSION TAG	UNP Q311G8
A	0	SER	-	EXPRESSION TAG	UNP Q311G8
A	1	LEU	-	EXPRESSION TAG	UNP Q311G8
A	131	GLU	-	EXPRESSION TAG	UNP Q311G8
A	132	GLY	-	EXPRESSION TAG	UNP Q311G8
A	133	HIS	-	EXPRESSION TAG	UNP Q311G8
A	134	HIS	-	EXPRESSION TAG	UNP Q311G8
A	135	HIS	-	EXPRESSION TAG	UNP Q311G8
A	136	HIS	-	EXPRESSION TAG	UNP Q311G8
A	137	HIS	-	EXPRESSION TAG	UNP Q311G8
A	138	HIS	-	EXPRESSION TAG	UNP Q311G8
B	-1	MET	-	EXPRESSION TAG	UNP Q311G8
B	0	SER	-	EXPRESSION TAG	UNP Q311G8
B	1	LEU	-	EXPRESSION TAG	UNP Q311G8
B	131	GLU	-	EXPRESSION TAG	UNP Q311G8
B	132	GLY	-	EXPRESSION TAG	UNP Q311G8
B	133	HIS	-	EXPRESSION TAG	UNP Q311G8
B	134	HIS	-	EXPRESSION TAG	UNP Q311G8
B	135	HIS	-	EXPRESSION TAG	UNP Q311G8
B	136	HIS	-	EXPRESSION TAG	UNP Q311G8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	137	HIS	-	EXPRESSION TAG	UNP Q311G8
B	138	HIS	-	EXPRESSION TAG	UNP Q311G8
C	-1	MET	-	EXPRESSION TAG	UNP Q311G8
C	0	SER	-	EXPRESSION TAG	UNP Q311G8
C	1	LEU	-	EXPRESSION TAG	UNP Q311G8
C	131	GLU	-	EXPRESSION TAG	UNP Q311G8
C	132	GLY	-	EXPRESSION TAG	UNP Q311G8
C	133	HIS	-	EXPRESSION TAG	UNP Q311G8
C	134	HIS	-	EXPRESSION TAG	UNP Q311G8
C	135	HIS	-	EXPRESSION TAG	UNP Q311G8
C	136	HIS	-	EXPRESSION TAG	UNP Q311G8
C	137	HIS	-	EXPRESSION TAG	UNP Q311G8
C	138	HIS	-	EXPRESSION TAG	UNP Q311G8
D	-1	MET	-	EXPRESSION TAG	UNP Q311G8
D	0	SER	-	EXPRESSION TAG	UNP Q311G8
D	1	LEU	-	EXPRESSION TAG	UNP Q311G8
D	131	GLU	-	EXPRESSION TAG	UNP Q311G8
D	132	GLY	-	EXPRESSION TAG	UNP Q311G8
D	133	HIS	-	EXPRESSION TAG	UNP Q311G8
D	134	HIS	-	EXPRESSION TAG	UNP Q311G8
D	135	HIS	-	EXPRESSION TAG	UNP Q311G8
D	136	HIS	-	EXPRESSION TAG	UNP Q311G8
D	137	HIS	-	EXPRESSION TAG	UNP Q311G8
D	138	HIS	-	EXPRESSION TAG	UNP Q311G8



- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	29	Total O 29 29	0	0
2	B	21	Total O 21 21	0	0
2	C	22	Total O 22 22	0	0
2	D	19	Total O 19 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Response regulator receiver modulated diguanylate cyclase with PAS/PAC sensor



- Molecule 1: Response regulator receiver modulated diguanylate cyclase with PAS/PAC sensor





- Molecule 1: Response regulator receiver modulated diguanylate cyclase with PAS/PAC sensor



HIS

- Molecule 1: Response regulator receiver modulated diguanylate cyclase with PAS/PAC sensor



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	163.02Å 51.63Å 67.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.15 34.91 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.00-2.15) 99.0 (34.91-2.15)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.3.0034	Depositor
R, R_{free}	0.232 , 0.288 0.226 , 0.284	Depositor DCC
R_{free} test set	994 reflections (3.24%)	DCC
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.157	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3990	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.43	0/1012	0.63	0/1371
1	B	0.44	0/1020	0.69	1/1381 (0.1%)
1	C	0.42	0/992	0.68	1/1344 (0.1%)
1	D	0.45	0/979	0.69	0/1327
All	All	0.44	0/4003	0.67	2/5423 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	51	LEU	CA-CB-CG	5.96	129.01	115.30
1	B	62	LEU	CA-CB-CG	5.33	127.55	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	88	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	983	0	1006	5	0
1	B	986	0	1028	7	0
1	C	970	0	996	8	0
1	D	960	0	984	4	0
2	A	29	0	0	1	0
2	B	21	0	0	0	0
2	C	22	0	0	0	0
2	D	19	0	0	0	0
All	All	3990	0	4014	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70[A]:GLU:HG3	1:B:74[A]:ARG:HE	1.46	0.79
1:C:13:VAL:HG12	1:C:62:LEU:HD11	1.76	0.67
1:D:67:ASP:OD1	1:D:97:ARG:NH1	2.34	0.60
1:B:70[A]:GLU:HG3	1:B:74[A]:ARG:NE	2.16	0.59
1:B:70[A]:GLU:CG	1:B:74[A]:ARG:HE	2.20	0.54
1:D:84:ILE:HG21	1:D:107:LEU:HD22	1.91	0.53
1:A:117:HIS:HB2	1:B:30:LEU:HD13	1.91	0.50
1:C:15[A]:ASP:HB2	1:C:62:LEU:HD23	1.95	0.49
1:C:84:ILE:HG21	1:C:107:LEU:HD22	1.95	0.47
1:C:53:PRO:HG2	1:C:81:LEU:HD21	1.98	0.46
1:D:48:ALA:O	1:D:53:PRO:HD3	2.18	0.44
1:C:82:PRO:HG3	1:C:127:LYS:HD2	2.00	0.43
1:A:41:GLY:HA3	1:A:71:THR:OG1	2.19	0.43
1:A:24:ARG:NH2	2:A:164:HOH:O	2.52	0.42
1:A:109:LYS:HD2	1:B:86:ILE:HG22	2.01	0.41
1:C:13:VAL:HG12	1:C:62:LEU:CD1	2.46	0.41
1:D:7:LEU:HA	1:D:8:PRO:HD3	1.78	0.41
1:A:113:ALA:HB1	1:B:30:LEU:HD11	2.01	0.41
1:B:57[B]:LEU:HD23	1:B:84:ILE:HB	2.02	0.40
1:C:62:LEU:HD23	1:C:62:LEU:HA	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/140 (92%)	126 (98%)	3 (2%)	0	100	100
1	B	130/140 (93%)	125 (96%)	5 (4%)	0	100	100
1	C	127/140 (91%)	124 (98%)	3 (2%)	0	100	100
1	D	126/140 (90%)	123 (98%)	3 (2%)	0	100	100
All	All	512/560 (91%)	498 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/113 (94%)	97 (92%)	9 (8%)	12	7
1	B	107/113 (95%)	102 (95%)	5 (5%)	30	26
1	C	104/113 (92%)	87 (84%)	17 (16%)	3	1
1	D	103/113 (91%)	94 (91%)	9 (9%)	12	6
All	All	420/452 (93%)	380 (90%)	40 (10%)	9	5

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	24	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	35	LEU
1	A	52	ARG
1	A	55	ILE
1	A	81	LEU
1	A	89	SER
1	A	99	LYS
1	A	114	ASP
1	B	11	LEU
1	B	13	VAL
1	B	34	VAL
1	B	35	LEU
1	B	100	ARG
1	C	7	LEU
1	C	15[A]	ASP
1	C	15[B]	ASP
1	C	28	GLU
1	C	34	VAL
1	C	35	LEU
1	C	40	ASN
1	C	43	GLU
1	C	51	LEU
1	C	61	MET
1	C	81	LEU
1	C	93	GLU
1	C	99	LYS
1	C	101	VAL
1	C	107	LEU
1	C	109	LYS
1	C	114	ASP
1	D	6	ASP
1	D	17	ARG
1	D	43	GLU
1	D	51	LEU
1	D	89	SER
1	D	107	LEU
1	D	114	ASP
1	D	124	ILE
1	D	127	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	102	ASN
1	D	80	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	126/140 (90%)	0.18	8 (6%) 21 26	28, 48, 78, 94	0
1	B	125/140 (89%)	0.06	2 (1%) 72 77	26, 46, 77, 86	0
1	C	126/140 (90%)	0.27	5 (3%) 39 46	32, 52, 81, 108	0
1	D	126/140 (90%)	0.25	7 (5%) 25 33	31, 54, 90, 107	0
All	All	503/560 (89%)	0.19	22 (4%) 35 43	26, 50, 82, 108	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	GLU	6.5
1	A	92	VAL	5.0
1	D	7	LEU	4.2
1	D	6	ASP	4.1
1	A	7	LEU	3.6
1	B	7	LEU	3.3
1	A	91[A]	ASP	3.3
1	D	100	ARG	3.2
1	B	77	ALA	3.1
1	A	89	SER	3.1
1	C	6	ASP	2.9
1	D	93	GLU	2.8
1	D	63	CYS	2.7
1	C	62	LEU	2.7
1	D	90	GLN	2.7
1	A	6	ASP	2.5
1	D	96	GLN	2.4
1	A	52	ARG	2.3
1	C	90	GLN	2.3
1	C	50	ASP	2.3
1	A	125[A]	HIS	2.2
1	C	63	CYS	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.