



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 10:29 am GMT

PDB ID : 3CGD
Title : Pyridine Nucleotide Complexes with Bacillus anthracis Coenzyme A-Disulfide Reductase: A Structural Analysis of Dual NAD(P)H Specificity
Authors : Wallen, J.R.
Deposited on : 2008-03-05
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1062 (2.26-2.26)
Clashscore	112137	1178 (2.26-2.26)
Ramachandran outliers	110173	1145 (2.26-2.26)
Sidechain outliers	110143	1146 (2.26-2.26)
RSRZ outliers	101464	1066 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	480	<div> <div>13%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>•</div> <div>8%</div> </div> </div>
1	B	480	<div> <div>11%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>8%</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7735 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyridine nucleotide-disulfide oxidoreductase, class I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	3	0
			3496	2208	603	670	15			
1	B	444	Total	C	N	O	S	0	2	0
			3489	2203	603	668	15			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-35	MET	-	EXPRESSION TAG	UNP Q81TK8
A	-34	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	-33	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	-32	SER	-	EXPRESSION TAG	UNP Q81TK8
A	-31	HIS	-	EXPRESSION TAG	UNP Q81TK8
A	-30	HIS	-	EXPRESSION TAG	UNP Q81TK8
A	-29	HIS	-	EXPRESSION TAG	UNP Q81TK8
A	-28	HIS	-	EXPRESSION TAG	UNP Q81TK8
A	-27	HIS	-	EXPRESSION TAG	UNP Q81TK8
A	-26	HIS	-	EXPRESSION TAG	UNP Q81TK8
A	-25	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	-24	MET	-	EXPRESSION TAG	UNP Q81TK8
A	-23	ALA	-	EXPRESSION TAG	UNP Q81TK8
A	-22	SER	-	EXPRESSION TAG	UNP Q81TK8
A	-21	MET	-	EXPRESSION TAG	UNP Q81TK8
A	-20	THR	-	EXPRESSION TAG	UNP Q81TK8
A	-19	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	-18	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	-17	GLN	-	EXPRESSION TAG	UNP Q81TK8
A	-16	GLN	-	EXPRESSION TAG	UNP Q81TK8
A	-15	MET	-	EXPRESSION TAG	UNP Q81TK8
A	-14	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	-13	ARG	-	EXPRESSION TAG	UNP Q81TK8
A	-12	THR	-	EXPRESSION TAG	UNP Q81TK8
A	-11	LEU	-	EXPRESSION TAG	UNP Q81TK8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	TYR	-	EXPRESSION TAG	UNP Q81TK8
A	-9	ASP	-	EXPRESSION TAG	UNP Q81TK8
A	-8	ASP	-	EXPRESSION TAG	UNP Q81TK8
A	-7	ASP	-	EXPRESSION TAG	UNP Q81TK8
A	-6	ASP	-	EXPRESSION TAG	UNP Q81TK8
A	-5	LYS	-	EXPRESSION TAG	UNP Q81TK8
A	-4	ASP	-	EXPRESSION TAG	UNP Q81TK8
A	-3	ARG	-	EXPRESSION TAG	UNP Q81TK8
A	-2	TRP	-	EXPRESSION TAG	UNP Q81TK8
A	-1	GLY	-	EXPRESSION TAG	UNP Q81TK8
A	0	SER	-	EXPRESSION TAG	UNP Q81TK8
B	-35	MET	-	EXPRESSION TAG	UNP Q81TK8
B	-34	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	-33	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	-32	SER	-	EXPRESSION TAG	UNP Q81TK8
B	-31	HIS	-	EXPRESSION TAG	UNP Q81TK8
B	-30	HIS	-	EXPRESSION TAG	UNP Q81TK8
B	-29	HIS	-	EXPRESSION TAG	UNP Q81TK8
B	-28	HIS	-	EXPRESSION TAG	UNP Q81TK8
B	-27	HIS	-	EXPRESSION TAG	UNP Q81TK8
B	-26	HIS	-	EXPRESSION TAG	UNP Q81TK8
B	-25	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	-24	MET	-	EXPRESSION TAG	UNP Q81TK8
B	-23	ALA	-	EXPRESSION TAG	UNP Q81TK8
B	-22	SER	-	EXPRESSION TAG	UNP Q81TK8
B	-21	MET	-	EXPRESSION TAG	UNP Q81TK8
B	-20	THR	-	EXPRESSION TAG	UNP Q81TK8
B	-19	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	-18	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	-17	GLN	-	EXPRESSION TAG	UNP Q81TK8
B	-16	GLN	-	EXPRESSION TAG	UNP Q81TK8
B	-15	MET	-	EXPRESSION TAG	UNP Q81TK8
B	-14	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	-13	ARG	-	EXPRESSION TAG	UNP Q81TK8
B	-12	THR	-	EXPRESSION TAG	UNP Q81TK8
B	-11	LEU	-	EXPRESSION TAG	UNP Q81TK8
B	-10	TYR	-	EXPRESSION TAG	UNP Q81TK8
B	-9	ASP	-	EXPRESSION TAG	UNP Q81TK8
B	-8	ASP	-	EXPRESSION TAG	UNP Q81TK8
B	-7	ASP	-	EXPRESSION TAG	UNP Q81TK8
B	-6	ASP	-	EXPRESSION TAG	UNP Q81TK8
B	-5	LYS	-	EXPRESSION TAG	UNP Q81TK8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	ASP	-	EXPRESSION TAG	UNP Q81TK8
B	-3	ARG	-	EXPRESSION TAG	UNP Q81TK8
B	-2	TRP	-	EXPRESSION TAG	UNP Q81TK8
B	-1	GLY	-	EXPRESSION TAG	UNP Q81TK8
B	0	SER	-	EXPRESSION TAG	UNP Q81TK8

- Molecule 2 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	16	3	0	0
2	B	1	48	21	7	16	3	0	0

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
3	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 4 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	229	Total	O	0	0
			229	229		
5	B	231	Total	O	0	0
			231	231		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyridine nucleotide-disulfide oxidoreductase, class I



- Molecule 1: Pyridine nucleotide-disulfide oxidoreductase, class I



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	169.61Å 81.17Å 98.05Å 90.00° 104.01° 90.00°	Depositor
Resolution (Å)	37.45 – 2.25 37.44 – 2.25	Depositor EDS
% Data completeness (in resolution range)	96.0 (37.45-2.25) 96.0 (37.44-2.25)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.206 , 0.248 0.206 , 0.244	Depositor DCC
R_{free} test set	3006 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	32.7	Xtriage
Anisotropy	0.611	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7735	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, FAD, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3563	0.59	0/4812
1	B	0.45	0/3553	0.59	0/4799
All	All	0.46	0/7116	0.59	0/9611

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	425	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3496	0	3527	20	0
1	B	3489	0	3519	18	0
2	A	48	0	32	0	0
2	B	48	0	32	0	0
3	A	53	0	31	0	0
3	B	53	0	31	0	0
4	A	44	0	26	1	0
4	B	44	0	26	2	0
5	A	229	0	0	4	0
5	B	231	0	0	2	0
All	All	7735	0	7224	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:LYS:HE2	5:A:977:HOH:O	1.60	1.01
1:B:137:ASP:OD1	1:B:140:ARG:NH2	2.36	0.58
1:A:244:VAL:HG13	4:A:818:NAD:H51N	1.90	0.53
1:A:395:GLU:HG2	5:A:833:HOH:O	2.09	0.53
1:A:288:HIS:HB3	1:A:291:LYS:HB2	1.92	0.52
1:A:5:ILE:HB	1:A:30:THR:HG22	1.91	0.52
1:A:120:GLU:HA	1:A:124:ARG:HD2	1.92	0.51
1:B:5:ILE:HB	1:B:30:THR:HG22	1.93	0.50
1:A:359:VAL:HG12	1:A:361:SER:HB2	1.94	0.50
1:A:114:VAL:HG13	1:A:282:ASP:HB3	1.94	0.49
1:B:156:ILE:HD12	1:B:214:VAL:HG21	1.94	0.49
1:B:372:LYS:HD3	1:B:395:GLU:OE2	2.12	0.48
1:A:384:THR:O	1:A:385:LYS:HB2	2.14	0.48
1:A:220:ASN:H	1:A:220:ASN:ND2	2.11	0.47
1:B:181:ARG:HE	1:B:182:ASN:ND2	2.12	0.47
1:A:200:GLU:O	1:A:204:HIS:HD2	1.97	0.47
1:B:254:THR:O	1:B:255[B]:ASN:HB3	2.15	0.46
1:B:163:LEU:HD21	1:B:185:ILE:HD12	1.97	0.45
1:A:104:SER:HB2	5:A:979:HOH:O	2.15	0.45
1:B:412:LYS:HA	5:B:911:HOH:O	2.17	0.45
1:A:220:ASN:CG	1:A:221:GLU:H	2.19	0.45
1:B:187:THR:O	1:B:345:LYS:HE3	2.17	0.45
1:B:288:HIS:HB3	1:B:291:LYS:HB2	1.99	0.44
1:A:46:TYR:HB2	1:A:52:ILE:HD12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:114:VAL:HG12	1:B:246:PRO:HA	2.00	0.44
1:B:22:ASN:HB2	1:B:315:LEU:HD13	2.00	0.44
1:A:116:PRO:HB2	1:A:131:LEU:HD22	2.00	0.44
1:B:161:ILE:CG1	4:B:819:NAD:H6N	2.48	0.43
1:B:33:LYS:HE2	1:B:33:LYS:HB3	1.79	0.43
1:A:370:ASN:O	1:A:372:LYS:HG2	2.19	0.42
1:B:73:ASP:OD1	1:B:75:LYS:HE3	2.18	0.42
1:A:345:LYS:HE3	5:A:970:HOH:O	2.19	0.42
1:A:44:LEU:HD22	1:A:138:ALA:HB2	2.02	0.42
1:B:54:SER:HB2	5:B:912:HOH:O	2.20	0.42
1:B:161:ILE:HG12	4:B:819:NAD:H6N	2.03	0.41
1:B:345:LYS:HG2	1:B:345:LYS:H	1.67	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/480 (93%)	429 (96%)	15 (3%)	1 (0%)	51	60
1	B	444/480 (92%)	425 (96%)	18 (4%)	1 (0%)	51	60
All	All	889/960 (93%)	854 (96%)	33 (4%)	2 (0%)	51	60

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	38	SER
1	B	38	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/398 (94%)	352 (94%)	21 (6%)	25	25
1	B	372/398 (94%)	355 (95%)	17 (5%)	31	35
All	All	745/796 (94%)	707 (95%)	38 (5%)	30	30

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	GLU
1	A	143	LYS
1	A	148	ASN
1	A	163	LEU
1	A	175	LYS
1	A	183	ASP
1	A	210	THR
1	A	221	GLU
1	A	241	SER
1	A	249[A]	ASP
1	A	249[B]	ASP
1	A	252	GLU
1	A	265[A]	GLU
1	A	265[B]	GLU
1	A	293	ILE
1	A	309	LEU
1	A	339	ARG
1	A	385	LYS
1	A	396	GLU
1	A	401	ARG
1	A	442	ARG
1	B	62	ASN
1	B	120	GLU
1	B	143	LYS
1	B	149	LYS
1	B	210	THR
1	B	211	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	223	VAL
1	B	249[A]	ASP
1	B	249[B]	ASP
1	B	339	ARG
1	B	357	VAL
1	B	372	LYS
1	B	396	GLU
1	B	401	ARG
1	B	412	LYS
1	B	414	SER
1	B	442	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	A	184	HIS
1	B	62	ASN
1	B	182	ASN
1	B	184	HIS
1	B	334	ASN
1	B	370	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COA	A	445	-	43,50,50	0.65	0	48,75,75	1.75	7 (14%)
3	FAD	A	446	-	51,58,58	0.99	4 (7%)	54,89,89	2.05	7 (12%)
4	NAD	A	818	-	41,48,48	1.65	3 (7%)	43,73,73	1.72	1 (2%)
2	COA	B	445	-	43,50,50	0.68	0	48,75,75	1.76	6 (12%)
3	FAD	B	446	-	51,58,58	0.97	2 (3%)	54,89,89	2.05	8 (14%)
4	NAD	B	819	-	41,48,48	1.65	3 (7%)	43,73,73	1.72	2 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	445	-	-	0/44/64/64	0/3/3/3
3	FAD	A	446	-	-	0/28/50/50	0/6/6/6
4	NAD	A	818	-	-	0/22/62/62	0/5/5/5
2	COA	B	445	-	-	0/44/64/64	0/3/3/3
3	FAD	B	446	-	-	0/28/50/50	0/6/6/6
4	NAD	B	819	-	-	0/22/62/62	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	446	FAD	C10-N1	2.15	1.36	1.33
3	B	446	FAD	C4X-N5	2.32	1.36	1.33
3	A	446	FAD	O4B-C1B	2.34	1.44	1.41
3	A	446	FAD	C4X-N5	2.48	1.36	1.33
4	B	819	NAD	C2A-N1A	2.65	1.38	1.33
4	A	818	NAD	C2A-N1A	2.66	1.38	1.33
3	A	446	FAD	C9A-N10	3.37	1.43	1.38
3	B	446	FAD	C9A-N10	3.65	1.43	1.38
4	B	819	NAD	C2A-N3A	4.00	1.38	1.32
4	A	818	NAD	C2A-N3A	4.33	1.39	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	818	NAD	C7N-N7N	7.82	1.48	1.33
4	B	819	NAD	C7N-N7N	7.99	1.48	1.33

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	818	NAD	N3A-C2A-N1A	-9.96	120.19	128.86
4	B	819	NAD	N3A-C2A-N1A	-9.95	120.19	128.86
3	A	446	FAD	N3A-C2A-N1A	-9.72	120.39	128.86
3	B	446	FAD	N3A-C2A-N1A	-8.87	121.13	128.86
2	B	445	COA	N3A-C2A-N1A	-8.67	121.31	128.86
2	A	445	COA	N3A-C2A-N1A	-8.31	121.62	128.86
2	A	445	COA	C6P-C7P-N8P	-3.41	104.81	111.87
3	B	446	FAD	C4X-C4-N3	-2.87	119.40	123.48
2	A	445	COA	C2P-C3P-N4P	-2.87	106.26	112.50
2	B	445	COA	C2P-C3P-N4P	-2.73	106.56	112.50
2	B	445	COA	C6P-C7P-N8P	-2.70	106.28	111.87
3	A	446	FAD	C4X-C4-N3	-2.70	119.64	123.48
2	B	445	COA	C7P-C6P-C5P	-2.59	108.05	112.22
3	B	446	FAD	C4A-C5A-N7A	-2.50	106.99	109.41
2	A	445	COA	C7P-C6P-C5P	-2.27	108.57	112.22
3	A	446	FAD	C4X-C10-N10	-2.22	118.98	120.52
4	B	819	NAD	C1B-N9A-C4A	-2.08	123.05	126.64
3	B	446	FAD	C4-C4X-N5	2.07	120.94	118.68
2	A	445	COA	C3P-N4P-C5P	2.07	126.83	122.84
3	A	446	FAD	C4-C4X-N5	2.17	121.06	118.68
3	B	446	FAD	C5X-C9A-N10	2.32	119.38	117.66
2	B	445	COA	C7P-N8P-C9P	2.33	126.93	122.59
3	A	446	FAD	C1'-N10-C9A	2.36	120.51	118.35
2	A	445	COA	O6A-CCP-CBP	2.37	114.35	110.55
2	B	445	COA	C3P-N4P-C5P	2.64	127.91	122.84
2	A	445	COA	C7P-N8P-C9P	2.70	127.62	122.59
3	B	446	FAD	C1'-N10-C9A	4.38	122.36	118.35
3	B	446	FAD	C4X-N5-C5X	5.13	122.18	116.76
3	A	446	FAD	C4X-N5-C5X	5.35	122.41	116.76
3	A	446	FAD	C4-N3-C2	6.59	120.92	115.16
3	B	446	FAD	C4-N3-C2	6.77	121.08	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	818	NAD	1	0
4	B	819	NAD	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	444/480 (92%)	0.95	62 (13%) 3 3	39, 46, 55, 62	0
1	B	444/480 (92%)	0.92	53 (11%) 5 5	40, 46, 54, 61	0
All	All	888/960 (92%)	0.93	115 (12%) 4 4	39, 46, 55, 62	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	147	THR	7.8
1	B	25	ASN	5.4
1	B	370	ASN	5.4
1	A	25	ASN	5.1
1	A	96	LYS	4.7
1	A	198	TYR	4.6
1	A	370	ASN	4.6
1	A	147	THR	4.5
1	A	221	GLU	4.4
1	B	260	HIS	4.4
1	A	150	VAL	4.4
1	A	95	THR	4.3
1	B	150	VAL	4.3
1	A	260	HIS	4.1
1	B	220	ASN	4.1
1	B	24	GLU	4.0
1	B	221	GLU	4.0
1	B	369	PRO	3.8
1	A	62	ASN	3.6
1	A	24	GLU	3.6
1	B	11	ALA	3.5
1	A	127	GLN	3.4
1	A	187	THR	3.4
1	B	198	TYR	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	69	LYS	3.3
1	A	206	ILE	3.3
1	A	128	GLY	3.2
1	B	62	ASN	3.2
1	B	10	ALA	3.2
1	B	222	ARG	3.2
1	A	255	ASN	3.1
1	A	149	LYS	3.1
1	B	144	THR	3.0
1	B	423	LEU	3.0
1	A	148	ASN	3.0
1	A	329	ILE	2.9
1	B	6	ILE	2.9
1	B	280	ALA	2.9
1	A	86	GLU	2.9
1	B	223	VAL	2.9
1	B	299	ILE	2.8
1	B	329	ILE	2.8
1	B	145	LEU	2.8
1	B	86	GLU	2.8
1	A	181	ARG	2.8
1	B	51	ALA	2.8
1	B	143	LYS	2.8
1	A	11	ALA	2.7
1	A	146	GLU	2.7
1	B	191	GLY	2.7
1	B	161	ILE	2.7
1	A	52	ILE	2.7
1	A	299	ILE	2.7
1	A	101	PHE	2.7
1	A	152	ASP	2.6
1	B	253	GLY	2.6
1	A	423	LEU	2.6
1	B	224	GLU	2.6
1	A	280	ALA	2.6
1	A	334	ASN	2.6
1	B	255[A]	ASN	2.6
1	A	56	GLU	2.6
1	B	211	ASN	2.6
1	A	69	LYS	2.6
1	B	330	ILE	2.5
1	B	127	GLN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	222	ARG	2.5
1	A	161	ILE	2.5
1	A	143	LYS	2.5
1	B	184	HIS	2.5
1	B	111	ALA	2.5
1	A	10	ALA	2.5
1	B	95	THR	2.5
1	A	31	LEU	2.4
1	A	171	GLU	2.4
1	A	425	TYR	2.4
1	A	163	LEU	2.4
1	B	183	ASP	2.4
1	B	101	PHE	2.4
1	B	210	THR	2.4
1	B	149	LYS	2.3
1	B	148	ASN	2.3
1	A	183	ASP	2.3
1	A	432	VAL	2.3
1	A	224	GLU	2.3
1	A	184	HIS	2.3
1	A	197	ILE	2.3
1	B	13	MET	2.3
1	B	31	LEU	2.3
1	A	173	GLY	2.3
1	A	192	ASP	2.3
1	B	192	ASP	2.2
1	B	433	TRP	2.2
1	A	371	ALA	2.2
1	A	6	ILE	2.2
1	B	187	THR	2.2
1	B	302	THR	2.2
1	A	397	GLY	2.2
1	B	146	GLU	2.2
1	A	217	PHE	2.2
1	A	191	GLY	2.2
1	A	302	THR	2.1
1	B	432	VAL	2.1
1	A	252	GLU	2.1
1	B	63	VAL	2.1
1	A	53	ALA	2.1
1	A	182	ASN	2.0
1	B	2	ASN	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	111	ALA	2.0
1	A	253	GLY	2.0
1	B	173	GLY	2.0
1	A	48	ILE	2.0
1	A	102	GLU	2.0
1	A	301	THR	2.0
1	A	354	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	NAD	A	818	44/44	0.90	0.19	-0.21	53,57,62,62	0
2	COA	B	445	48/48	0.89	0.19	-0.54	48,58,69,70	0
2	COA	A	445	48/48	0.89	0.17	-0.72	50,59,70,70	0
4	NAD	B	819	44/44	0.92	0.16	-0.91	53,56,60,60	0
3	FAD	B	446	53/53	0.93	0.15	-1.47	36,43,52,52	0
3	FAD	A	446	53/53	0.94	0.13	-1.65	37,45,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.