



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 15, 2017 – 03:42 am GMT

PDB ID : 4COX
Title : CYCLOOXYGENASE-2 (PROSTAGLANDIN SYNTHASE-2) COM-
PLEXED WITH A NON-SELECTIVE INHIBITOR, INDOMETHACIN
Authors : Kurumbail, R.; Stallings, W.
Deposited on : 1996-12-18
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	587	
1	B	587	
1	C	587	
1	D	587	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	681	-	-	-	X
2	NAG	C	661	-	-	-	X
2	NAG	C	681	-	-	-	X
4	IMN	B	701	-	-	X	X
4	IMN	D	701	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18332 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLOOXYGENASE-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	B	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	C	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			
1	D	552	Total	C	N	O	S	0	0	0
			4473	2886	748	814	25			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	310	GLN	ASN	CONFLICT	UNP Q05769
A	333	LYS	ARG	CONFLICT	UNP Q05769
B	310	GLN	ASN	CONFLICT	UNP Q05769
B	333	LYS	ARG	CONFLICT	UNP Q05769
C	310	GLN	ASN	CONFLICT	UNP Q05769
C	333	LYS	ARG	CONFLICT	UNP Q05769
D	310	GLN	ASN	CONFLICT	UNP Q05769
D	333	LYS	ARG	CONFLICT	UNP Q05769

- Molecule 2 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 4 is INDOMETHACIN (three-letter code: IMN) (formula: $C_{19}H_{16}ClNO_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			25	19	1	1	4		
4	B	1	Total	C	Cl	N	O	0	0
			25	19	1	1	4		
4	C	1	Total	C	Cl	N	O	0	0
			25	19	1	1	4		
4	D	1	Total	C	Cl	N	O	0	0
			25	19	1	1	4		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYCLOOXYGENASE-2




• Molecule 1: CYCLOOXYGENASE-2





• Molecule 1: CYCLOOXYGENASE-2



• Molecule 1: CYCLOOXYGENASE-2



F556	K557	I558	A562	S563	I564	I568	V572	K573	G574	C575	P576	F577	T578	S579	F580	H581	V582	Q583	ASP	PRO	GLN	PRO	THR	LYS	THR	ALA	THR	ILE	ASN	ALA	SER	ALA	SER	HIS	SER	ARG	LEU	ASP	ASP	ILE	ASN	PRO	THR	VAL	LEU	ILE	LYS	ARG	ARG	SER	THR	GLU	LEU		
E416	H417	G418	L419	T420	Q421	E422	V423	L424	E424	T427	R428	Q429	I430	A431	R432	R433	V434	R438	M439	V440	P441	I442	A443	V444	Q445	A446	V447	A450	S451	L452	D453	Q454	S455	R456	E457	M458	K459	S462	L463	M464	E465	Y466	R467	K468	R469	F470	S471	L472	Y475	T476	S477	F478	E479	V480	L481
K342	T345	Y348	Y349	Q350	H351	L352	S353	G354	G354	F357	K358	L366	F367	A368	Q369	Q372	Y373	Q374	H375	R376	R377	A378	S379	E380	F381	N382	L383	L384	Y385	H386	W387	H388	P389	L390	L391	T394	F395	N396	I397	E398	D399	Q400	E401	Y402	K405	Y409	M410	N411	S412	I413	L414	L415			
T269	Q270	V271	Y275	Y276	P277	H278	L279	P280	E281	N282	L283	Q284	F285	A286	V287	G288	Q289	E290	V291	F292	G293	L294	L298	M299	M300	T303	L306	R307	E308	H309	R310	R311	T315	L316	E319	H320	P321	E322	D325	E326	F329	Q330	K333	L336	L337	Y338	G338	E339	T340	L341					
D190	P191	Q192	N195	Y196	F198	A199	T206	S126	P127	F209	R210	K211	T212	D213	H214	K215	R216	G217	P218	Y136	K137	S138	F142	L230	N231	H232	I233	T234	G235	E236	T237	L238	Q241	H242	K243	R244	R245	M163	G164	V165	E170	D173	L178	E179	V180	V181	G259	E260	L182	Y261	Y262	V266	K267	D268	
A33	N34	C37	S38	N39	P40	G41	Q42	N43	R44	G45	E46	S49	T50	K56	C57	F64	C69	T70	T71	P72	E73	F74	L75	T76	R77	I78	L82	K83	P84	T85	T88	V89	H90	I91	I92	L93	T94	H95	F96	M104	N105	I105A	P106	F107	L108	R109	S110	L111	I112	M113	K114				

4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	179.80Å 133.60Å 118.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.90 19.94 – 2.90	Depositor EDS
% Data completeness (in resolution range)	68.0 (8.00-2.90) 75.7 (19.94-2.90)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.88Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.219 , 0.309 0.228 , 0.299	Depositor DCC
R_{free} test set	4857 reflections (11.21%)	DCC
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.735	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 58.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	18332	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 30.54 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.2860e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, NAG, IMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/4600	0.84	6/6237 (0.1%)
1	B	0.70	0/4600	0.85	5/6237 (0.1%)
1	C	0.68	0/4600	0.84	5/6237 (0.1%)
1	D	0.68	0/4600	0.84	5/6237 (0.1%)
All	All	0.68	0/18400	0.84	21/24948 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	C	0	1
1	D	0	1
All	All	0	4

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	418	GLY	N-CA-C	6.00	128.09	113.10
1	D	418	GLY	N-CA-C	5.97	128.02	113.10
1	B	258	GLY	N-CA-C	-5.97	98.19	113.10
1	C	418	GLY	N-CA-C	5.92	127.90	113.10
1	D	258	GLY	N-CA-C	-5.85	98.48	113.10

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	147	TYR	Sidechain
1	A	466	TYR	Sidechain
1	C	147	TYR	Sidechain
1	D	147	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4473	0	4375	284	0
1	B	4473	0	4375	277	0
1	C	4473	0	4375	278	0
1	D	4473	0	4375	283	0
2	A	42	0	39	1	0
2	B	42	0	39	1	0
2	C	42	0	39	1	0
2	D	42	0	39	0	0
3	A	43	0	30	4	0
3	B	43	0	30	5	0
3	C	43	0	30	5	0
3	D	43	0	30	5	0
4	A	25	0	15	8	0
4	B	25	0	15	9	0
4	C	25	0	15	8	0
4	D	25	0	15	9	0
All	All	18332	0	17836	1109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

The worst 5 of 1109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:477:SER:HB2	1:B:479:GLU:HG2	1.37	1.07
1:A:579:SER:HB2	1:D:267:LYS:HZ1	1.23	1.03
1:C:477:SER:HB2	1:C:479:GLU:HG2	1.41	1.01
1:B:183:LEU:HD22	1:B:442:ILE:HD13	1.44	0.98

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:574:GLY:O	1:D:576:PRO:HD3	1.63	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	550/587 (94%)	456 (83%)	74 (14%)	20 (4%)	4	17
1	B	550/587 (94%)	455 (83%)	74 (14%)	21 (4%)	4	15
1	C	550/587 (94%)	454 (82%)	78 (14%)	18 (3%)	4	18
1	D	550/587 (94%)	451 (82%)	76 (14%)	23 (4%)	3	12
All	All	2200/2348 (94%)	1816 (82%)	302 (14%)	82 (4%)	4	16

5 of 82 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	SER
1	A	282	ASN
1	A	398	GLU
1	A	514	PRO
1	B	138	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	493/525 (94%)	423 (86%)	70 (14%)	4	11
1	B	493/525 (94%)	421 (85%)	72 (15%)	3	11
1	C	493/525 (94%)	424 (86%)	69 (14%)	4	12
1	D	493/525 (94%)	426 (86%)	67 (14%)	4	12
All	All	1972/2100 (94%)	1694 (86%)	278 (14%)	4	12

5 of 278 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	412	SER
1	C	120	ARG
1	D	382	ASN
1	B	441	PRO
1	B	583	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 62 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	133	HIS
1	C	282	ASN
1	D	374	GLN
1	C	232	HIS
1	C	351	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	661	1	14,14,15	0.58	0	15,19,21	0.88	1 (6%)
2	NAG	A	671	1	14,14,15	0.57	0	15,19,21	1.07	0
2	NAG	A	681	1	14,14,15	0.43	0	15,19,21	0.92	1 (6%)
3	HEM	A	682	1	28,50,50	2.02	7 (25%)	17,82,82	1.05	0
4	IMN	A	701	-	21,27,27	2.40	9 (42%)	26,39,39	1.33	2 (7%)
2	NAG	B	661	1	14,14,15	0.79	0	15,19,21	0.87	1 (6%)
2	NAG	B	671	1	14,14,15	0.69	0	15,19,21	1.09	2 (13%)
2	NAG	B	681	1	14,14,15	0.46	0	15,19,21	0.89	0
3	HEM	B	682	1	28,50,50	1.92	6 (21%)	17,82,82	1.21	2 (11%)
4	IMN	B	701	-	21,27,27	2.36	11 (52%)	26,39,39	1.20	2 (7%)
2	NAG	C	661	1	14,14,15	0.79	0	15,19,21	0.86	1 (6%)
2	NAG	C	671	1	14,14,15	0.83	0	15,19,21	1.25	2 (13%)
2	NAG	C	681	1	14,14,15	0.48	0	15,19,21	0.85	0
3	HEM	C	682	1	28,50,50	1.91	6 (21%)	17,82,82	1.11	2 (11%)
4	IMN	C	701	-	21,27,27	2.37	9 (42%)	26,39,39	1.35	3 (11%)
2	NAG	D	661	1	14,14,15	0.73	0	15,19,21	0.94	1 (6%)
2	NAG	D	671	1	14,14,15	0.63	0	15,19,21	1.15	3 (20%)
2	NAG	D	681	1	14,14,15	0.78	0	15,19,21	1.02	0
3	HEM	D	682	1	28,50,50	1.92	6 (21%)	17,82,82	1.13	2 (11%)
4	IMN	D	701	-	21,27,27	2.40	8 (38%)	26,39,39	1.30	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	661	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	671	1	-	0/6/23/26	0/1/1/1
2	NAG	A	681	1	-	0/6/23/26	0/1/1/1
3	HEM	A	682	1	-	0/6/54/54	0/0/8/8
4	IMN	A	701	-	-	0/8/14/14	0/3/3/3
2	NAG	B	661	1	-	0/6/23/26	0/1/1/1
2	NAG	B	671	1	-	0/6/23/26	0/1/1/1
2	NAG	B	681	1	-	0/6/23/26	0/1/1/1
3	HEM	B	682	1	-	0/6/54/54	0/0/8/8
4	IMN	B	701	-	-	0/8/14/14	0/3/3/3
2	NAG	C	661	1	-	0/6/23/26	0/1/1/1
2	NAG	C	671	1	-	0/6/23/26	0/1/1/1
2	NAG	C	681	1	-	0/6/23/26	0/1/1/1
3	HEM	C	682	1	-	0/6/54/54	0/0/8/8
4	IMN	C	701	-	-	0/8/14/14	0/3/3/3
2	NAG	D	661	1	-	0/6/23/26	0/1/1/1
2	NAG	D	671	1	-	0/6/23/26	0/1/1/1
2	NAG	D	681	1	-	0/6/23/26	0/1/1/1
3	HEM	D	682	1	-	0/6/54/54	0/0/8/8
4	IMN	D	701	-	-	0/8/14/14	0/3/3/3

The worst 5 of 62 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	682	HEM	C3C-CAC	-5.17	1.37	1.47
3	A	682	HEM	C3C-CAC	-4.74	1.38	1.47
3	A	682	HEM	C3C-C2C	-4.71	1.34	1.40
3	B	682	HEM	C3C-C2C	-4.67	1.34	1.40
3	A	682	HEM	C3B-CAB	-4.60	1.38	1.47

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	671	NAG	C2-N2-C7	-2.68	119.03	122.94
3	B	682	HEM	CMA-C3A-C4A	-2.52	124.58	128.46
2	A	661	NAG	O5-C1-C2	-2.45	108.06	111.47
3	D	682	HEM	CMA-C3A-C4A	-2.44	124.72	128.46
2	D	671	NAG	C2-N2-C7	-2.41	119.42	122.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 56 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	661	NAG	1	0
3	A	682	HEM	4	0
4	A	701	IMN	8	0
2	B	661	NAG	1	0
3	B	682	HEM	5	0
4	B	701	IMN	9	0
2	C	661	NAG	1	0
3	C	682	HEM	5	0
4	C	701	IMN	8	0
3	D	682	HEM	5	0
4	D	701	IMN	9	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	552/587 (94%)	-0.27	4 (0%) 87 86	2, 10, 22, 35	0
1	B	552/587 (94%)	-0.27	0 100 100	2, 10, 22, 34	0
1	C	552/587 (94%)	-0.30	2 (0%) 92 92	2, 10, 22, 34	0
1	D	552/587 (94%)	-0.23	3 (0%) 90 90	2, 11, 24, 31	0
All	All	2208/2348 (94%)	-0.27	9 (0%) 92 92	2, 10, 22, 35	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	74	PHE	3.3
1	A	280	PRO	2.7
1	A	74	PHE	2.5
1	A	278	HIS	2.4
1	D	582	VAL	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAG	C	661	14/15	0.87	0.27	6.17	20,23,29,30	0
2	NAG	C	681	14/15	0.85	0.38	4.14	19,26,28,32	0
2	NAG	A	681	14/15	0.86	0.25	3.85	18,24,27,32	0
4	IMN	B	701	25/25	0.90	0.21	2.17	5,18,24,25	0
4	IMN	D	701	25/25	0.90	0.20	1.92	8,22,23,26	0
2	NAG	B	681	14/15	0.87	0.24	1.73	14,27,29,31	0
2	NAG	D	661	14/15	0.85	0.26	1.04	16,22,29,29	0
2	NAG	D	681	14/15	0.87	0.23	0.72	19,27,29,33	0
4	IMN	C	701	25/25	0.93	0.17	0.49	3,20,23,24	0
2	NAG	B	671	14/15	0.92	0.19	0.40	2,8,19,24	0
2	NAG	B	661	14/15	0.82	0.28	0.34	15,22,29,31	0
3	HEM	B	682	43/43	0.94	0.17	0.28	2,10,20,33	0
2	NAG	D	671	14/15	0.88	0.20	0.26	2,10,17,21	0
2	NAG	C	671	14/15	0.90	0.17	0.16	2,10,20,25	0
2	NAG	A	671	14/15	0.91	0.18	0.10	2,8,20,24	0
4	IMN	A	701	25/25	0.94	0.16	0.09	2,18,21,22	0
3	HEM	A	682	43/43	0.94	0.16	-0.04	2,8,23,32	0
3	HEM	C	682	43/43	0.94	0.16	-0.09	2,7,19,27	0
3	HEM	D	682	43/43	0.95	0.15	-0.34	4,10,20,33	0
2	NAG	A	661	14/15	0.82	0.33	-	19,23,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.