



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2017 – 12:10 PM EDT

PDB ID : 5CQ1
Title : Disproportionating enzyme 1 from Arabidopsis - cycloamylose soak
Authors : O'Neill, E.C.; Stevenson, C.E.M.; Tantanarat, K.; Latousakis, D.; Donaldson, M.I.; Rejzek, M.; Limpaseni, T.; Smith, A.M.; Field, R.A.; Lawson, D.M.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20029824
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	564	<div> <div>2%</div> <div> <div></div> <div>88%</div> <div>9%</div> </div> </div>
1	B	564	<div> <div>4%</div> <div> <div></div> <div>86%</div> <div>5%</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GLC	A	602	-	-	-	X
4	EDO	A	604	-	-	-	X
4	EDO	A	605	-	-	-	X
4	EDO	A	606	-	-	-	X
4	EDO	B	606	-	-	-	X
4	EDO	B	608	-	-	-	X
4	EDO	B	609	-	-	-	X
4	EDO	B	611	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-alpha-glucanotransferase DPE1, chloroplastic/amyloplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	514	Total	C	N	O	S	0	1	0
			4006	2570	678	747	11			
1	B	511	Total	C	N	O	S	0	1	0
			3982	2557	678	735	12			

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	MET	-	initiating methionine	UNP Q9LV91
A	14	HIS	-	expression tag	UNP Q9LV91
A	15	HIS	-	expression tag	UNP Q9LV91
A	16	HIS	-	expression tag	UNP Q9LV91
A	17	HIS	-	expression tag	UNP Q9LV91
A	18	HIS	-	expression tag	UNP Q9LV91
A	19	HIS	-	expression tag	UNP Q9LV91
A	20	GLY	-	expression tag	UNP Q9LV91
A	21	LYS	-	expression tag	UNP Q9LV91
A	22	PRO	-	expression tag	UNP Q9LV91
A	23	ILE	-	expression tag	UNP Q9LV91
A	24	PRO	-	expression tag	UNP Q9LV91
A	25	ASN	-	expression tag	UNP Q9LV91
A	26	PRO	-	expression tag	UNP Q9LV91
A	27	LEU	-	expression tag	UNP Q9LV91
A	28	LEU	-	expression tag	UNP Q9LV91
A	29	GLY	-	expression tag	UNP Q9LV91
A	30	LEU	-	expression tag	UNP Q9LV91
A	31	ASP	-	expression tag	UNP Q9LV91
A	32	SER	-	expression tag	UNP Q9LV91
A	33	THR	-	expression tag	UNP Q9LV91
A	34	GLU	-	expression tag	UNP Q9LV91
A	35	ASN	-	expression tag	UNP Q9LV91
A	36	LEU	-	expression tag	UNP Q9LV91
A	37	TYR	-	expression tag	UNP Q9LV91

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	38	PHE	-	expression tag	UNP Q9LV91
A	39	GLN	-	expression tag	UNP Q9LV91
A	40	GLY	-	expression tag	UNP Q9LV91
A	41	ILE	-	expression tag	UNP Q9LV91
A	42	ASP	-	expression tag	UNP Q9LV91
A	43	PRO	-	expression tag	UNP Q9LV91
A	44	PHE	-	expression tag	UNP Q9LV91
A	45	THR	-	expression tag	UNP Q9LV91
B	13	MET	-	initiating methionine	UNP Q9LV91
B	14	HIS	-	expression tag	UNP Q9LV91
B	15	HIS	-	expression tag	UNP Q9LV91
B	16	HIS	-	expression tag	UNP Q9LV91
B	17	HIS	-	expression tag	UNP Q9LV91
B	18	HIS	-	expression tag	UNP Q9LV91
B	19	HIS	-	expression tag	UNP Q9LV91
B	20	GLY	-	expression tag	UNP Q9LV91
B	21	LYS	-	expression tag	UNP Q9LV91
B	22	PRO	-	expression tag	UNP Q9LV91
B	23	ILE	-	expression tag	UNP Q9LV91
B	24	PRO	-	expression tag	UNP Q9LV91
B	25	ASN	-	expression tag	UNP Q9LV91
B	26	PRO	-	expression tag	UNP Q9LV91
B	27	LEU	-	expression tag	UNP Q9LV91
B	28	LEU	-	expression tag	UNP Q9LV91
B	29	GLY	-	expression tag	UNP Q9LV91
B	30	LEU	-	expression tag	UNP Q9LV91
B	31	ASP	-	expression tag	UNP Q9LV91
B	32	SER	-	expression tag	UNP Q9LV91
B	33	THR	-	expression tag	UNP Q9LV91
B	34	GLU	-	expression tag	UNP Q9LV91
B	35	ASN	-	expression tag	UNP Q9LV91
B	36	LEU	-	expression tag	UNP Q9LV91
B	37	TYR	-	expression tag	UNP Q9LV91
B	38	PHE	-	expression tag	UNP Q9LV91
B	39	GLN	-	expression tag	UNP Q9LV91
B	40	GLY	-	expression tag	UNP Q9LV91
B	41	ILE	-	expression tag	UNP Q9LV91
B	42	ASP	-	expression tag	UNP Q9LV91
B	43	PRO	-	expression tag	UNP Q9LV91
B	44	PHE	-	expression tag	UNP Q9LV91
B	45	THR	-	expression tag	UNP Q9LV91

- Molecule 2 is BETA-D-GLUCOSE (three-letter code: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 3 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: $C_6H_{12}O_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		
3	B	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			11	6	5		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0
4	B	1	Total 4	C 2	O 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	158	Total 158	O 158	0	0
5	B	151	Total 151	O 151	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 4-alpha-glucanotransferase DPE1, chloroplastic/amyloplastic



- Molecule 1: 4-alpha-glucanotransferase DPE1, chloroplastic/amyloplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	69.89Å 73.35Å 79.39Å 65.28° 69.49° 66.86°	Depositor
Resolution (Å)	32.03 – 2.30 32.03 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.4 (32.03-2.30) 82.0 (32.03-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.176 , 0.205 0.183 , 0.208	Depositor DCC
R_{free} test set	2688 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.651	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8417	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BGC, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/4122	0.69	0/5615
1	B	0.59	0/4099	0.70	1/5584 (0.0%)
All	All	0.59	0/8221	0.70	1/11199 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	280	ARG	NE-CZ-NH2	-6.21	117.20	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4006	0	3812	8	0
1	B	3982	0	3784	12	0
2	A	11	0	9	0	0
2	B	11	0	9	0	0
3	A	11	0	10	0	0
3	B	23	0	22	1	0
4	A	32	0	48	3	0
4	B	32	0	48	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	158	0	0	3	0
5	B	151	0	0	3	0
All	All	8417	0	7742	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:610:EDO:H11	5:A:733:HOH:O	1.64	0.96
4:B:611:EDO:H22	5:B:754:HOH:O	1.69	0.90
4:A:610:EDO:C1	5:A:733:HOH:O	2.34	0.67
1:B:371:ARG:NH1	5:B:704:HOH:O	2.30	0.62
4:B:611:EDO:C2	5:B:754:HOH:O	2.36	0.60
1:B:474:ASN:O	4:B:609:EDO:H11	2.04	0.58
1:B:200:GLU:HG2	1:B:201:LEU:N	2.21	0.56
1:A:371:ARG:NH1	5:A:705:HOH:O	2.44	0.51
1:B:294:MET:O	1:B:294:MET:HG3	2.11	0.51
1:A:332:SER:O	1:A:333:GLU:HB2	2.12	0.50
1:A:486:GLN:O	1:A:487:GLU:HB2	2.13	0.49
1:A:458:PRO:HD2	1:A:495[A]:TYR:CE1	2.51	0.46
1:B:420:GLU:OE1	3:B:601:GLC:O3	2.14	0.45
1:B:324:SER:HA	1:B:342:LEU:HD21	2.00	0.42
1:A:294:MET:HG3	1:A:294:MET:O	2.18	0.42
1:B:324:SER:CA	1:B:342:LEU:HD21	2.49	0.42
1:A:324:SER:HA	1:A:342:LEU:HD21	2.01	0.42
1:A:324:SER:CA	1:A:342:LEU:HD21	2.51	0.41
1:B:555:SER:O	1:B:558:ASN:HB2	2.21	0.41
1:B:211:ASP:OD1	1:B:211:ASP:C	2.60	0.41
1:B:246:ARG:HG2	1:B:306:TRP:CZ2	2.56	0.41
1:A:474:ASN:O	4:A:609:EDO:H22	2.21	0.40
1:B:124:LEU:HB3	1:B:125:PRO:HD2	2.03	0.40
1:B:184:SER:N	1:B:185:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	511/564 (91%)	501 (98%)	9 (2%)	1 (0%)	51	63
1	B	508/564 (90%)	500 (98%)	8 (2%)	0	100	100
All	All	1019/1128 (90%)	1001 (98%)	17 (2%)	1 (0%)	55	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/480 (86%)	401 (98%)	10 (2%)	54	72
1	B	405/480 (84%)	395 (98%)	10 (2%)	53	70
All	All	816/960 (85%)	796 (98%)	20 (2%)	53	70

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	SER
1	A	172	ASP
1	A	241	GLU
1	A	260	GLU
1	A	336	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	376	ARG
1	A	482	ASP
1	A	487	GLU
1	A	503	ASP
1	A	560	GLU
1	B	67	SER
1	B	200	GLU
1	B	241	GLU
1	B	260	GLU
1	B	336	GLN
1	B	376	ARG
1	B	482	ASP
1	B	485	ASP
1	B	503	ASP
1	B	553	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

21 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	A	601	1,3	11,11,12	0.29	0	13,15,17	1.09	1 (7%)
3	GLC	A	602	2	11,11,12	0.44	0	13,15,17	1.26	2 (15%)
4	EDO	A	603	-	3,3,3	0.57	0	2,2,2	0.50	0
4	EDO	A	604	-	3,3,3	0.86	0	2,2,2	0.49	0
4	EDO	A	605	-	3,3,3	0.44	0	2,2,2	0.19	0
4	EDO	A	606	-	3,3,3	0.50	0	2,2,2	0.37	0
4	EDO	A	607	-	3,3,3	0.66	0	2,2,2	0.25	0
4	EDO	A	608	-	3,3,3	0.72	0	2,2,2	0.15	0
4	EDO	A	609	-	3,3,3	0.74	0	2,2,2	0.35	0
4	EDO	A	610	-	3,3,3	0.73	0	2,2,2	0.20	0
3	GLC	B	601	-	12,12,12	0.54	0	17,17,17	0.90	0
2	BGC	B	602	1,3	11,11,12	0.41	0	13,15,17	1.97	2 (15%)
3	GLC	B	603	2	11,11,12	0.35	0	13,15,17	0.88	1 (7%)
4	EDO	B	604	-	3,3,3	0.50	0	2,2,2	0.43	0
4	EDO	B	605	-	3,3,3	0.65	0	2,2,2	0.36	0
4	EDO	B	606	-	3,3,3	1.02	0	2,2,2	0.06	0
4	EDO	B	607	-	3,3,3	0.76	0	2,2,2	0.30	0
4	EDO	B	608	-	3,3,3	0.72	0	2,2,2	0.23	0
4	EDO	B	609	-	3,3,3	0.94	0	2,2,2	0.28	0
4	EDO	B	610	-	3,3,3	0.55	0	2,2,2	0.47	0
4	EDO	B	611	-	3,3,3	0.33	0	2,2,2	0.42	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	601	1,3	-	0/2/19/22	0/1/1/1
3	GLC	A	602	2	-	0/2/19/22	0/1/1/1
4	EDO	A	603	-	-	0/1/1/1	0/0/0/0
4	EDO	A	604	-	-	0/1/1/1	0/0/0/0
4	EDO	A	605	-	-	0/1/1/1	0/0/0/0
4	EDO	A	606	-	-	0/1/1/1	0/0/0/0
4	EDO	A	607	-	-	0/1/1/1	0/0/0/0
4	EDO	A	608	-	-	0/1/1/1	0/0/0/0
4	EDO	A	609	-	-	0/1/1/1	0/0/0/0
4	EDO	A	610	-	-	0/1/1/1	0/0/0/0
3	GLC	B	601	-	-	0/2/22/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	602	1,3	-	0/2/19/22	0/1/1/1
3	GLC	B	603	2	-	0/2/19/22	0/1/1/1
4	EDO	B	604	-	-	0/1/1/1	0/0/0/0
4	EDO	B	605	-	-	0/1/1/1	0/0/0/0
4	EDO	B	606	-	-	0/1/1/1	0/0/0/0
4	EDO	B	607	-	-	0/1/1/1	0/0/0/0
4	EDO	B	608	-	-	0/1/1/1	0/0/0/0
4	EDO	B	609	-	-	0/1/1/1	0/0/0/0
4	EDO	B	610	-	-	0/1/1/1	0/0/0/0
4	EDO	B	611	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	602	BGC	O4-C4-C5	-2.23	103.67	109.28
3	A	602	GLC	C1-C2-C3	2.50	112.82	109.65
3	B	603	GLC	C1-O5-C5	2.51	115.62	112.17
2	A	601	BGC	C1-O5-C5	2.94	116.22	112.17
3	A	602	GLC	C1-O5-C5	3.03	116.34	112.17
2	B	602	BGC	C1-O5-C5	6.27	120.80	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	609	EDO	1	0
4	A	610	EDO	2	0
3	B	601	GLC	1	0
4	B	609	EDO	1	0
4	B	611	EDO	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	514/564 (91%)	-0.27	14 (2%) 55 62	19, 38, 72, 111	0
1	B	511/564 (90%)	-0.01	24 (4%) 32 39	20, 40, 72, 105	0
All	All	1025/1128 (90%)	-0.14	38 (3%) 42 49	19, 39, 72, 111	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	393	VAL	6.3
1	A	75	VAL	4.4
1	A	67	SER	4.4
1	A	68	GLU	4.3
1	A	65	PHE	4.1
1	B	75	VAL	4.0
1	A	71	GLN	3.8
1	B	386	SER	3.7
1	B	337	LEU	3.5
1	A	332	SER	3.3
1	B	67	SER	3.3
1	B	66	PRO	3.2
1	A	334	THR	3.2
1	A	72	TRP	3.1
1	B	321	LEU	2.9
1	B	73	LEU	2.9
1	B	385	PRO	2.9
1	B	234	TYR	2.8
1	B	132	GLY	2.6
1	A	64	ASP	2.5
1	A	390	VAL	2.5
1	A	66	PRO	2.5
1	A	234	TYR	2.5
1	B	65	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	424	VAL	2.4
1	B	390	VAL	2.4
1	B	76	PRO	2.3
1	B	392	MET	2.3
1	B	294	MET	2.2
1	B	125	PRO	2.2
1	B	126	LEU	2.2
1	B	322	LEU	2.2
1	A	69	TYR	2.2
1	B	395	ARG	2.2
1	B	72	TRP	2.2
1	A	199	GLY	2.1
1	B	388	ALA	2.1
1	B	163	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	A	606	4/4	0.95	0.27	9.39	62,62,64,65	0
4	EDO	B	608	4/4	0.73	0.24	6.11	66,66,68,68	0
3	GLC	A	602	11/12	0.80	0.23	4.01	61,69,74,76	0
4	EDO	B	609	4/4	0.93	0.16	2.74	34,40,40,43	0
4	EDO	B	606	4/4	0.86	0.27	2.32	47,49,50,50	0
4	EDO	A	604	4/4	0.88	0.16	2.26	48,49,53,53	0
4	EDO	A	605	4/4	0.93	0.19	2.19	52,53,55,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	EDO	B	611	4/4	0.90	0.19	2.08	48,48,51,53	0
4	EDO	B	605	4/4	0.90	0.23	1.48	54,62,63,68	0
4	EDO	A	610	4/4	0.94	0.15	1.13	43,44,45,45	0
3	GLC	B	603	11/12	0.83	0.20	1.04	59,66,72,73	0
4	EDO	B	610	4/4	0.95	0.16	1.04	52,57,60,65	0
4	EDO	A	607	4/4	0.97	0.13	0.86	44,44,45,48	0
2	BGC	A	601	11/12	0.90	0.15	0.72	59,70,85,92	0
4	EDO	A	609	4/4	0.97	0.10	0.16	38,43,43,45	0
4	EDO	B	607	4/4	0.94	0.13	-0.10	44,46,46,47	0
3	GLC	B	601	12/12	0.82	0.18	-0.11	64,71,80,83	0
2	BGC	B	602	11/12	0.89	0.15	-0.18	52,63,67,69	0
4	EDO	A	603	4/4	0.98	0.09	-	39,40,41,41	0
4	EDO	A	608	4/4	0.88	0.15	-	47,52,54,55	0
4	EDO	B	604	4/4	0.95	0.10	-	42,43,44,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.