



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:38 pm GMT

PDB ID : 1CT1  
Title : CHOLERA TOXIN B-PENTAMER MUTANT G33R BOUND TO RECEPTOR PENTASACCHARIDE  
Authors : Merritt, E.A.; Hol, W.G.J.  
Deposited on : 1997-06-03  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	103	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>19%</div> <div>.</div> </div> </div>
1	E	103	<div> <div>8%</div> <div> <div></div> <div>81%</div> <div>17%</div> <div>.</div> </div> </div>
1	F	103	<div> <div>%</div> <div> <div></div> <div>76%</div> <div>20%</div> <div>..</div> </div> </div>
1	G	103	<div> <div></div> <div> <div></div> <div>72%</div> <div>25%</div> <div>..</div> </div> </div>
1	H	103	<div> <div>5%</div> <div> <div></div> <div>85%</div> <div>12%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHOLERA TOXIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	103	Total	C	N	O	S	0	0	0
			821	515	145	156	5			
1	E	103	Total	C	N	O	S	0	0	0
			821	515	145	156	5			
1	F	103	Total	C	N	O	S	0	0	0
			821	515	145	156	5			
1	G	103	Total	C	N	O	S	0	0	0
			821	515	145	156	5			
1	H	103	Total	C	N	O	S	0	0	0
			821	515	145	156	5			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	18	HIS	TYR	CONFLICT	UNP P01556
D	33	ARG	GLY	ENGINEERED	UNP P01556
D	47	THR	ILE	CONFLICT	UNP P01556
E	18	HIS	TYR	CONFLICT	UNP P01556
E	33	ARG	GLY	ENGINEERED	UNP P01556
E	47	THR	ILE	CONFLICT	UNP P01556
F	18	HIS	TYR	CONFLICT	UNP P01556
F	33	ARG	GLY	ENGINEERED	UNP P01556
F	47	THR	ILE	CONFLICT	UNP P01556
G	18	HIS	TYR	CONFLICT	UNP P01556
G	33	ARG	GLY	ENGINEERED	UNP P01556
G	47	THR	ILE	CONFLICT	UNP P01556
H	18	HIS	TYR	CONFLICT	UNP P01556
H	33	ARG	GLY	ENGINEERED	UNP P01556
H	47	THR	ILE	CONFLICT	UNP P01556

- Molecule 2 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	F	5	Total	C	N	O	0	0
			68	37	2	29		
2	G	5	Total	C	N	O	0	0
			68	37	2	29		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	25	Total	O	0	0
			25	25		
4	E	22	Total	O	0	0
			22	22		
4	F	35	Total	O	0	0
			35	35		
4	G	37	Total	O	0	0
			37	37		
4	H	32	Total	O	0	0
			32	32		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: CHOLERA TOXIN




#### • Molecule 1: CHOLERA TOXIN




#### • Molecule 1: CHOLERA TOXIN



#### • Molecule 1: CHOLERA TOXIN



#### • Molecule 1: CHOLERA TOXIN



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.40Å 67.61Å 101.70Å 90.00° 131.66° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30 25.63 – 2.28	Depositor EDS
% Data completeness (in resolution range)	92.0 (15.00-2.30) 92.2 (25.63-2.28)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.41 (at 2.29Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.182 , 0.250 0.185 , 0.243	Depositor DCC
$R_{free}$ test set	1527 reflections (7.39%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.0	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 77.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4393	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.99% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, SIA, BGC, NGA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	D	0.49	0/835	0.72	1/1127 (0.1%)
1	E	0.49	0/835	0.70	1/1127 (0.1%)
1	F	0.58	1/835 (0.1%)	0.79	2/1127 (0.2%)
1	G	0.71	1/835 (0.1%)	0.78	2/1127 (0.2%)
1	H	0.61	2/835 (0.2%)	0.75	2/1127 (0.2%)
All	All	0.58	4/4175 (0.1%)	0.75	8/5635 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	68	MET	CG-SD	-12.28	1.49	1.81
1	H	68	MET	SD-CE	7.30	2.18	1.77
1	F	68	MET	SD-CE	7.11	2.17	1.77
1	H	68	MET	CG-SD	-6.50	1.64	1.81

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	31	LEU	CA-CB-CG	7.58	132.72	115.30
1	H	31	LEU	CA-CB-CG	6.60	130.48	115.30
1	G	31	LEU	CA-CB-CG	6.55	130.37	115.30
1	F	68	MET	CG-SD-CE	-6.44	89.90	100.20
1	H	68	MET	CG-SD-CE	-5.90	90.76	100.20
1	G	50	VAL	N-CA-C	-5.82	95.30	111.00
1	E	31	LEU	CA-CB-CG	5.49	127.92	115.30
1	D	50	VAL	N-CA-C	-5.19	96.99	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	821	0	827	11	0
1	E	821	0	827	13	0
1	F	821	0	827	25	0
1	G	821	0	827	30	0
1	H	821	0	827	15	0
2	F	68	0	58	4	0
2	G	68	0	57	1	0
3	H	1	0	0	0	0
4	D	25	0	0	1	0
4	E	22	0	0	2	0
4	F	35	0	0	4	0
4	G	37	0	0	4	0
4	H	32	0	0	3	0
All	All	4393	0	4250	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:68:MET:CE	1:F:68:MET:SD	2.17	1.31
1:H:68:MET:SD	1:H:68:MET:CE	2.18	1.31
1:G:54:GLY:H	1:G:57:HIS:HD2	1.12	0.94
1:G:23:LYS:HA	4:G:135:HOH:O	1.71	0.89
1:H:9:CYS:SG	4:H:126:HOH:O	2.35	0.84
2:G:108:SIA:H92	1:H:33:ARG:CZ	2.14	0.76
1:E:83:GLU:HG2	1:E:84:LYS:HG3	1.70	0.73
1:G:54:GLY:N	1:G:57:HIS:HD2	1.87	0.72
2:F:108:SIA:H111	1:G:34:LYS:HD2	1.73	0.70
1:H:17:ILE:HG12	4:H:126:HOH:O	1.92	0.69
1:F:33:ARG:HH11	1:F:33:ARG:HG2	1.56	0.69
1:G:54:GLY:H	1:G:57:HIS:CD2	2.02	0.68
1:H:68:MET:CE	1:H:68:MET:CG	2.75	0.64
1:D:102:ALA:HB2	1:E:26:SER:OG	1.97	0.64

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:63:LYS:HD3	1:D:64:ALA:N	2.13	0.63
1:D:83:GLU:HG2	1:D:84:LYS:HG3	1.81	0.63
1:G:44:ASN:HB2	4:G:134:HOH:O	1.99	0.61
1:F:4:ASN:ND2	1:F:7:ASP:H	1.98	0.61
1:E:97:ALA:HA	1:F:31:LEU:HD22	1.83	0.60
1:G:26:SER:OG	1:G:41:THR:HB	2.02	0.60
1:F:68:MET:CE	1:F:68:MET:CG	2.79	0.60
2:F:108:SIA:C9	1:G:33:ARG:HD3	2.33	0.58
1:F:33:ARG:HH11	1:F:33:ARG:CG	2.16	0.58
1:G:82:VAL:HG13	4:G:135:HOH:O	2.05	0.57
1:F:4:ASN:ND2	1:F:6:THR:HB	2.21	0.56
1:F:57:HIS:CE1	4:F:129:HOH:O	2.59	0.54
1:F:4:ASN:HD22	1:F:6:THR:N	2.06	0.54
1:F:4:ASN:HD22	1:F:6:THR:H	1.55	0.54
1:G:1:THR:HB	1:G:2:PRO:CD	2.37	0.54
1:D:16:GLN:OE1	1:D:89:ASN:HB3	2.08	0.54
1:E:9:CYS:SG	1:E:15:THR:HB	2.49	0.53
1:G:82:VAL:CG1	4:G:135:HOH:O	2.56	0.53
1:G:74:ILE:HD11	1:H:73:ARG:HD2	1.91	0.52
1:D:33:ARG:HG3	1:D:34:LYS:HG2	1.92	0.52
2:F:108:SIA:H91	1:G:33:ARG:HD3	1.92	0.51
1:E:36:GLU:O	1:E:52:VAL:HG22	2.10	0.51
1:F:4:ASN:HD21	1:F:7:ASP:H	1.57	0.50
1:F:63:LYS:HG3	1:F:64:ALA:N	2.26	0.50
1:F:58:ILE:HG12	1:G:33:ARG:HD2	1.96	0.48
1:G:71:THR:HA	1:G:74:ILE:HD11	1.95	0.47
1:F:97:ALA:HA	1:G:31:LEU:HD22	1.97	0.47
4:F:133:HOH:O	1:G:53:PRO:HD2	2.14	0.47
1:E:56:GLN:HG3	1:E:57:HIS:N	2.29	0.46
1:F:79:GLU:HA	4:F:113:HOH:O	2.14	0.46
1:D:56:GLN:HG3	1:D:57:HIS:N	2.31	0.46
2:F:108:SIA:H92	1:G:33:ARG:HD3	1.96	0.46
1:G:58:ILE:CD1	1:H:33:ARG:HD2	2.45	0.46
1:F:30:SER:O	1:F:36:GLU:HA	2.16	0.45
1:G:44:ASN:H	1:G:44:ASN:HD22	1.63	0.45
1:E:43:LYS:HB2	4:E:114:HOH:O	2.16	0.44
1:E:18:HIS:O	1:E:84:LYS:HA	2.17	0.44
1:D:35:ARG:NH2	1:H:11:GLU:HG2	2.33	0.43
1:G:83:GLU:HG2	1:G:84:LYS:HG3	2.00	0.43
1:F:33:ARG:O	1:F:34:LYS:HB2	2.17	0.43
1:H:78:THR:O	1:H:79:GLU:HB2	2.18	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:79:GLU:HA	4:E:110:HOH:O	2.18	0.43
1:G:3:GLN:HG2	1:H:93:PRO:HG3	2.00	0.43
1:F:33:ARG:NH1	1:F:33:ARG:CG	2.78	0.43
1:H:57:HIS:HB3	1:H:61:GLN:CB	2.49	0.43
1:D:58:ILE:O	1:D:61:GLN:HB2	2.19	0.43
1:D:13:HIS:HB3	4:D:124:HOH:O	2.18	0.42
1:F:61:GLN:O	1:F:65:ILE:HG13	2.19	0.42
1:D:30:SER:O	1:D:36:GLU:HA	2.20	0.42
1:G:33:ARG:O	1:G:34:LYS:HB2	2.20	0.41
1:F:1:THR:HG23	1:F:2:PRO:HD2	2.02	0.41
1:H:57:HIS:HB3	1:H:61:GLN:HB3	2.01	0.41
1:D:41:THR:HA	1:D:46:ALA:O	2.20	0.41
1:F:103:ASN:ND2	1:G:23:LYS:NZ	2.69	0.41
1:F:57:HIS:HE1	4:F:129:HOH:O	2.00	0.41
1:F:15:THR:HA	1:F:87:VAL:O	2.20	0.41
1:G:34:LYS:O	1:G:35:ARG:HD3	2.21	0.41
1:G:70:ASP:O	1:G:74:ILE:HG13	2.21	0.41
1:H:17:ILE:N	4:H:126:HOH:O	2.54	0.41
1:F:40:ILE:HG13	1:F:40:ILE:O	2.20	0.41
1:E:12:TYR:HB2	1:E:15:THR:HG21	2.02	0.41
1:E:15:THR:HA	1:E:87:VAL:O	2.21	0.40
1:G:44:ASN:N	1:G:44:ASN:HD22	2.19	0.40
1:E:3:GLN:HE21	1:E:3:GLN:HB3	1.71	0.40
1:E:12:TYR:OH	1:F:35:ARG:HB2	2.20	0.40
1:G:97:ALA:O	1:H:31:LEU:HD22	2.21	0.40
1:G:74:ILE:H	1:G:74:ILE:HG13	1.54	0.40
1:G:74:ILE:HD13	1:H:77:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	101/103 (98%)	97 (96%)	4 (4%)	0	100	100
1	E	101/103 (98%)	97 (96%)	4 (4%)	0	100	100
1	F	101/103 (98%)	99 (98%)	2 (2%)	0	100	100
1	G	101/103 (98%)	98 (97%)	3 (3%)	0	100	100
1	H	101/103 (98%)	95 (94%)	6 (6%)	0	100	100
All	All	505/515 (98%)	486 (96%)	19 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	90/90 (100%)	86 (96%)	4 (4%)	33	45
1	E	90/90 (100%)	85 (94%)	5 (6%)	25	33
1	F	90/90 (100%)	86 (96%)	4 (4%)	33	45
1	G	90/90 (100%)	82 (91%)	8 (9%)	11	13
1	H	90/90 (100%)	86 (96%)	4 (4%)	33	45
All	All	450/450 (100%)	425 (94%)	25 (6%)	25	33

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	33	ARG
1	D	63	LYS
1	D	82	VAL
1	D	88	TRP
1	E	26	SER
1	E	31	LEU
1	E	35	ARG
1	E	82	VAL
1	E	103	ASN
1	F	31	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	33	ARG
1	F	43	LYS
1	F	103	ASN
1	G	16	GLN
1	G	31	LEU
1	G	44	ASN
1	G	55	SER
1	G	74	ILE
1	G	77	LEU
1	G	81	LYS
1	G	103	ASN
1	H	11	GLU
1	H	31	LEU
1	H	58	ILE
1	H	91	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	14	ASN
1	D	94	HIS
1	D	103	ASN
1	E	21	ASN
1	E	103	ASN
1	F	4	ASN
1	F	103	ASN
1	G	3	GLN
1	G	21	ASN
1	G	44	ASN
1	G	57	HIS
1	H	18	HIS
1	H	21	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

10 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	GAL	F	104	2	11,11,12	1.95	2 (18%)	13,15,17	2.23	5 (38%)
2	NGA	F	105	2	14,14,15	0.84	0	15,19,21	1.13	2 (13%)
2	GAL	F	106	2	11,11,12	1.41	1 (9%)	13,15,17	1.78	3 (23%)
2	BGC	F	107	2	12,12,12	1.05	0	17,17,17	1.47	4 (23%)
2	SIA	F	108	2	17,20,21	1.24	2 (11%)	19,28,31	1.42	4 (21%)
2	GAL	G	104	2	11,11,12	2.42	3 (27%)	13,15,17	0.87	0
2	NGA	G	105	2	14,14,15	1.67	4 (28%)	15,19,21	1.66	4 (26%)
2	GAL	G	106	2	11,11,12	2.47	2 (18%)	13,15,17	2.60	4 (30%)
2	BGC	G	107	2	12,12,12	1.31	2 (16%)	17,17,17	1.93	6 (35%)
2	SIA	G	108	2	17,20,21	2.11	4 (23%)	19,28,31	1.94	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GAL	F	104	2	-	0/2/19/22	0/1/1/1
2	NGA	F	105	2	-	0/6/23/26	0/1/1/1
2	GAL	F	106	2	-	0/2/19/22	0/1/1/1
2	BGC	F	107	2	-	0/2/22/22	0/1/1/1
2	SIA	F	108	2	-	0/14/34/38	0/1/1/1
2	GAL	G	104	2	-	0/2/19/22	0/1/1/1
2	NGA	G	105	2	-	0/6/23/26	0/1/1/1
2	GAL	G	106	2	-	0/2/19/22	0/1/1/1
2	BGC	G	107	2	-	0/2/22/22	0/1/1/1
2	SIA	G	108	2	-	0/14/34/38	0/1/1/1

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	106	GAL	O5-C1	-4.72	1.36	1.43
2	G	108	SIA	O8-C8	-3.01	1.36	1.43
2	G	107	BGC	C4-C3	2.09	1.57	1.52
2	F	108	SIA	C4-C5	2.19	1.55	1.53
2	G	104	GAL	C1-C2	2.25	1.57	1.52
2	G	108	SIA	O6-C6	2.30	1.47	1.43
2	G	105	NGA	O5-C5	2.38	1.48	1.43
2	G	105	NGA	O5-C1	2.45	1.47	1.43
2	G	105	NGA	C3-C2	2.50	1.57	1.52
2	F	104	GAL	C4-C5	2.77	1.58	1.53
2	G	107	BGC	C4-C5	2.88	1.59	1.53
2	F	108	SIA	C8-C7	2.96	1.59	1.53
2	G	104	GAL	C4-C5	3.13	1.59	1.53
2	G	105	NGA	C4-C3	3.35	1.60	1.52
2	F	106	GAL	C4-C5	3.65	1.60	1.53
2	G	108	SIA	C8-C7	3.94	1.61	1.53
2	F	104	GAL	C2-C3	4.80	1.59	1.52
2	G	108	SIA	C4-C5	5.79	1.58	1.53
2	G	106	GAL	C1-C2	6.03	1.66	1.52
2	G	104	GAL	C2-C3	6.53	1.61	1.52

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	104	GAL	C1-C2-C3	-5.84	102.25	109.65
2	G	108	SIA	C8-C7-C6	-3.38	106.48	113.04
2	G	107	BGC	C1-O5-C5	-3.01	107.97	113.39
2	G	107	BGC	C1-C2-C3	-2.91	105.40	110.65
2	F	105	NGA	O3-C3-C2	-2.70	103.61	109.39
2	G	106	GAL	O2-C2-C3	-2.59	105.09	110.17
2	G	107	BGC	O4-C4-C5	-2.57	102.81	109.28
2	F	107	BGC	O1-C1-C2	-2.56	102.19	109.42
2	G	107	BGC	C4-C3-C2	-2.50	106.42	110.84
2	F	104	GAL	O3-C3-C4	-2.46	105.02	110.36
2	F	106	GAL	O3-C3-C2	-2.36	105.74	110.02
2	G	105	NGA	O7-C7-C8	-2.34	117.80	122.06
2	G	105	NGA	O3-C3-C2	-2.30	104.45	109.39
2	F	107	BGC	C6-C5-C4	-2.22	107.81	113.00
2	F	108	SIA	O7-C7-C6	-2.21	104.53	109.46
2	F	108	SIA	C8-C7-C6	-2.14	108.87	113.04
2	F	107	BGC	O4-C4-C5	-2.06	104.08	109.28
2	F	105	NGA	C6-C5-C4	-2.01	108.30	113.00
2	F	108	SIA	C6-C5-N5	2.05	114.60	111.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	108	SIA	C6-C5-N5	2.05	114.61	111.00
2	G	105	NGA	O4-C4-C3	2.07	114.87	110.36
2	F	104	GAL	C2-C3-C4	2.16	114.64	110.88
2	G	106	GAL	O3-C3-C4	2.20	115.15	110.36
2	F	104	GAL	C6-C5-C4	2.26	118.29	113.00
2	G	107	BGC	C6-C5-C4	2.48	118.81	113.00
2	F	107	BGC	O5-C5-C6	2.83	113.19	106.41
2	F	104	GAL	O2-C2-C3	2.97	116.00	110.17
2	F	106	GAL	O5-C1-C2	3.24	115.87	110.79
2	G	107	BGC	C3-C4-C5	3.25	115.95	110.22
2	G	106	GAL	O5-C1-C2	3.26	115.90	110.79
2	F	108	SIA	O8-C8-C7	3.38	117.48	109.09
2	G	108	SIA	O9-C9-C8	3.65	119.16	111.11
2	F	106	GAL	C1-O5-C5	3.75	117.34	112.17
2	G	105	NGA	C1-O5-C5	4.06	117.76	112.17
2	G	108	SIA	O8-C8-C7	4.69	120.74	109.09
2	G	106	GAL	C1-O5-C5	7.47	122.46	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	108	SIA	4	0
2	G	108	SIA	1	0

## 5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	D	103/103 (100%)	0.03	6 (5%) 24 30	13, 23, 59, 86	0
1	E	103/103 (100%)	0.17	8 (7%) 14 19	11, 27, 61, 80	0
1	F	103/103 (100%)	-0.26	1 (0%) 82 86	11, 21, 38, 56	0
1	G	103/103 (100%)	-0.33	0 100 100	9, 20, 40, 60	0
1	H	103/103 (100%)	-0.05	5 (4%) 30 37	11, 20, 56, 90	0
All	All	515/515 (100%)	-0.09	20 (3%) 40 47	9, 23, 53, 90	0

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	103	ASN	7.5
1	D	56	GLN	4.5
1	H	59	ASP	4.5
1	D	59	ASP	4.4
1	E	56	GLN	4.2
1	E	55	SER	4.1
1	H	55	SER	3.9
1	H	56	GLN	3.6
1	E	13	HIS	3.1
1	H	57	HIS	3.1
1	F	103	ASN	3.1
1	D	58	ILE	3.0
1	E	54	GLY	2.9
1	D	55	SER	2.8
1	H	54	GLY	2.8
1	E	59	ASP	2.6
1	E	58	ILE	2.4
1	D	1	THR	2.4
1	E	57	HIS	2.1
1	D	103	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	G	107	12/12	0.87	0.17	1.85	41,43,47,52	0
2	SIA	G	108	20/21	0.81	0.22	1.72	43,49,54,55	0
2	BGC	F	107	12/12	0.85	0.17	1.48	40,45,47,49	0
2	NGA	G	105	14/15	0.81	0.20	1.07	37,41,43,43	0
2	SIA	F	108	20/21	0.87	0.17	0.39	40,46,52,54	0
2	GAL	G	104	11/12	0.90	0.13	0.29	27,29,32,34	0
2	NGA	F	105	14/15	0.89	0.14	0.05	28,32,40,44	0
2	GAL	F	104	11/12	0.92	0.12	-0.21	22,25,29,31	0
2	GAL	F	106	11/12	0.92	0.18	-	36,37,38,39	0
2	GAL	G	106	11/12	0.89	0.18	-	43,45,46,47	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	H	104	1/1	0.98	0.12	-	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.