



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 05:20 am GMT

PDB ID : 4CTT
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with (S)-6-(2-amino-2-(3-(2-(4-methylpyridin-2-yl)ethyl)phenyl)ethyl)-4-methylpyridin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-03-15
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>18%</div> <div>• •</div> </div> </div>
1	B	422	<div> <div>2%</div> <div> <div></div> <div>83%</div> <div>14%</div> <div>•</div> </div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6969 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, BRAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	408	Total	C	N	O	S	0	1	1
			3319	2124	568	606	21			
1	B	411	Total	C	N	O	S	0	0	0
			3345	2140	574	610	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is (S)-6-(2-AMINO-2-(3-(2-(4-METHYLPYRIDIN-2-YL)ETHYL)PHENYL)ETHYL)-4-METHYLPYRIDIN-2-AMINE (three-letter code: S7K) (formula: C₂₂H₂₆N₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N		0	0
			26	22	4			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	0	0
			26	22	4		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	61	Total	O	0	0
			61	61		
7	B	63	Total	O	0	0
			63	63		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



• Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.69Å 111.33Å 164.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.19 – 2.30 49.13 – 2.28	Depositor EDS
% Data completeness (in resolution range)	99.9 (92.19-2.30) 99.4 (49.13-2.28)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.195 , 0.249 0.194 , 0.249	Depositor DCC
R_{free} test set	2134 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	47.9	Xtriage
Anisotropy	0.734	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6969	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: S7K, ZN, H4B, HEM, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	8/3415 (0.2%)	0.70	0/4634
1	B	0.72	4/3438 (0.1%)	0.72	0/4661
All	All	0.72	12/6853 (0.2%)	0.71	0/9295

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	374	SER	CB-OG	6.49	1.50	1.42
1	A	561	TRP	CD2-CE2	5.56	1.48	1.41
1	B	409	TRP	CD2-CE2	5.45	1.47	1.41
1	A	421	TRP	CD2-CE2	5.45	1.47	1.41
1	B	716	TRP	CD2-CE2	5.42	1.47	1.41
1	A	306	TRP	CD2-CE2	5.40	1.47	1.41
1	A	475	TRP	CD2-CE2	5.39	1.47	1.41
1	A	716	TRP	CD2-CE2	5.30	1.47	1.41
1	B	711	TRP	CD2-CE2	5.27	1.47	1.41
1	A	510	TRP	CD2-CE2	5.27	1.47	1.41
1	B	306	TRP	CD2-CE2	5.14	1.47	1.41
1	A	676	TRP	CD2-CE2	5.10	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3319	0	3227	53	0
1	B	3345	0	3259	36	0
2	A	43	0	30	6	0
2	B	43	0	30	5	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	26	0	26	3	0
4	B	26	0	26	3	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0
7	A	61	0	0	3	0
7	B	63	0	0	1	0
All	All	6969	0	6634	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.45	0.98
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.57	0.86
1:A:307:GLU:HG3	7:B:2048:HOH:O	1.80	0.80
1:A:487:GLN:HB3	1:A:488:PRO:HD2	1.71	0.72
1:A:550:LYS:HB2	1:A:550:LYS:NZ	2.05	0.71
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.76	0.66
2:A:750:HEM:CMC	2:A:750:HEM:HBC2	2.22	0.62
7:A:2052:HOH:O	1:B:622:SER:HB3	2.00	0.61
1:A:299:ARG:HG3	1:A:318:LEU:HD21	1.83	0.61
1:B:587:TRP:H	2:B:750:HEM:HAB	1.67	0.59
1:A:587:TRP:H	2:A:750:HEM:HAB	1.67	0.58
1:B:325:GLY:O	1:B:332:MET:HG3	2.02	0.58
2:B:750:HEM:HMC1	2:B:750:HEM:HBC2	1.86	0.58
1:B:388:ILE:O	1:B:392:SER:N	2.36	0.56
1:A:523:LEU:CD2	1:A:531:PRO:HB2	2.33	0.55
2:B:750:HEM:HBC2	2:B:750:HEM:CMC	2.36	0.55
1:A:380:ARG:HD3	1:A:400:GLU:OE2	2.06	0.55
1:A:674:ALA:HB3	1:A:695:MET:HB3	1.87	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:567:VAL:HG23	4:A:800:S7K:H25	1.87	0.54
1:A:321:THR:HG22	7:A:2001:HOH:O	2.07	0.54
1:A:553:TRP:HZ3	1:A:557:LEU:HD11	1.73	0.54
1:B:525:GLN:HG3	1:B:529:ASN:O	2.08	0.54
1:A:620:LYS:HB2	1:A:620:LYS:NZ	2.23	0.52
1:B:300:PHE:HD1	1:B:315:THR:HG22	1.74	0.52
1:B:706:TYR:OH	2:B:750:HEM:O1D	2.23	0.52
1:B:595:VAL:O	1:B:599:CYS:HB2	2.10	0.52
1:B:362:LEU:HD11	1:B:384:VAL:HG21	1.92	0.52
1:A:507:GLN:HB2	7:A:2034:HOH:O	2.10	0.51
1:A:496:PRO:O	1:A:499:VAL:HG23	2.13	0.49
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.95	0.49
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.48	0.48
1:A:608:GLU:HB2	1:A:609:GLU:OE1	2.13	0.48
1:A:548:HIS:CE1	1:A:551:PHE:HD2	2.31	0.48
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.49	0.48
2:A:750:HEM:HBD2	2:A:750:HEM:HMD2	1.96	0.48
1:B:299:ARG:HB3	1:B:299:ARG:CZ	2.44	0.47
1:B:701:THR:HA	1:B:702:PRO:C	2.35	0.47
1:A:675:ASP:O	1:A:679:ILE:HG12	2.15	0.47
1:A:566:ALA:HB2	1:A:585:SER:HB3	1.97	0.47
1:B:485:TYR:CZ	1:B:514:ARG:HA	2.50	0.47
1:A:306:TRP:CD2	1:B:336:MET:HE3	2.49	0.47
1:A:494:GLY:O	1:A:496:PRO:HD3	2.15	0.46
1:A:470:HIS:HB3	1:A:527:ASN:ND2	2.31	0.46
1:A:686:SER:HA	1:A:691:PHE:CG	2.51	0.46
1:A:299:ARG:O	1:A:317:HIS:CE1	2.68	0.46
1:B:355:PHE:N	1:B:356:PRO:HD2	2.30	0.46
1:B:567:VAL:HG23	4:B:800:S7K:H25	1.97	0.46
1:A:684:SER:HB3	1:A:687:ILE:CG1	2.46	0.46
1:A:317:HIS:O	1:A:320:SER:HB2	2.16	0.45
1:A:550:LYS:HB2	1:A:550:LYS:HZ1	1.79	0.45
1:A:592:GLU:OE2	4:A:800:S7K:N21	2.50	0.45
1:A:525:GLN:HG3	1:A:529:ASN:O	2.17	0.45
1:B:477:SER:HB2	1:B:569:ASN:ND2	2.32	0.45
1:A:696:LEU:HD22	1:B:330:ILE:HD11	1.99	0.44
1:B:593:ILE:HA	1:B:597:ASP:HB2	2.00	0.44
1:A:566:ALA:HA	1:A:584:PHE:O	2.17	0.44
1:A:587:TRP:H	2:A:750:HEM:CAB	2.31	0.44
1:A:706:TYR:OH	4:A:800:S7K:H18	2.17	0.44
1:A:380:ARG:NH1	1:A:397:LYS:HG2	2.33	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:524:LEU:O	1:B:531:PRO:HA	2.17	0.44
1:A:631:VAL:HG11	1:B:628:GLN:CG	2.46	0.44
1:B:300:PHE:HD1	1:B:315:THR:CG2	2.30	0.44
1:A:322:LEU:HB2	1:A:699:ARG:HB2	2.00	0.44
2:A:750:HEM:HBD2	2:A:750:HEM:CMD	2.48	0.44
1:A:306:TRP:CD2	1:B:336:MET:CE	3.01	0.44
1:B:618:MET:HB3	1:B:618:MET:HE2	1.85	0.43
1:A:403:TYR:CE2	1:A:407:HIS:CE1	3.06	0.43
1:B:299:ARG:HB3	1:B:299:ARG:NH1	2.33	0.43
1:A:524:LEU:O	1:A:531:PRO:HA	2.17	0.43
1:B:403:TYR:CE2	1:B:407:HIS:CE1	3.07	0.43
1:B:504:ILE:O	1:B:508:GLN:HG2	2.19	0.43
1:A:371:ARG:CG	1:A:371:ARG:HH11	2.30	0.43
2:B:750:HEM:C4B	4:B:800:S7K:H23	2.54	0.43
1:B:567:VAL:HG21	4:B:800:S7K:C4	2.49	0.42
1:A:485:TYR:HE2	1:A:512:ALA:HB1	1.84	0.42
1:A:306:TRP:CG	1:B:336:MET:HE2	2.54	0.42
1:A:548:HIS:NE2	1:A:632:GLU:OE1	2.51	0.41
1:A:684:SER:HB3	1:A:687:ILE:HG12	2.01	0.41
1:B:546:ILE:HG12	1:B:560:LYS:HA	2.02	0.41
1:A:475:TRP:CZ2	1:A:531:PRO:HG3	2.54	0.41
1:B:425:GLN:HG2	1:B:448:TYR:CZ	2.55	0.41
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.51	0.41
1:A:330:ILE:HD11	1:B:696:LEU:HD13	2.02	0.40
1:A:686:SER:OG	1:B:595:VAL:HG12	2.22	0.40
1:A:448:TYR:CD2	1:A:448:TYR:C	2.94	0.40
1:A:462:PHE:HB3	1:A:463:PRO:CD	2.52	0.40
1:B:715:VAL:O	1:B:715:VAL:HG12	2.20	0.40
1:B:485:TYR:CE2	1:B:514:ARG:HA	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/422 (96%)	380 (94%)	24 (6%)	1 (0%)	51	63
1	B	407/422 (96%)	391 (96%)	15 (4%)	1 (0%)	51	63
All	All	812/844 (96%)	771 (95%)	39 (5%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	352	ASP
1	A	333	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/377 (97%)	346 (95%)	18 (5%)	29	39
1	B	366/377 (97%)	352 (96%)	14 (4%)	38	52
All	All	730/754 (97%)	698 (96%)	32 (4%)	33	45

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	332	MET
1	A	336	MET
1	A	337	LEU
1	A	371	ARG
1	A	454	ASN
1	A	476	ASN
1	A	486	LYS
1	A	507	GLN
1	A	523	LEU
1	A	527	ASN
1	A	547	ARG
1	A	550	LYS
1	A	552	ASP
1	A	569	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	615	ASP
1	A	622	SER
1	A	645	LYS
1	A	653	SER
1	B	321	THR
1	B	353	GLN
1	B	392	SER
1	B	423	LYS
1	B	454	ASN
1	B	455	LEU
1	B	532	GLU
1	B	535	GLN
1	B	540	LEU
1	B	547	ARG
1	B	555	LYS
1	B	608	GLU
1	B	621	THR
1	B	668	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	364	GLN
1	A	425	GLN
1	A	454	ASN
1	A	508	GLN
1	A	527	ASN
1	A	529	ASN
1	A	569	ASN
1	A	605	ASN
1	A	642	GLN
1	A	697	ASN
1	B	353	GLN
1	B	454	ASN
1	B	527	ASN
1	B	535	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	750	1	28,50,50	2.86	11 (39%)	17,82,82	3.19	9 (52%)
3	H4B	A	760	-	14,18,18	0.90	1 (7%)	12,26,26	2.17	4 (33%)
4	S7K	A	800	-	27,28,28	1.02	2 (7%)	35,38,38	1.56	7 (20%)
5	ACT	A	860	-	1,3,3	0.95	0	0,3,3	0.00	-
2	HEM	B	750	1	28,50,50	2.85	10 (35%)	17,82,82	3.57	10 (58%)
3	H4B	B	760	-	14,18,18	1.05	1 (7%)	12,26,26	2.05	4 (33%)
4	S7K	B	800	-	27,28,28	0.85	0	35,38,38	1.56	6 (17%)
5	ACT	B	860	-	1,3,3	1.13	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/6/54/54	0/0/8/8

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	S7K	A	800	-	-	0/13/13/13	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1	-	0/6/54/54	0/0/8/8
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	S7K	B	800	-	-	0/13/13/13	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	800	S7K	C6-C29	-2.98	1.48	1.51
2	B	750	HEM	C4D-ND	-2.28	1.34	1.36
3	A	760	H4B	C4-C4A	-2.08	1.38	1.41
4	A	800	S7K	C26-N21	2.29	1.38	1.34
3	B	760	H4B	C4-N3	2.30	1.37	1.33
2	A	750	HEM	C1D-CHD	2.43	1.46	1.40
2	A	750	HEM	C4A-CHB	2.72	1.47	1.40
2	A	750	HEM	C4B-CHC	2.74	1.47	1.40
2	A	750	HEM	C2A-C3A	2.95	1.46	1.37
2	B	750	HEM	C4B-CHC	3.00	1.48	1.40
2	A	750	HEM	C3B-C2B	3.10	1.44	1.40
2	B	750	HEM	C2A-C3A	3.19	1.47	1.37
2	A	750	HEM	C3D-C2D	3.41	1.47	1.37
2	B	750	HEM	C3D-C2D	3.87	1.49	1.37
2	A	750	HEM	C4A-NA	3.92	1.44	1.36
2	B	750	HEM	C1A-NA	4.23	1.45	1.36
2	B	750	HEM	C3B-C2B	4.26	1.46	1.40
2	A	750	HEM	C1A-NA	4.39	1.45	1.36
2	B	750	HEM	C4A-NA	4.61	1.46	1.36
2	B	750	HEM	C4C-NC	5.64	1.43	1.36
2	B	750	HEM	C3C-C2C	5.86	1.48	1.40
2	A	750	HEM	C3C-C2C	6.01	1.48	1.40
2	A	750	HEM	C1C-NC	6.64	1.44	1.36
2	B	750	HEM	C1C-NC	7.11	1.45	1.36
2	A	750	HEM	C4C-NC	7.16	1.45	1.36

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	C1D-C2D-C3D	-9.55	100.35	107.00
2	A	750	HEM	C1D-C2D-C3D	-8.15	101.33	107.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CAD-CBD-CGD	-5.10	103.95	112.66
2	B	750	HEM	CBA-CAA-C2A	-4.47	103.94	112.48
2	A	750	HEM	CBA-CAA-C2A	-3.83	105.16	112.48
4	B	800	S7K	C13-C12-N11	-3.81	119.53	123.92
2	B	750	HEM	C4A-C3A-C2A	-3.52	104.55	107.00
4	A	800	S7K	C13-C12-N11	-3.33	120.08	123.92
3	A	760	H4B	N3-C2-N1	-3.21	120.24	125.45
3	B	760	H4B	N3-C2-N1	-3.13	120.37	125.45
4	B	800	S7K	C25-C26-N21	-2.87	119.80	122.91
4	A	800	S7K	C25-C26-N21	-2.71	119.98	122.91
2	B	750	HEM	CAD-CBD-CGD	-2.70	108.04	112.66
4	B	800	S7K	C18-C19-C2	-2.09	105.80	113.31
2	A	750	HEM	C4A-C3A-C2A	-2.00	105.60	107.00
3	B	760	H4B	C4-N3-C2	2.02	118.97	116.06
2	B	750	HEM	CMC-C2C-C3C	2.03	128.66	124.89
2	A	750	HEM	CMC-C2C-C3C	2.18	128.94	124.89
2	A	750	HEM	C4C-C3C-C2C	2.29	108.50	106.90
4	A	800	S7K	C25-C24-C23	2.37	121.01	118.09
2	B	750	HEM	CAD-C3D-C2D	2.39	135.83	129.00
4	A	800	S7K	N22-C22-N21	2.48	120.81	116.64
3	A	760	H4B	C2-N1-C8A	2.60	120.36	114.51
4	A	800	S7K	C28-C26-N21	2.64	121.28	117.12
4	B	800	S7K	C19-C18-C16	2.82	118.58	112.71
4	A	800	S7K	C22-N21-C26	3.00	120.29	118.17
2	B	750	HEM	C4C-C3C-C2C	3.00	108.99	106.90
4	A	800	S7K	C12-N11-C16	3.01	121.46	117.40
2	A	750	HEM	CMD-C2D-C3D	3.07	130.74	124.94
3	B	760	H4B	C2-N1-C8A	3.08	121.45	114.51
2	A	750	HEM	CMB-C2B-C3B	3.26	130.94	124.89
3	A	760	H4B	C4-N3-C2	3.51	121.11	116.06
4	B	800	S7K	C22-N21-C26	3.64	120.75	118.17
3	A	760	H4B	C4-C4A-C8A	3.97	118.16	114.56
2	B	750	HEM	CMD-C2D-C3D	3.99	132.47	124.94
2	B	750	HEM	C3B-C4B-NB	4.01	114.40	109.21
4	B	800	S7K	C12-N11-C16	4.02	122.82	117.40
3	B	760	H4B	C4-C4A-C8A	4.34	118.49	114.56
2	A	750	HEM	C3B-C4B-NB	5.08	115.78	109.21
2	B	750	HEM	CMB-C2B-C3B	5.49	135.08	124.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	750	HEM	6	0
4	A	800	S7K	3	0
2	B	750	HEM	5	0
4	B	800	S7K	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	408/422 (96%)	0.17	14 (3%) 46 53	40, 67, 112, 146	0
1	B	411/422 (97%)	-0.01	7 (1%) 70 76	39, 56, 85, 114	0
All	All	819/844 (97%)	0.08	21 (2%) 56 63	39, 60, 106, 146	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	717	LYS	4.6
1	B	348	VAL	3.6
1	A	493	LEU	3.1
1	A	553	TRP	2.7
1	B	300	PHE	2.7
1	B	350	THR	2.6
1	B	349	ARG	2.6
1	A	373	GLY	2.5
1	A	715	VAL	2.5
1	A	557	LEU	2.4
1	B	351	LYS	2.4
1	A	322	LEU	2.3
1	A	716	TRP	2.3
1	A	355	PHE	2.2
1	B	389	GLU	2.2
1	A	554	PHE	2.2
1	A	366	TYR	2.1
1	A	551	PHE	2.1
1	A	516	ARG	2.1
1	A	604	TYR	2.1
1	B	718	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	S7K	B	800	26/26	0.94	0.17	1.52	42,52,89,90	0
5	ACT	A	860	4/4	0.97	0.17	1.48	75,75,75,78	0
2	HEM	B	750	43/43	0.98	0.16	1.40	41,46,53,60	0
4	S7K	A	800	26/26	0.95	0.15	1.14	40,52,97,99	0
5	ACT	B	860	4/4	0.95	0.16	0.76	58,60,64,66	0
6	ZN	A	900	1/1	0.98	0.13	0.65	53,53,53,53	0
2	HEM	A	750	43/43	0.98	0.14	0.42	40,46,55,61	0
3	H4B	A	760	17/17	0.95	0.12	-0.22	48,50,53,54	0
3	H4B	B	760	17/17	0.94	0.12	-0.79	46,49,53,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.