



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2017 – 04:44 AM EDT

PDB ID : 5CVS
Title : GlgE isoform 1 from Streptomyces coelicolor E423A mutant soaked in maltoheptaose
Authors : Rashid, A.M.; Syson, K.; Koliwer-Brandl, H.; van de Weerd, R.; Stevenson, C.E.M.; Batey, S.F.D.; Miah, F.; Alber, M.; Ioerger, T.R.; Chandra, G.; Appelmek, B.J.; Nartowski, K.P.; Khimyak, Y.Z.; Lawson, D.M.; Jacobs, W.R.; Geurtsen, J.; Kalscheuer, R.; Bornemann, S.
Deposited on : unknown
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20029824
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20029824


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4130 (2.30-2.30)
Clashscore	112137	4751 (2.30-2.30)
Ramachandran outliers	110173	4705 (2.30-2.30)
Sidechain outliers	110143	4704 (2.30-2.30)
RSRZ outliers	101464	4156 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	675	
1	B	675	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLC	A	708	-	-	-	X

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	649	Total	C	N	O	S	0	6	0
			5139	3254	937	938	10			
1	B	649	Total	C	N	O	S	0	7	0
			5100	3232	914	944	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	423	ALA	GLU	conflict	UNP Q9L1K2
B	423	ALA	GLU	conflict	UNP Q9L1K2

- Molecule 2 is ALPHA-D-GLUCOSE (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			12	6	6		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	A	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			11	6	5		
2	B	1	Total	C	O	0	0
			11	6	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	281	Total	O	0	0
			281	281		
3	B	203	Total	O	0	0
			203	203		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase 1



- Molecule 1: Alpha-1,4-glucan:maltose-1-phosphate maltosyltransferase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	114.08Å 114.08Å 314.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	62.83 – 2.30 57.04 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (62.83-2.30) 99.9 (57.04-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.181 , 0.209 0.184 , 0.211	Depositor DCC
R_{free} test set	4636 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	49.8	Xtriage
Anisotropy	0.309	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10991	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/5311	0.74	3/7265 (0.0%)
1	B	0.55	0/5276	0.74	2/7228 (0.0%)
All	All	0.56	0/10587	0.74	5/14493 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	25	ASP	CB-CG-OD2	-6.71	112.26	118.30
1	A	32	ARG	NE-CZ-NH1	5.15	122.87	120.30
1	A	606	ARG	NE-CZ-NH1	5.05	122.82	120.30
1	B	71	ARG	NE-CZ-NH1	5.02	122.81	120.30
1	B	32	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5139	0	4976	4	0
1	B	5100	0	4881	6	0
2	A	134	0	114	2	0
2	B	134	0	114	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	281	0	0	1	0
3	B	203	0	0	1	0
All	All	10991	0	10085	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:301[B]:ASP:OD1	1:B:305[B]:HIS:NE2	2.30	0.65
2:A:710:GLC:O3	2:A:711:GLC:O2	2.12	0.52
1:A:395:ASN:ND2	2:A:706:GLC:O5	2.46	0.48
1:A:103:THR:HG21	3:A:897:HOH:O	2.14	0.48
1:B:584:GLU:HG3	1:B:604:ASP:HB2	1.95	0.47
1:B:558:ARG:HA	1:B:558:ARG:NE	2.30	0.47
1:B:395:ASN:ND2	2:B:706:GLC:O5	2.48	0.46
1:B:103:THR:HG21	3:B:935:HOH:O	2.15	0.46
1:B:489:GLY:HA3	1:B:493:ALA:HB2	2.02	0.42
1:A:423:ALA:HA	1:A:443:TYR:CD1	2.55	0.41
1:A:489:GLY:HA3	1:A:493:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	653/675 (97%)	644 (99%)	8 (1%)	1 (0%)	51	63
1	B	654/675 (97%)	644 (98%)	9 (1%)	1 (0%)	51	63
All	All	1307/1350 (97%)	1288 (98%)	17 (1%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	616	ASP
1	A	616	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	532/557 (96%)	526 (99%)	6 (1%)	78	89
1	B	526/557 (94%)	520 (99%)	6 (1%)	78	89
All	All	1058/1114 (95%)	1046 (99%)	12 (1%)	78	89

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	153	ARG
1	A	154	GLU
1	A	238	ARG
1	A	251	TYR
1	A	352	ASN
1	A	610	GLU
1	B	23	VAL
1	B	152	ARG
1	B	251	TYR
1	B	352	ASN
1	B	584	GLU
1	B	610	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	A	701	2	12,12,12	0.27	0	17,17,17	0.47	0
2	GLC	A	702	2	11,11,12	0.28	0	13,15,17	0.48	0
2	GLC	A	703	2	11,11,12	0.35	0	13,15,17	0.83	0
2	GLC	A	704	2	11,11,12	0.32	0	13,15,17	0.40	0
2	GLC	A	705	2	11,11,12	0.35	0	13,15,17	0.89	0
2	GLC	A	706	2	11,11,12	0.41	0	13,15,17	0.98	1 (7%)
2	GLC	A	707	2	11,11,12	0.37	0	13,15,17	0.51	0
2	GLC	A	708	2	12,12,12	0.33	0	17,17,17	0.58	0
2	GLC	A	709	2	11,11,12	0.35	0	13,15,17	0.44	0
2	GLC	A	710	2	11,11,12	0.27	0	13,15,17	0.52	0
2	GLC	A	711	2	11,11,12	0.29	0	13,15,17	0.63	0
2	GLC	A	712	2	11,11,12	0.27	0	13,15,17	0.39	0
2	GLC	B	701	2	12,12,12	0.26	0	17,17,17	0.36	0
2	GLC	B	702	2	11,11,12	0.33	0	13,15,17	0.44	0
2	GLC	B	703	2	11,11,12	0.35	0	13,15,17	0.65	0
2	GLC	B	704	2	11,11,12	0.29	0	13,15,17	0.58	0
2	GLC	B	705	2	11,11,12	0.31	0	13,15,17	0.69	0
2	GLC	B	706	2	11,11,12	0.37	0	13,15,17	0.87	0
2	GLC	B	707	2	11,11,12	0.34	0	13,15,17	0.44	0
2	GLC	B	708	2	12,12,12	0.33	0	17,17,17	0.57	0
2	GLC	B	709	2	11,11,12	0.34	0	13,15,17	0.42	0
2	GLC	B	710	2	11,11,12	0.26	0	13,15,17	0.39	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	711	2	11,11,12	0.35	0	13,15,17	0.69	0
2	GLC	B	712	2	11,11,12	0.28	0	13,15,17	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	A	701	2	-	0/2/22/22	0/1/1/1
2	GLC	A	702	2	-	0/2/19/22	0/1/1/1
2	GLC	A	703	2	-	0/2/19/22	0/1/1/1
2	GLC	A	704	2	-	0/2/19/22	0/1/1/1
2	GLC	A	705	2	-	0/2/19/22	0/1/1/1
2	GLC	A	706	2	-	0/2/19/22	0/1/1/1
2	GLC	A	707	2	-	0/2/19/22	0/1/1/1
2	GLC	A	708	2	-	0/2/19/22	0/1/1/1
2	GLC	A	709	2	-	0/2/19/22	0/1/1/1
2	GLC	A	710	2	-	0/2/19/22	0/1/1/1
2	GLC	A	711	2	-	0/2/19/22	0/1/1/1
2	GLC	A	712	2	-	0/2/19/22	0/1/1/1
2	GLC	B	701	2	-	0/2/22/22	0/1/1/1
2	GLC	B	702	2	-	0/2/19/22	0/1/1/1
2	GLC	B	703	2	-	0/2/19/22	0/1/1/1
2	GLC	B	704	2	-	0/2/19/22	0/1/1/1
2	GLC	B	705	2	-	0/2/19/22	0/1/1/1
2	GLC	B	706	2	-	0/2/19/22	0/1/1/1
2	GLC	B	707	2	-	0/2/19/22	0/1/1/1
2	GLC	B	708	2	-	0/2/22/22	0/1/1/1
2	GLC	B	709	2	-	0/2/19/22	0/1/1/1
2	GLC	B	710	2	-	0/2/19/22	0/1/1/1
2	GLC	B	711	2	-	0/2/19/22	0/1/1/1
2	GLC	B	712	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	706	GLC	C1-O5-C5	2.01	114.94	112.17

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	706	GLC	1	0
2	A	710	GLC	1	0
2	A	711	GLC	1	0
2	B	706	GLC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	649/675 (96%)	-0.08	21 (3%)	48 55	36, 52, 95, 116	0
1	B	649/675 (96%)	-0.14	12 (1%)	69 74	37, 59, 100, 133	0
All	All	1298/1350 (96%)	-0.11	33 (2%)	58 65	36, 56, 97, 133	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	PRO	8.1
1	A	176	ALA	4.7
1	A	144	ALA	3.6
1	A	181	VAL	3.4
1	A	641	TRP	3.2
1	B	628	VAL	3.2
1	B	624	TRP	3.0
1	A	15	PRO	3.0
1	A	598	LEU	2.8
1	A	130	LEU	2.7
1	A	213	ALA	2.7
1	B	354	PRO	2.6
1	B	641	TRP	2.6
1	A	156	LEU	2.5
1	B	275	ASP	2.5
1	B	305[A]	HIS	2.5
1	A	305[A]	HIS	2.4
1	B	627	SER	2.4
1	A	212	GLY	2.3
1	A	155	LEU	2.3
1	A	56	HIS	2.3
1	A	141	ARG	2.3
1	B	639	TYR	2.2
1	A	122	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	354	PRO	2.2
1	A	226[A]	HIS	2.1
1	A	153	ARG	2.1
1	B	598	LEU	2.1
1	A	149	ARG	2.0
1	B	622	LEU	2.0
1	B	226	HIS	2.0
1	A	138	LEU	2.0
1	A	145	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GLC	A	708	12/12	0.90	0.21	2.28	76,82,90,91	0
2	GLC	A	712	11/12	0.85	0.25	1.48	86,92,97,99	0
2	GLC	B	707	11/12	0.96	0.16	1.00	57,60,63,63	0
2	GLC	B	712	11/12	0.83	0.25	0.85	108,117,120,121	0
2	GLC	B	702	11/12	0.96	0.11	0.30	61,63,68,68	0
2	GLC	A	707	11/12	0.96	0.13	0.23	45,50,52,53	0
2	GLC	A	706	11/12	0.92	0.12	-0.24	46,52,55,59	0
2	GLC	B	706	11/12	0.93	0.12	-0.41	57,63,66,66	0
2	GLC	A	705	11/12	0.97	0.12	-0.60	46,50,55,61	0
2	GLC	B	704	11/12	0.95	0.09	-0.89	63,65,66,67	0
2	GLC	A	702	11/12	0.96	0.10	-0.95	50,58,61,63	0
2	GLC	B	705	11/12	0.96	0.09	-0.98	54,60,66,66	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GLC	A	704	11/12	0.98	0.08	-1.27	50,52,54,56	0
2	GLC	A	701	12/12	0.84	0.18	-	71,88,94,96	0
2	GLC	B	709	11/12	0.83	0.29	-	117,119,123,125	0
2	GLC	B	708	12/12	0.79	0.30	-	108,111,120,126	0
2	GLC	B	701	12/12	0.87	0.13	-	77,94,98,101	0
2	GLC	A	703	11/12	0.97	0.10	-	47,48,49,51	0
2	GLC	A	710	11/12	0.92	0.13	-	81,84,86,90	0
2	GLC	B	710	11/12	0.86	0.30	-	114,120,122,127	0
2	GLC	B	703	11/12	0.96	0.07	-	63,65,67,68	0
2	GLC	A	709	11/12	0.89	0.18	-	81,83,85,87	0
2	GLC	A	711	11/12	0.95	0.23	-	93,96,99,99	0
2	GLC	B	711	11/12	0.81	0.43	-	118,126,128,129	0

6.5 Other polymers [i](#)

There are no such residues in this entry.