



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 12, 2017 – 08:20 pm GMT

PDB ID : 1CWH
Title : HUMAN CYCLOPHILIN A COMPLEXED WITH 3-D-SER CYCLOSPORIN
Authors : Mikol, V.; Kallen, J.; Taylor, P.; Walkinshaw, M.D.
Deposited on : 1998-05-07
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	112137	2083 (1.86-1.86)
Ramachandran outliers	110173	2060 (1.86-1.86)
Sidechain outliers	110143	2060 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	165	 81% 18% .
2	C	11	 45% 55%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1439 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEPTIDYL-PROLYL CIS-TRANS ISOMERASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	165	Total	C	N	O	S	0	0	0
			1266	802	218	237	9			

- Molecule 2 is a protein called CYCLOSPORIN A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	0	0	0
			87	63	11	13			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	7	DSE	SAR	ENGINEERED MUTATION	NOR NOR00033

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	83	Total	O	0	0
			83	83		
3	C	3	Total	O	0	0
			3	3		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: PEPTIDYL-PROLYL CIS-TRANS ISOMERASE A



- Molecule 2: CYCLOSPORIN A



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	62.90 Å 65.30 Å 40.80 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 1.86	Depositor
% Data completeness (in resolution range)	95.7 (8.00-1.86)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.194 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1439	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ABA, MLE, DAL, MVA, BMT, DSE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.90	0/1294	1.46	19/1733 (1.1%)
2	C	0.56	0/10	1.25	0/11
All	All	0.90	0/1304	1.46	19/1744 (1.1%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	ARG	NE-CZ-NH1	9.51	125.06	120.30
1	A	144	ARG	NE-CZ-NH2	-9.23	115.69	120.30
1	A	69	ARG	NE-CZ-NH2	-9.11	115.75	120.30
1	A	19	ARG	CB-CG-CD	-8.50	89.51	111.60
1	A	121	TRP	CD1-CG-CD2	8.15	112.82	106.30
1	A	121	TRP	CE2-CD2-CG	-7.31	101.45	107.30
1	A	148	ARG	NE-CZ-NH2	-7.21	116.69	120.30
1	A	69	ARG	NE-CZ-NH1	6.96	123.78	120.30
1	A	19	ARG	NE-CZ-NH2	-6.61	116.99	120.30
1	A	144	ARG	NE-CZ-NH1	6.47	123.54	120.30
1	A	100	MET	CG-SD-CE	-6.46	89.86	100.20
1	A	19	ARG	NE-CZ-NH1	6.34	123.47	120.30
1	A	37	ARG	NE-CZ-NH1	6.12	123.36	120.30
1	A	79	TYR	CB-CG-CD2	-5.55	117.67	121.00
1	A	48	TYR	CB-CG-CD2	-5.36	117.79	121.00
1	A	138	ILE	CG1-CB-CG2	-5.30	99.73	111.40
1	A	19	ARG	CA-CB-CG	5.25	124.96	113.40
1	A	121	TRP	CG-CD1-NE1	-5.09	105.01	110.10
1	A	158	ILE	N-CA-C	-5.04	97.39	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1266	0	1237	12	0
2	C	87	0	112	3	0
3	A	83	0	0	3	0
3	C	3	0	0	0	0
All	All	1439	0	1349	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:MET:SD	3:A:2062:HOH:O	2.41	0.78
1:A:70:HIS:HD2	3:A:2027:HOH:O	1.83	0.62
1:A:25:PHE:HD2	1:A:90:LEU:HD13	1.74	0.53
1:A:25:PHE:CD2	1:A:90:LEU:HD13	2.46	0.51
1:A:56:ILE:O	1:A:150:GLY:HA2	2.12	0.49
1:A:9:ASP:OD1	1:A:19:ARG:HD2	2.15	0.46
1:A:68:THR:HB	1:A:75:GLY:H	1.79	0.46
1:A:100:MET:CE	3:A:2011:HOH:O	2.66	0.44
1:A:82:LYS:HD2	1:A:107:THR:HA	2.00	0.43
1:A:4:PRO:HG2	1:A:24:LEU:HB2	2.01	0.41
1:A:25:PHE:HD2	1:A:90:LEU:CD1	2.33	0.41
1:A:55:ARG:HA	1:A:150:GLY:O	2.21	0.41
2:C:9:VAL:HA	2:C:10:MLE:HN1	1.86	0.41
2:C:1:DAL:HA	2:C:2:MLE:HN1	1.76	0.40
2:C:4:MVA:HA	2:C:5:BMT:HN1	1.83	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/165 (99%)	158 (97%)	5 (3%)	0	100	100
2	C	1/11 (9%)	1 (100%)	0	0	100	100
All	All	164/176 (93%)	159 (97%)	5 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/133 (100%)	124 (93%)	9 (7%)	18	5
2	C	1/1 (100%)	1 (100%)	0	100	100
All	All	134/134 (100%)	125 (93%)	9 (7%)	19	5

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	13	ASP
1	A	29	VAL
1	A	61	MET
1	A	81	GLU
1	A	102	ASN
1	A	120	GLU
1	A	148	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	154	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	HIS
1	A	102	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	DAL	C	1	2	4,4,5	0.99	0	1,4,6	0.53	0
2	MLE	C	10	2	8,8,9	1.05	1 (12%)	7,9,11	1.47	2 (28%)
2	MLE	C	2	2	8,8,9	1.03	1 (12%)	7,9,11	1.67	2 (28%)
2	MLE	C	3	2	8,8,9	0.96	0	7,9,11	1.24	1 (14%)
2	MVA	C	4	2	7,7,8	1.51	1 (14%)	7,8,10	2.03	2 (28%)
2	BMT	C	5	2	12,12,13	1.24	1 (8%)	11,14,16	1.07	1 (9%)
2	ABA	C	6	2	5,5,6	0.95	0	3,5,7	1.02	0
2	DSE	C	7	2	6,6,7	0.91	0	5,6,8	1.75	2 (40%)
2	MLE	C	8	2	8,8,9	1.07	1 (12%)	7,9,11	1.07	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DAL	C	1	2	-	0/0/2/4	0/0/0/0
2	MLE	C	10	2	-	0/4/8/10	0/0/0/0
2	MLE	C	2	2	-	0/4/8/10	0/0/0/0
2	MLE	C	3	2	-	0/4/8/10	0/0/0/0
2	MVA	C	4	2	-	0/5/8/10	0/0/0/0
2	BMT	C	5	2	-	0/13/16/18	0/0/0/0
2	ABA	C	6	2	-	0/2/4/6	0/0/0/0
2	DSE	C	7	2	-	0/2/6/8	0/0/0/0
2	MLE	C	8	2	-	0/4/8/10	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	10	MLE	CA-C	2.07	1.53	1.50
2	C	8	MLE	CA-C	2.17	1.53	1.50
2	C	5	BMT	CG2-CB	2.24	1.57	1.53
2	C	2	MLE	CA-C	2.27	1.53	1.50
2	C	4	MVA	CB-CA	3.08	1.59	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	4	MVA	CB-CA-C	-3.80	108.24	113.07
2	C	2	MLE	O-C-CA	-3.37	117.30	125.15
2	C	7	DSE	O-C-CA	-3.15	117.80	125.15
2	C	8	MLE	O-C-CA	-2.25	119.90	125.15
2	C	2	MLE	CN-N-CA	2.02	120.20	113.60
2	C	4	MVA	CN-N-CA	2.04	119.77	113.58
2	C	3	MLE	CG-CB-CA	2.14	120.91	115.27
2	C	7	DSE	CN-N-CA	2.21	120.79	113.60
2	C	5	BMT	CN-N-CA	2.32	120.61	113.58
2	C	10	MLE	CN-N-CA	2.46	121.63	113.60
2	C	10	MLE	CG-CB-CA	2.60	122.12	115.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	DAL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	10	MLE	1	0
2	C	2	MLE	1	0
2	C	4	MVA	1	0
2	C	5	BMT	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.