



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2017 – 04:11 PM EDT

PDB ID : 3CXN
Title : Structure of the Urease Accessory Protein UreF from Helicobacter pylori
Authors : Lam, R.; Johns, K.; Romanov, V.; Dong, A.; Wu-Brown, J.; Guthrie, J.; Dharamsi, A.; Thambipillai, D.; Mansoury, K.; Edwards, A.M.; Pai, E.F.; Chirgadze, N.Y.
Deposited on : unknown
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1088 (1.56-1.56)
Clashscore	112137	1132 (1.56-1.56)
Ramachandran outliers	110173	1110 (1.56-1.56)
Sidechain outliers	110143	1108 (1.56-1.56)
RSRZ outliers	101464	1089 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div> <div>13%</div> <div>68%</div> <div>7%</div> <div>24%</div> </div>
1	B	274	<div> <div>7%</div> <div>71%</div> <div>5%</div> <div>24%</div> </div>
1	C	274	<div> <div>14%</div> <div>64%</div> <div>10%</div> <div>26%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	255	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5446 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Urease accessory protein ureF.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	Se	0	9	0
			1688	1084	272	322	2	8			
1	B	209	Total	C	N	O	S	Se	0	14	0
			1733	1115	278	330	2	8			
1	C	204	Total	C	N	O	S	Se	0	2	0
			1618	1031	265	313	2	7			

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MSE	-	EXPRESSION TAG	UNP Q09065
A	-18	GLY	-	EXPRESSION TAG	UNP Q09065
A	-17	SER	-	EXPRESSION TAG	UNP Q09065
A	-16	SER	-	EXPRESSION TAG	UNP Q09065
A	-15	HIS	-	EXPRESSION TAG	UNP Q09065
A	-14	HIS	-	EXPRESSION TAG	UNP Q09065
A	-13	HIS	-	EXPRESSION TAG	UNP Q09065
A	-12	HIS	-	EXPRESSION TAG	UNP Q09065
A	-11	HIS	-	EXPRESSION TAG	UNP Q09065
A	-10	HIS	-	EXPRESSION TAG	UNP Q09065
A	-9	SER	-	EXPRESSION TAG	UNP Q09065
A	-8	SER	-	EXPRESSION TAG	UNP Q09065
A	-7	GLY	-	EXPRESSION TAG	UNP Q09065
A	-6	LEU	-	EXPRESSION TAG	UNP Q09065
A	-5	VAL	-	EXPRESSION TAG	UNP Q09065
A	-4	PRO	-	EXPRESSION TAG	UNP Q09065
A	-3	ARG	-	EXPRESSION TAG	UNP Q09065
A	-2	GLY	-	EXPRESSION TAG	UNP Q09065
A	-1	SER	-	EXPRESSION TAG	UNP Q09065
A	0	HIS	-	EXPRESSION TAG	UNP Q09065
A	1	MSE	-	EXPRESSION TAG	UNP Q09065
B	-19	MSE	-	EXPRESSION TAG	UNP Q09065
B	-18	GLY	-	EXPRESSION TAG	UNP Q09065

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	SER	-	EXPRESSION TAG	UNP Q09065
B	-16	SER	-	EXPRESSION TAG	UNP Q09065
B	-15	HIS	-	EXPRESSION TAG	UNP Q09065
B	-14	HIS	-	EXPRESSION TAG	UNP Q09065
B	-13	HIS	-	EXPRESSION TAG	UNP Q09065
B	-12	HIS	-	EXPRESSION TAG	UNP Q09065
B	-11	HIS	-	EXPRESSION TAG	UNP Q09065
B	-10	HIS	-	EXPRESSION TAG	UNP Q09065
B	-9	SER	-	EXPRESSION TAG	UNP Q09065
B	-8	SER	-	EXPRESSION TAG	UNP Q09065
B	-7	GLY	-	EXPRESSION TAG	UNP Q09065
B	-6	LEU	-	EXPRESSION TAG	UNP Q09065
B	-5	VAL	-	EXPRESSION TAG	UNP Q09065
B	-4	PRO	-	EXPRESSION TAG	UNP Q09065
B	-3	ARG	-	EXPRESSION TAG	UNP Q09065
B	-2	GLY	-	EXPRESSION TAG	UNP Q09065
B	-1	SER	-	EXPRESSION TAG	UNP Q09065
B	0	HIS	-	EXPRESSION TAG	UNP Q09065
B	1	MSE	-	EXPRESSION TAG	UNP Q09065
C	-19	MSE	-	EXPRESSION TAG	UNP Q09065
C	-18	GLY	-	EXPRESSION TAG	UNP Q09065
C	-17	SER	-	EXPRESSION TAG	UNP Q09065
C	-16	SER	-	EXPRESSION TAG	UNP Q09065
C	-15	HIS	-	EXPRESSION TAG	UNP Q09065
C	-14	HIS	-	EXPRESSION TAG	UNP Q09065
C	-13	HIS	-	EXPRESSION TAG	UNP Q09065
C	-12	HIS	-	EXPRESSION TAG	UNP Q09065
C	-11	HIS	-	EXPRESSION TAG	UNP Q09065
C	-10	HIS	-	EXPRESSION TAG	UNP Q09065
C	-9	SER	-	EXPRESSION TAG	UNP Q09065
C	-8	SER	-	EXPRESSION TAG	UNP Q09065
C	-7	GLY	-	EXPRESSION TAG	UNP Q09065
C	-6	LEU	-	EXPRESSION TAG	UNP Q09065
C	-5	VAL	-	EXPRESSION TAG	UNP Q09065
C	-4	PRO	-	EXPRESSION TAG	UNP Q09065
C	-3	ARG	-	EXPRESSION TAG	UNP Q09065
C	-2	GLY	-	EXPRESSION TAG	UNP Q09065
C	-1	SER	-	EXPRESSION TAG	UNP Q09065
C	0	HIS	-	EXPRESSION TAG	UNP Q09065
C	1	MSE	-	EXPRESSION TAG	UNP Q09065

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	118	Total	O	0	0
			118	118		
3	B	185	Total	O	0	0
			185	185		
3	C	98	Total	O	0	0
			98	98		

- Molecule 1: Urease accessory protein ureF



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	135.22Å 89.42Å 66.04Å 90.00° 94.03° 90.00°	Depositor
Resolution (Å)	26.77 – 1.55 26.77 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.3 (26.77-1.55) 99.3 (26.77-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 1.55Å)	Xtriage
Refinement program	REFMAC refmac_5.2.0019	Depositor
R, R_{free}	0.187 , 0.211 0.192 , 0.213	Depositor DCC
R_{free} test set	5637 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	18.8	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5446	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/1736	0.60	0/2330
1	B	0.54	0/1796	0.64	0/2408
1	C	0.45	0/1641	0.57	0/2203
All	All	0.50	0/5173	0.60	0/6941

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1688	0	1734	15	0
1	B	1733	0	1795	13	0
1	C	1618	0	1648	23	0
2	B	6	0	8	0	0
3	A	118	0	0	1	0
3	B	185	0	0	3	0
3	C	98	0	0	2	0
All	All	5446	0	5185	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:42:VAL:HG11	1:C:191:ILE:HD11	1.47	0.95
1:C:123:ALA:O	1:C:126:LYS:HE2	1.70	0.92
1:B:85[B]:GLU:HG2	1:B:181[B]:TYR:OH	1.71	0.90
1:B:85[B]:GLU:OE2	1:B:181[B]:TYR:OH	1.92	0.86
1:C:73:LEU:HD22	1:C:189:MSE:HE3	1.70	0.73
1:B:85[B]:GLU:CG	1:B:181[B]:TYR:OH	2.37	0.71
1:A:42:VAL:HG11	1:C:191:ILE:CD1	2.18	0.70
1:C:126:LYS:HE3	3:C:306:HOH:O	1.91	0.70
1:B:76:ASN:ND2	3:B:405:HOH:O	2.27	0.67
1:C:32:GLU:OE1	1:C:176:LYS:NZ	2.29	0.65
1:C:186:THR:HG21	1:C:215:PHE:CZ	2.33	0.64
1:C:82:LEU:HB2	1:C:218:LEU:HD11	1.83	0.61
1:B:85[B]:GLU:CD	1:B:181[B]:TYR:OH	2.39	0.60
1:B:112:MSE:HG2	1:B:121:ARG:HD2	1.84	0.60
1:A:32:GLU:OE2	1:A:176:LYS:NZ	2.35	0.59
1:C:173:GLU:OE2	1:C:175:LYS:HB3	2.03	0.58
1:A:54:LEU:HD13	1:A:72:TYR:CE2	2.41	0.55
1:A:126:LYS:O	1:A:129:ASN:ND2	2.40	0.54
1:B:218[A]:LEU:HG	3:B:333:HOH:O	2.08	0.53
1:C:179:ARG:HG2	1:C:219:ILE:HD13	1.92	0.51
1:A:51:SER:HA	1:A:192:ASN:ND2	2.26	0.51
1:C:73:LEU:CD2	1:C:189:MSE:HE3	2.41	0.50
1:A:213[B]:SER:HB3	1:A:214:PRO:HD3	1.94	0.50
1:A:54:LEU:HG	1:A:58:ILE:CD1	2.41	0.49
1:A:42:VAL:CG1	1:C:191:ILE:HD11	2.30	0.49
1:C:221:LYS:HE3	1:C:225:LEU:HD21	1.92	0.49
1:B:60:GLN:C	1:B:61:LYS:HG2	2.34	0.47
1:C:186:THR:HG21	1:C:215:PHE:CE1	2.49	0.47
1:A:179[B]:ARG:HG3	1:C:33:PHE:CE1	2.50	0.47
1:A:54:LEU:HD23	1:A:192:ASN:ND2	2.30	0.47
1:C:186:THR:HG21	1:C:215:PHE:HZ	1.75	0.46
1:B:103:LYS:HE3	3:B:380:HOH:O	2.16	0.44
1:A:90[A]:LYS:HG3	1:A:174:LEU:HD21	2.00	0.44
1:C:123:ALA:O	1:C:126:LYS:HG3	2.18	0.43
1:B:82:LEU:HB2	1:B:218[B]:LEU:HD11	2.00	0.43
1:A:153:LYS:HB3	1:A:153:LYS:HE3	1.88	0.42
1:C:118:MSE:CE	1:C:119:GLU:HG3	2.49	0.42
1:C:118:MSE:HE3	1:C:119:GLU:N	2.34	0.42
1:C:28:HIS:H	1:C:28:HIS:CD2	2.38	0.42
1:B:50:HIS:HE1	1:B:85[A]:GLU:OE1	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:LEU:HG	3:A:350:HOH:O	2.20	0.41
1:C:186:THR:HG23	1:C:189:MSE:HE2	2.02	0.41
1:C:213:SER:HB3	1:C:214:PRO:HD3	2.03	0.41
1:C:67:GLU:HG3	3:C:305:HOH:O	2.19	0.41
1:A:57:TYR:CD2	1:B:118[A]:MSE:HE1	2.55	0.41
1:B:100:ASP:OD2	1:B:103:LYS:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/274 (78%)	211 (99%)	3 (1%)	0	100	100
1	B	221/274 (81%)	218 (99%)	3 (1%)	0	100	100
1	C	202/274 (74%)	196 (97%)	6 (3%)	0	100	100
All	All	637/822 (78%)	625 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/234 (82%)	190 (98%)	3 (2%)	68	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	200/234 (86%)	197 (98%)	3 (2%)	70	41
1	C	183/234 (78%)	176 (96%)	7 (4%)	38	7
All	All	576/702 (82%)	563 (98%)	13 (2%)	57	23

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	MSE
1	A	129	ASN
1	A	166	PHE
1	B	118[A]	MSE
1	B	118[B]	MSE
1	B	166	PHE
1	C	72	TYR
1	C	76	ASN
1	C	80	GLN
1	C	118	MSE
1	C	121	ARG
1	C	126	LYS
1	C	166	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	ASN
1	A	188	ASN
1	A	192	ASN
1	B	76	ASN
1	C	25	ASN
1	C	28	HIS
1	C	80	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	255	-	5,5,5	0.30	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	255	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	201/274 (73%)	1.06	35 (17%) 2 2	17, 24, 50, 57	1 (0%)
1	B	203/274 (74%)	0.54	18 (8%) 10 12	15, 20, 35, 47	1 (0%)
1	C	198/274 (72%)	1.02	37 (18%) 1 2	20, 29, 43, 46	0
All	All	602/822 (73%)	0.87	90 (14%) 3 2	15, 25, 44, 57	2 (0%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	52	PHE	13.3
1	A	120	LEU	7.8
1	A	27	ALA	7.1
1	B	59	GLN	6.7
1	B	25	ASN	6.0
1	A	122	LEU	5.8
1	A	28	HIS	5.6
1	A	48	TYR	5.5
1	B	52	PHE	5.5
1	C	25	ASN	5.5
1	A	119	GLU	5.3
1	C	233	ALA	5.3
1	A	47	SER	5.3
1	A	53	GLY	5.1
1	A	121	ARG	4.8
1	A	51	SER	4.8
1	C	120	LEU	4.8
1	C	47	SER	4.6
1	B	56	THR	4.6
1	A	29	VAL	4.5
1	B	60	GLN	4.1
1	C	122	LEU	4.0
1	A	54	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	GLU	3.9
1	A	58	ILE	3.8
1	A	123	ALA	3.8
1	C	77	LEU	3.7
1	C	224	GLU	3.7
1	C	46	GLY	3.7
1	C	232	THR	3.7
1	C	231	CYS	3.7
1	C	83	TYR	3.6
1	A	59	GLN	3.6
1	A	49	THR	3.5
1	C	119	GLU	3.5
1	B	55	GLU	3.4
1	C	121	ARG	3.4
1	A	117	PRO	3.4
1	B	58	ILE	3.4
1	C	113	LEU	3.3
1	C	53	GLY	3.3
1	B	53	GLY	3.2
1	A	125	GLN	3.2
1	A	127	LEU	3.1
1	A	139	ASN	3.1
1	A	126	LYS	3.1
1	C	225	LEU	3.0
1	A	181[A]	TYR	3.0
1	A	61	LYS	3.0
1	A	140	GLU	3.0
1	B	181[A]	TYR	2.9
1	C	229	HIS	2.9
1	C	57	TYR	2.8
1	C	55	GLU	2.7
1	C	198	PRO	2.7
1	C	76	ASN	2.7
1	C	78	SER	2.7
1	C	27	ALA	2.7
1	A	33	PHE	2.7
1	C	98	GLN	2.6
1	B	38	VAL	2.6
1	C	116	SER	2.6
1	A	50	HIS	2.5
1	C	226	ASP	2.5
1	C	45	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	60	GLN	2.5
1	C	195	LYS	2.4
1	B	27	ALA	2.4
1	B	47	SER	2.3
1	C	36	LEU	2.3
1	C	28	HIS	2.3
1	A	182	LEU	2.3
1	B	183	TYR	2.3
1	C	34	LEU	2.2
1	A	183	TYR	2.2
1	A	137	ALA	2.2
1	C	148	ASN	2.2
1	C	81	PHE	2.2
1	C	135	LEU	2.2
1	B	163	TYR	2.2
1	B	51	SER	2.2
1	C	99	GLN	2.2
1	A	133	LYS	2.1
1	B	182	LEU	2.1
1	A	178	LEU	2.1
1	A	56	THR	2.1
1	C	56	THR	2.1
1	B	57	TYR	2.0
1	B	61	LYS	2.0
1	C	38	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GOL	B	255	6/6	0.53	0.24	4.71	53,53,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.