



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 24, 2017 – 05:38 PM EDT

PDB ID : 5CYP
Title : GTPase domain of Septin 9 in complex with GTP-gamma-S
Authors : Matos, S.S.; Leonardo, D.A.; Pereira, H.M.; Horjales, E.; Araujo, A.P.U.; Garratt, R.C.
Deposited on : unknown
Resolution : 2.89 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20030345
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.89 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	
1	B	274	
1	C	274	
1	D	274	

2 Entry composition ⓘ


There are 3 unique types of molecules in this entry. The entry contains 7249 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Septin-9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	248	Total	C	N	O	S	0	0	0
			1907	1211	337	350	9			
1	B	227	Total	C	N	O	S	0	0	0
			1697	1086	289	313	9			
1	C	238	Total	C	N	O	S	0	0	0
			1810	1151	313	336	10			
1	D	227	Total	C	N	O	S	0	0	0
			1703	1087	295	312	9			

- Molecule 2 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C₁₀H₁₆N₅O₁₃P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			32	10	5	13	3	1		


- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Septin-9



• Molecule 1: Septin-9




• Molecule 1: Septin-9



• Molecule 1: Septin-9





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.59 Å 79.16 Å 108.22 Å 90.00° 100.38° 90.00°	Depositor
Resolution (Å)	45.65 – 2.89 45.65 – 2.89	Depositor EDS
% Data completeness (in resolution range)	97.3 (45.65-2.89) 97.6 (45.65-2.89)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.91 Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.252 , 0.297 0.253 , 0.297	Depositor DCC
R_{free} test set	1361 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	50.5	Xtriage
Anisotropy	0.786	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7249	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.32 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.1650e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.21	0/1939	0.39	0/2624
1	B	0.21	0/1727	0.39	0/2348
1	C	0.21	0/1841	0.37	0/2498
1	D	0.21	0/1733	0.37	0/2357
All	All	0.21	0/7240	0.38	0/9827

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1907	0	1829	14	0
1	B	1697	0	1582	13	0
1	C	1810	0	1716	20	0
1	D	1703	0	1587	16	0
2	A	32	0	12	5	0
2	B	32	0	12	4	0
2	C	32	0	12	6	0
2	D	32	0	12	4	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
All	All	7249	0	6762	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:46:SER:OG	1:A:262:ARG:NH1	1.97	0.97
1:D:185:ARG:NH1	1:D:189:ASP:OD1	2.18	0.76
2:A:301:GSP:O3G	2:A:301:GSP:O1B	2.05	0.74
2:D:301:GSP:O1B	2:D:301:GSP:O2G	2.10	0.70
1:B:185:ARG:NH1	1:B:189:ASP:OD1	2.26	0.68
1:C:23:GLU:HG2	1:C:82:LYS:HE3	1.79	0.64
2:C:301:GSP:O3G	2:C:301:GSP:O1B	2.17	0.63
1:C:119:GLU:OE1	1:C:286:ARG:NH1	2.33	0.62
1:A:43:LEU:HD21	1:A:83:LEU:HD21	1.84	0.59
1:A:93:ASP:OD1	1:A:94:HIS:ND1	2.36	0.59
1:D:93:ASP:OD1	1:D:94:HIS:ND1	2.35	0.59
2:B:301:GSP:O3G	2:B:301:GSP:O2B	2.20	0.58
1:C:93:ASP:OD1	1:C:94:HIS:ND1	2.36	0.58
1:C:178:GLU:OE2	2:D:301:GSP:O2'	2.21	0.58
1:B:93:ASP:OD1	1:B:94:HIS:ND1	2.37	0.57
2:C:301:GSP:O2B	2:C:301:GSP:O2G	2.23	0.57
2:B:301:GSP:O1B	2:B:301:GSP:O2G	2.21	0.57
1:C:43:LEU:HD21	1:C:83:LEU:HD21	1.88	0.56
1:D:167:VAL:HG12	1:D:222:PHE:HB2	1.87	0.55
1:C:167:VAL:HG12	1:C:222:PHE:HB2	1.88	0.55
1:A:183:LYS:NZ	1:A:257:GLU:OE2	2.40	0.55
2:D:301:GSP:O1A	2:D:301:GSP:O2B	2.23	0.54
1:C:226:GLY:N	2:C:301:GSP:O6	2.33	0.54
1:C:116:LEU:HD13	1:C:286:ARG:HG2	1.91	0.53
1:D:43:LEU:HD12	1:D:262:ARG:HH11	1.74	0.52
1:B:134:HIS:CE1	1:B:274:LYS:NZ	2.78	0.51
1:C:199:PRO:HD3	1:C:216:PHE:CD1	2.45	0.51
1:A:31:GLN:CD	1:A:148:ARG:HH22	2.14	0.51
1:C:60:ARG:NH1	1:C:62:PRO:HA	2.25	0.51
1:C:31:GLN:CD	1:C:148:ARG:HH22	2.14	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:20:GLN:HB3	1:C:80:ARG:HE	1.77	0.49
1:A:241:ARG:HB3	1:A:248:ILE:HB	1.94	0.48
1:B:36:LYS:NZ	2:B:301:GSP:O2G	2.38	0.48
1:C:107:PHE:O	1:C:111:GLN:HG2	2.13	0.48
1:D:199:PRO:HD3	1:D:216:PHE:CD1	2.48	0.48
2:D:301:GSP:O2B	2:D:301:GSP:O3G	2.32	0.48
1:B:199:PRO:HD3	1:B:216:PHE:CD1	2.48	0.47
1:A:204:ASP:OD1	1:A:213:ASN:ND2	2.39	0.47
1:C:60:ARG:HH11	1:C:62:PRO:HA	1.80	0.47
1:D:216:PHE:CD2	1:D:264:LEU:HD13	2.49	0.47
1:D:36:LYS:NZ	1:D:90:GLY:HA2	2.31	0.46
1:C:186:ILE:HD12	1:C:221:PRO:HG3	1.98	0.46
1:B:182:PHE:O	1:B:186:ILE:HG12	2.16	0.46
1:D:30:GLY:O	1:D:36:LYS:NZ	2.46	0.46
2:C:301:GSP:O2'	1:D:178:GLU:OE2	2.33	0.46
1:D:71:THR:O	1:D:72:HIS:ND1	2.49	0.46
1:A:30:GLY:O	1:A:36:LYS:NZ	2.47	0.45
2:A:301:GSP:O2'	1:B:178:GLU:OE2	2.33	0.45
1:A:199:PRO:HD3	1:A:216:PHE:CD1	2.52	0.45
1:A:107:PHE:O	1:A:111:GLN:HG2	2.17	0.44
1:A:170:LYS:HE2	2:A:301:GSP:C4	2.52	0.44
1:D:182:PHE:O	1:D:186:ILE:HG12	2.16	0.44
2:A:301:GSP:O1A	1:B:145:HIS:NE2	2.38	0.43
1:C:183:LYS:NZ	1:C:257:GLU:OE2	2.52	0.43
1:B:107:PHE:O	1:B:111:GLN:HG2	2.19	0.43
1:D:107:PHE:O	1:D:111:GLN:HG2	2.20	0.42
1:A:178:GLU:OE2	2:B:301:GSP:O2'	2.37	0.42
1:B:23:GLU:OE2	1:B:130:ASP:N	2.51	0.42
1:A:31:GLN:HE21	1:A:92:GLY:HA3	1.85	0.41
1:B:137:LEU:HD21	1:B:265:LEU:HD11	2.01	0.41
1:C:140:ILE:HD13	1:C:147:LEU:HD21	2.02	0.41
1:D:158:LEU:HB3	1:D:164:ILE:HD11	2.02	0.41
1:C:182:PHE:CZ	1:C:186:ILE:HD11	2.56	0.41
2:C:301:GSP:O1A	1:D:145:HIS:NE2	2.39	0.41
1:D:172:ASP:OD1	1:D:172:ASP:N	2.53	0.41
2:A:301:GSP:O2G	2:A:301:GSP:O2B	2.39	0.41
1:C:71:THR:O	1:C:72:HIS:ND1	2.54	0.41
1:B:158:LEU:HB3	1:B:164:ILE:HD11	2.02	0.40
1:D:118:GLU:O	1:D:124:ARG:NE	2.39	0.40
1:A:182:PHE:O	1:A:186:ILE:HG12	2.20	0.40
1:C:63:LYS:HG3	2:C:301:GSP:S1G	2.62	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:43:LEU:HD21	1:B:83:LEU:HD21	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/274 (87%)	227 (95%)	11 (5%)	0	100	100
1	B	219/274 (80%)	210 (96%)	9 (4%)	0	100	100
1	C	230/274 (84%)	220 (96%)	10 (4%)	0	100	100
1	D	219/274 (80%)	212 (97%)	7 (3%)	0	100	100
All	All	906/1096 (83%)	869 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/254 (79%)	196 (98%)	4 (2%)	60	87
1	B	170/254 (67%)	168 (99%)	2 (1%)	75	93
1	C	189/254 (74%)	187 (99%)	2 (1%)	78	94
1	D	171/254 (67%)	170 (99%)	1 (1%)	89	97

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	730/1016 (72%)	721 (99%)	9 (1%)	75	93

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	127	ARG
1	A	160	LYS
1	A	167	VAL
1	B	118	GLU
1	B	167	VAL
1	C	60	ARG
1	C	80	ARG
1	D	265	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	134	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GSP	A	301	3	26,34,34	1.65	4 (15%)	24,54,54	2.66	7 (29%)
2	GSP	B	301	3	26,34,34	1.67	4 (15%)	24,54,54	2.68	6 (25%)
2	GSP	C	301	3	26,34,34	1.64	4 (15%)	24,54,54	2.66	7 (29%)
2	GSP	D	301	3	26,34,34	1.69	4 (15%)	24,54,54	2.67	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GSP	A	301	3	-	0/17/38/38	0/3/3/3
2	GSP	B	301	3	-	0/17/38/38	0/3/3/3
2	GSP	C	301	3	-	0/17/38/38	0/3/3/3
2	GSP	D	301	3	-	0/17/38/38	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	GSP	C2-N2	2.18	1.38	1.34
2	D	301	GSP	C4-N3	2.22	1.39	1.35
2	C	301	GSP	C2-N2	2.23	1.38	1.34
2	B	301	GSP	C4-N3	2.24	1.39	1.35
2	B	301	GSP	C2-N2	2.27	1.38	1.34
2	D	301	GSP	C2-N2	2.28	1.38	1.34
2	C	301	GSP	C4-N3	2.29	1.39	1.35
2	A	301	GSP	C4-N3	2.32	1.39	1.35
2	C	301	GSP	C2-N1	2.85	1.40	1.35
2	D	301	GSP	C2-N1	2.95	1.40	1.35
2	A	301	GSP	C2-N1	2.95	1.40	1.35
2	B	301	GSP	C2-N1	2.99	1.40	1.35
2	C	301	GSP	C6-N1	5.72	1.43	1.33
2	A	301	GSP	C6-N1	5.79	1.43	1.33
2	B	301	GSP	C6-N1	5.88	1.43	1.33
2	D	301	GSP	C6-N1	5.93	1.43	1.33

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	301	GSP	C5-C6-N1	-7.98	112.12	123.48
2	B	301	GSP	C5-C6-N1	-7.95	112.16	123.48
2	A	301	GSP	C5-C6-N1	-7.91	112.23	123.48
2	C	301	GSP	C5-C6-N1	-7.85	112.30	123.48
2	B	301	GSP	PB-O3B-PG	-6.96	109.85	132.35
2	C	301	GSP	PB-O3B-PG	-6.96	109.86	132.35
2	D	301	GSP	PB-O3B-PG	-6.88	110.11	132.35
2	A	301	GSP	PB-O3B-PG	-6.87	110.16	132.35
2	A	301	GSP	C2-N3-C4	-4.38	110.05	115.16
2	C	301	GSP	C2-N3-C4	-4.33	110.10	115.16
2	B	301	GSP	C2-N3-C4	-4.32	110.11	115.16
2	D	301	GSP	C2-N3-C4	-4.31	110.12	115.16
2	C	301	GSP	O2B-PB-O1B	2.04	122.86	112.28
2	A	301	GSP	O2A-PA-O1A	2.05	122.89	112.28
2	C	301	GSP	O2A-PA-O1A	2.05	122.90	112.28
2	D	301	GSP	O2A-PA-O1A	2.06	122.94	112.28
2	A	301	GSP	O2B-PB-O1B	2.06	122.96	112.28
2	B	301	GSP	O2B-PB-O1B	2.07	123.00	112.28
2	D	301	GSP	C4-C5-N7	2.35	111.68	109.41
2	B	301	GSP	C4-C5-N7	2.45	111.77	109.41
2	A	301	GSP	C4-C5-N7	2.52	111.84	109.41
2	C	301	GSP	C4-C5-N7	2.53	111.86	109.41
2	C	301	GSP	C6-N1-C2	3.49	121.09	116.06
2	A	301	GSP	C6-N1-C2	3.51	121.11	116.06
2	B	301	GSP	C6-N1-C2	3.53	121.14	116.06
2	D	301	GSP	C6-N1-C2	3.54	121.15	116.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	GSP	5	0
2	B	301	GSP	4	0
2	C	301	GSP	6	0
2	D	301	GSP	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/274 (90%)	0.20	10 (4%) 39 34	32, 60, 99, 171	0
1	B	227/274 (82%)	0.22	9 (3%) 39 34	42, 64, 110, 149	0
1	C	238/274 (86%)	0.08	10 (4%) 37 32	25, 50, 100, 137	0
1	D	227/274 (82%)	0.06	10 (4%) 35 30	29, 53, 97, 146	0
All	All	940/1096 (85%)	0.14	39 (4%) 38 32	25, 58, 102, 171	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	228	ASP	6.3
1	A	248	ILE	4.6
1	D	100	CYS	4.3
1	A	229	HIS	4.1
1	A	255	HIS	3.8
1	D	80	ARG	3.6
1	B	100	CYS	3.5
1	C	100	CYS	3.5
1	C	255	HIS	3.5
1	B	79	VAL	3.5
1	A	100	CYS	3.5
1	A	254	THR	3.2
1	A	99	ASN	3.1
1	C	253	THR	2.9
1	B	81	MET	2.9
1	C	252	ASN	2.7
1	A	247	THR	2.7
1	B	75	GLU	2.6
1	B	287	VAL	2.6
1	C	256	CYS	2.6
1	D	259	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	123	ASN	2.5
1	D	263	ASP	2.5
1	A	18	MET	2.5
1	D	122	ILE	2.5
1	C	173	THR	2.4
1	D	260	TYR	2.4
1	C	96	ASN	2.3
1	D	42	THR	2.3
1	A	230	GLU	2.3
1	B	42	THR	2.3
1	C	227	SER	2.2
1	B	174	LEU	2.1
1	C	99	ASN	2.1
1	C	75	GLU	2.1
1	D	79	VAL	2.1
1	B	172	ASP	2.1
1	D	290	LEU	2.0
1	B	121	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	GSP	D	301	32/32	0.95	0.16	-0.91	38,43,47,54	0
2	GSP	A	301	32/32	0.96	0.16	-1.00	31,38,43,44	0
2	GSP	C	301	32/32	0.96	0.14	-1.12	27,42,46,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	GSP	B	301	32/32	0.96	0.13	-1.47	35,47,54,56	0
3	MG	A	302	1/1	0.95	0.11	-	33,33,33,33	0
3	MG	D	302	1/1	0.95	0.12	-	49,49,49,49	0
3	MG	B	302	1/1	0.95	0.11	-	38,38,38,38	0
3	MG	C	302	1/1	0.98	0.15	-	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.