



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 09:56 am GMT

PDB ID : 4D0O
Title : AKAP13 (AKAP-Lbc) DH domain
Authors : Abdul Azeez, K.R.; Shrestha, L.; Krojer, T.; Allerston, C.; von Delft, F.;
Bountra, C.; Arrowsmith, C.; Edwards, A.M.; Knapp, S.; Klussmann, E.;
Elkins, J.M.
Deposited on : 2014-04-29
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	3666 (2.80-2.72)
Clashscore	112137	4174 (2.80-2.72)
Ramachandran outliers	110173	4103 (2.80-2.72)
Sidechain outliers	110143	4106 (2.80-2.72)
RSRZ outliers	101464	3697 (2.80-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div>6%</div> <div>• •</div> </div> </div>
1	B	244	<div> <div>5%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div>•</div> </div> </div>

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3692 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called A-KINASE ANCHOR PROTEIN 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	234	Total	C	N	O	S	0	0	0
			1826	1177	303	335	11			
1	B	240	Total	C	N	O	S	0	0	0
			1866	1200	311	344	11			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1968	GLU	-	EXPRESSION TAG	UNP Q12802
A	1969	ASN	-	EXPRESSION TAG	UNP Q12802
A	1970	LEU	-	EXPRESSION TAG	UNP Q12802
A	1971	TYR	-	EXPRESSION TAG	UNP Q12802
A	1972	PHE	-	EXPRESSION TAG	UNP Q12802
A	1973	GLN	-	EXPRESSION TAG	UNP Q12802
A	1974	SER	-	EXPRESSION TAG	UNP Q12802
A	1975	MET	-	EXPRESSION TAG	UNP Q12802
B	1968	GLU	-	EXPRESSION TAG	UNP Q12802
B	1969	ASN	-	EXPRESSION TAG	UNP Q12802
B	1970	LEU	-	EXPRESSION TAG	UNP Q12802
B	1971	TYR	-	EXPRESSION TAG	UNP Q12802
B	1972	PHE	-	EXPRESSION TAG	UNP Q12802
B	1973	GLN	-	EXPRESSION TAG	UNP Q12802
B	1974	SER	-	EXPRESSION TAG	UNP Q12802
B	1975	MET	-	EXPRESSION TAG	UNP Q12802


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: A-KINASE ANCHOR PROTEIN 13



• Molecule 1: A-KINASE ANCHOR PROTEIN 13



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.13Å 94.84Å 109.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.55 – 2.75 42.13 – 2.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (71.55-2.75) 99.9 (42.13-2.75)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.236 , 0.308 0.238 , 0.310	Depositor DCC
R_{free} test set	711 reflections (5.12%)	DCC
Wilson B-factor (Å ²)	49.6	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 35.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3692	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/1855	0.46	0/2501
1	B	0.33	0/1896	0.48	0/2558
All	All	0.32	0/3751	0.47	0/5059

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	2139	ARG	Peptide
1	B	2139	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1826	0	1766	9	0
1	B	1866	0	1795	10	0
All	All	3692	0	3561	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1983:ILE:HD13	1:A:2076:PHE:CE2	2.33	0.64
1:A:2152:GLN:HE21	1:B:2152:GLN:CG	2.18	0.57
1:A:2057:PHE:CE2	1:A:2061:LEU:HD11	2.41	0.55
1:B:2057:PHE:CE2	1:B:2061:LEU:HD11	2.42	0.55
1:A:2152:GLN:HE21	1:B:2152:GLN:HG3	1.71	0.54
1:A:2163:ARG:NH2	1:B:2197:ALA:HB1	2.24	0.53
1:A:2152:GLN:CG	1:B:2152:GLN:HE21	2.25	0.50
1:A:2152:GLN:HG3	1:B:2152:GLN:HE21	1.78	0.48
1:A:2007:MET:HE2	1:A:2007:MET:O	2.14	0.47
1:B:1968:GLU:O	1:B:1970:LEU:N	2.48	0.46
1:B:2096:ARG:O	1:B:2100:THR:OG1	2.25	0.45
1:B:2143:ILE:N	1:B:2144:PRO:CD	2.84	0.41
1:A:2143:ILE:N	1:A:2144:PRO:CD	2.84	0.40
1:B:2044:PRO:HG2	1:B:2104:PHE:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/244 (94%)	225 (98%)	3 (1%)	2 (1%)	20	49
1	B	238/244 (98%)	231 (97%)	4 (2%)	3 (1%)	14	37

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	468/488 (96%)	456 (97%)	7 (2%)	5 (1%)	17	43

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1969	ASN
1	A	2139	ARG
1	B	2139	ARG
1	A	2108	HIS
1	B	2108	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/228 (82%)	181 (96%)	7 (4%)	39	71
1	B	191/228 (84%)	187 (98%)	4 (2%)	59	85
All	All	379/456 (83%)	368 (97%)	11 (3%)	48	78

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2001	GLU
1	A	2007	MET
1	A	2048	GLU
1	A	2066	GLU
1	A	2080	ARG
1	A	2117	ASP
1	A	2180	GLN
1	B	2001	GLU
1	B	2048	GLU
1	B	2080	ARG
1	B	2180	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	2008	GLN
1	A	2012	HIS
1	A	2152	GLN
1	B	2152	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/244 (95%)	0.45	12 (5%) 29 23	37, 58, 85, 101	0
1	B	240/244 (98%)	0.38	11 (4%) 33 27	25, 44, 78, 116	0
All	All	474/488 (97%)	0.41	23 (4%) 30 24	25, 53, 84, 116	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2206	ASN	4.4
1	B	2207	GLU	4.2
1	B	2205	LEU	3.6
1	A	2167	CYS	3.5
1	A	2170	ASP	3.5
1	B	1986	SER	3.4
1	A	2069	VAL	3.1
1	B	2120	ALA	3.0
1	B	2122	ASP	2.8
1	B	2115	PHE	2.7
1	A	2076	PHE	2.5
1	B	2069	VAL	2.5
1	A	1996	VAL	2.4
1	A	2062	GLU	2.4
1	B	1985	ASP	2.3
1	A	1991	GLN	2.3
1	B	2038	MET	2.2
1	A	2117	ASP	2.2
1	B	2199	TYR	2.2
1	A	2176	GLU	2.2
1	A	2065	LYS	2.2
1	A	1985	ASP	2.1
1	A	2203	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.