



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 06:54 am GMT

PDB ID : 2D1Z  
Title : Crystal structure of catalytic-site mutant xylanase from *Streptomyces olivaceoviridis* E-86  
Authors : Suzuki, R.; Kuno, A.; Fujimoto, Z.; Ito, S.; Kawahara, S.I.; Kaneko, S.; Hasegawa, T.; Taira, K.  
Deposited on : 2005-09-02  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	<div> <div>6%</div> <div>87%</div> <div>11%</div> <div>••</div> </div>
1	B	436	<div> <div>12%</div> <div>84%</div> <div>14%</div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1442	-	-	-	X
3	GOL	A	1443	-	-	-	X
3	GOL	A	1444	-	-	-	X
3	GOL	A	1445	-	-	-	X
3	GOL	A	1446	-	-	-	X
3	GOL	B	1942	-	-	-	X
3	GOL	B	1943	-	-	-	X
3	GOL	B	1944	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ENDO-1,4-BETA-D-XYLANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	0	0
			3233	1988	588	641	16			
1	B	427	Total	C	N	O	S	0	0	0
			3233	1988	588	641	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	SER	ASN	ENGINEERED MUTATION	UNP Q7SI98
A	128	HIS	GLU	ENGINEERED MUTATION	UNP Q7SI98
B	627	SER	ASN	ENGINEERED MUTATION	UNP Q7SI98
B	628	HIS	GLU	ENGINEERED MUTATION	UNP Q7SI98

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	427	Total 427	O 427	0	0
4	B	474	Total 474	O 474	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ENDO-1,4-BETA-D-XYLANASE



#### • Molecule 1: ENDO-1,4-BETA-D-XYLANASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	78.51Å 94.06Å 139.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.03 – 1.60 47.03 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.03-1.60) 99.4 (47.03-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.01 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.191 , 0.207 0.191 , 0.207	Depositor DCC
$R_{free}$ test set	6894 reflections (5.08%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.1	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7438	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.29	0/3298	0.62	0/4473
1	B	0.29	0/3298	0.61	0/4473
All	All	0.29	0/6596	0.62	0/8946

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	669	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3233	0	3043	30	0
1	B	3233	0	3040	52	0
2	A	5	0	0	0	0
3	A	30	0	40	5	0
3	B	36	0	48	3	0
4	A	427	0	0	5	0
4	B	474	0	0	6	0
All	All	7438	0	6171	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:540:MET:HE3	1:B:541:VAL:N	1.88	0.87
1:B:540:MET:CE	1:B:579:ARG:HG3	2.07	0.84
1:B:540:MET:HE1	1:B:579:ARG:HG3	1.59	0.84
1:B:911:GLY:H	3:B:1946:GOL:H31	1.41	0.83
1:A:336:GLN:HB2	1:A:376:LYS:HE3	1.63	0.80
1:A:139:ARG:HB2	3:A:1442:GOL:H11	1.66	0.76
1:B:816:LYS:HE2	1:B:936:THR:HG21	1.68	0.76
1:A:421:GLN:HE22	3:A:1445:GOL:H32	1.53	0.73
1:B:540:MET:HE3	1:B:541:VAL:H	1.55	0.72
1:B:816:LYS:HG2	1:B:823:CYS:SG	2.36	0.65
1:B:910:VAL:HA	1:B:921:GLN:HE21	1.63	0.63
1:B:935:ARG:HA	1:B:936:THR:HB	1.79	0.63
1:A:238:ASP:HB2	1:A:280:PRO:HB2	1.81	0.63
1:B:549:ILE:HD12	4:B:2168:HOH:O	2.00	0.62
1:A:421:GLN:NE2	3:A:1445:GOL:H32	2.18	0.59
1:A:97:SER:O	1:A:100:THR:HG22	2.03	0.58
1:B:789:LYS:HD2	1:B:891:ARG:HH12	1.68	0.58
1:B:837:VAL:HG23	1:B:920:ILE:HB	1.85	0.57
1:A:127:SER:HA	1:A:170:ASN:O	2.05	0.57
1:A:1:ALA:HB1	4:A:1545:HOH:O	2.06	0.56
1:A:389:LYS:HE3	4:A:1585:HOH:O	2.05	0.56
1:B:936:THR:HG22	1:B:936:THR:OXT	2.05	0.56
1:B:738:ASP:HB2	1:B:780:PRO:HB2	1.87	0.56
1:A:411:GLY:H	3:A:1445:GOL:H31	1.71	0.54
1:B:754:CYS:SG	1:B:763:ILE:HD11	2.48	0.54
1:B:899:VAL:HG22	1:B:906:CYS:SG	2.47	0.54
1:B:676:ASN:HB3	1:B:679:TRP:CD2	2.43	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:720:THR:HG23	4:B:2298:HOH:O	2.09	0.53
1:B:789:LYS:HD2	1:B:891:ARG:NH1	2.24	0.52
1:B:778:ASP:O	1:B:779:THR:C	2.48	0.51
1:B:935:ARG:HA	1:B:936:THR:C	2.31	0.51
1:A:234:ILE:HD12	1:A:263:ILE:HG12	1.92	0.50
1:B:779:THR:N	1:B:780:PRO:HD3	2.27	0.50
1:A:55:GLN:HB2	1:A:58:GLN:HB2	1.92	0.50
1:A:337:VAL:HG23	1:A:420:ILE:HB	1.94	0.50
1:B:925:CYS:HA	1:B:931:GLN:OE1	2.11	0.50
1:A:142:ASN:O	1:A:146:THR:HG23	2.11	0.50
1:A:15:TYR:CG	1:A:40:MET:HG3	2.47	0.50
1:A:279:THR:N	1:A:280:PRO:HD3	2.27	0.50
1:B:627:SER:HA	1:B:670:ASN:O	2.12	0.49
1:A:278:ASP:O	1:A:279:THR:C	2.50	0.49
1:B:820:SER:HA	1:B:914:THR:HB	1.93	0.49
1:B:822:ARG:HG2	1:B:822:ARG:HH11	1.77	0.49
1:B:874:GLY:HA2	1:B:921:GLN:OE1	2.13	0.49
1:A:85:TRP:CZ3	1:A:87:SER:HB3	2.48	0.48
1:B:935:ARG:CA	1:B:936:THR:HB	2.43	0.48
1:B:910:VAL:HG22	1:B:921:GLN:NE2	2.28	0.48
1:A:190:ARG:O	1:A:194:GLN:HG3	2.14	0.48
1:A:80:GLY:HA3	1:A:123:TRP:CE3	2.48	0.48
1:B:911:GLY:H	3:B:1946:GOL:C3	2.17	0.48
1:B:909:ALA:HB3	1:B:930:ASN:HB2	1.96	0.48
1:B:694:GLN:HG3	4:B:2173:HOH:O	2.15	0.47
1:A:179:TRP:HE3	4:A:1682:HOH:O	1.97	0.47
1:A:1:ALA:HB3	1:A:7:ALA:HB1	1.96	0.47
1:B:501:ALA:HB3	4:B:2119:HOH:O	2.13	0.47
1:B:754:CYS:CB	1:B:763:ILE:HD11	2.45	0.47
1:A:436:THR:OXT	1:A:436:THR:HG22	2.15	0.46
1:B:827:PRO:HG3	1:B:838:GLN:HG2	1.97	0.46
1:B:580:GLY:HA3	1:B:623:TRP:CE3	2.51	0.46
1:B:816:LYS:HE2	1:B:936:THR:CG2	2.43	0.46
1:B:719:ARG:O	1:B:723:GLN:HG3	2.16	0.46
1:B:540:MET:HE2	1:B:579:ARG:HG3	1.95	0.45
1:A:426:SER:O	1:A:427:ASN:HB2	2.15	0.45
1:B:921:GLN:NE2	3:B:1946:GOL:H32	2.31	0.45
3:A:1446:GOL:H32	4:A:1660:HOH:O	2.16	0.45
1:A:141:SER:O	1:A:145:ARG:HG3	2.16	0.45
1:B:502:GLU:HB2	1:B:801:ASN:OD1	2.17	0.45
1:A:144:GLN:HE22	1:A:148:ASN:HA	1.82	0.45

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:831:THR:HG22	4:B:2306:HOH:O	2.16	0.44
1:B:845:ALA:O	1:B:849:GLN:HG2	2.18	0.43
1:A:432:ARG:HD3	4:A:1552:HOH:O	2.19	0.43
1:B:519:ALA:HB2	1:B:766:TRP:CE3	2.54	0.42
1:B:910:VAL:HA	1:B:921:GLN:NE2	2.32	0.42
1:B:672:TYR:HB3	1:B:705:GLN:NE2	2.34	0.42
1:B:842:CYS:HA	1:B:848:GLN:OE1	2.19	0.41
1:B:763:ILE:HD12	1:B:763:ILE:N	2.35	0.41
1:A:52:THR:O	1:A:59:PHE:HA	2.20	0.41
1:B:789:LYS:CD	1:B:891:ARG:NH1	2.84	0.41
1:A:172:TYR:HB3	1:A:205:GLN:NE2	2.35	0.41
1:A:168:CYS:HA	1:A:201:CYS:O	2.20	0.41
1:B:822:ARG:HE	1:B:915:ALA:HA	1.84	0.41
1:B:822:ARG:HG2	1:B:822:ARG:NH1	2.36	0.41
1:B:694:GLN:HG2	4:B:2253:HOH:O	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/436 (97%)	411 (97%)	12 (3%)	0	100	100
1	B	423/436 (97%)	411 (97%)	12 (3%)	0	100	100
All	All	846/872 (97%)	822 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	334/341 (98%)	329 (98%)	5 (2%)	70	48
1	B	334/341 (98%)	328 (98%)	6 (2%)	64	40
All	All	668/682 (98%)	657 (98%)	11 (2%)	68	45

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	MET
1	A	85	TRP
1	A	247	TYR
1	A	274	TRP
1	A	280	PRO
1	B	540	MET
1	B	585	TRP
1	B	601	LEU
1	B	747	TYR
1	B	780	PRO
1	B	927	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	55	GLN
1	A	74	ASN
1	A	223	GLN
1	A	284	ASN
1	A	421	GLN
1	B	511	GLN
1	B	828	ASN
1	B	921	GLN
1	B	927	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1441	-	4,4,4	0.31	0	6,6,6	0.09	0
3	GOL	A	1442	-	5,5,5	0.15	0	5,5,5	0.29	0
3	GOL	A	1443	-	5,5,5	0.13	0	5,5,5	0.29	0
3	GOL	A	1444	-	5,5,5	0.15	0	5,5,5	0.29	0
3	GOL	A	1445	-	5,5,5	0.19	0	5,5,5	0.31	0
3	GOL	A	1446	-	5,5,5	0.20	0	5,5,5	0.34	0
3	GOL	B	1941	-	5,5,5	0.15	0	5,5,5	0.27	0
3	GOL	B	1942	-	5,5,5	0.15	0	5,5,5	0.27	0
3	GOL	B	1943	-	5,5,5	0.14	0	5,5,5	0.29	0
3	GOL	B	1944	-	5,5,5	0.13	0	5,5,5	0.29	0
3	GOL	B	1945	-	5,5,5	0.16	0	5,5,5	0.30	0
3	GOL	B	1946	-	5,5,5	0.16	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1441	-	-	0/0/0/0	0/0/0/0
3	GOL	A	1442	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1443	-	-	0/4/4/4	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1444	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1445	-	-	0/4/4/4	0/0/0/0
3	GOL	A	1446	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1941	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1942	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1943	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1944	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1945	-	-	0/4/4/4	0/0/0/0
3	GOL	B	1946	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1442	GOL	1	0
3	A	1445	GOL	3	0
3	A	1446	GOL	1	0
3	B	1946	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	427/436 (97%)	0.37	27 (6%) 21 20	9, 15, 33, 44	0
1	B	427/436 (97%)	0.57	52 (12%) 5 4	8, 13, 46, 58	0
All	All	854/872 (97%)	0.47	79 (9%) 9 9	8, 14, 39, 58	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	819	GLY	10.9
1	A	436	THR	10.6
1	B	936	THR	10.0
1	A	303	GLY	8.2
1	B	818	VAL	7.8
1	B	501	ALA	7.2
1	B	822	ARG	7.2
1	B	844	SER	7.1
1	B	935	ARG	6.6
1	B	927	ASN	6.5
1	B	910	VAL	6.1
1	B	814	GLN	5.6
1	B	842	CYS	5.6
1	A	92	TRP	5.5
1	B	914	THR	5.4
1	B	934	THR	5.3
1	B	820	SER	5.0
1	B	912	GLY	4.8
1	B	816	LYS	4.7
1	B	923	TYR	4.6
1	B	823	CYS	4.5
1	B	911	GLY	4.5
1	B	821	GLY	4.4
1	A	1	ALA	4.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	843	HIS	4.3
1	B	928	GLY	4.2
1	A	344	SER	4.2
1	B	915	ALA	4.2
1	B	839	LEU	4.2
1	A	100	THR	4.0
1	A	97	SER	4.0
1	B	932	ARG	4.0
1	B	845	ALA	4.0
1	B	894	SER	3.9
1	B	918	THR	3.9
1	B	925	CYS	3.8
1	B	929	SER	3.6
1	A	140	ASP	3.6
1	A	91	GLY	3.5
1	A	145	ARG	3.5
1	B	926	SER	3.4
1	A	56	ARG	3.4
1	B	849	GLN	3.3
1	B	841	ASP	3.2
1	B	840	TYR	3.2
1	A	95	SER	3.2
1	A	435	ARG	3.1
1	A	147	GLY	3.0
1	B	817	GLY	3.0
1	A	58	GLN	2.9
1	B	909	ALA	2.9
1	B	896	GLY	2.9
1	A	94	GLN	2.9
1	B	917	GLY	2.7
1	B	916	ASN	2.7
1	B	828	ASN	2.6
1	A	179	TRP	2.6
1	B	874	GLY	2.6
1	A	133	ASP	2.6
1	A	98	GLY	2.6
1	A	143	LEU	2.6
1	B	924	SER	2.5
1	B	930	ASN	2.5
1	B	876	LYS	2.5
1	A	96	LEU	2.5
1	B	824	LEU	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	89	GLN	2.4
1	B	813	GLY	2.4
1	B	922	LEU	2.2
1	B	895	ASP	2.2
1	A	86	HIS	2.2
1	A	141	SER	2.2
1	A	101	LEU	2.2
1	B	919	LEU	2.2
1	A	146	THR	2.2
1	A	85	TRP	2.2
1	B	906	CYS	2.1
1	B	899	VAL	2.1
1	B	908	ASP	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	1943	6/6	0.77	0.23	16.35	44,45,45,46	0
3	GOL	B	1942	6/6	0.70	0.23	15.15	41,43,43,43	0
3	GOL	A	1445	6/6	0.76	0.24	12.06	28,33,34,40	0
3	GOL	A	1446	6/6	0.73	0.26	6.44	21,27,32,33	0
3	GOL	A	1444	6/6	0.68	0.20	6.23	36,37,37,38	0
3	GOL	A	1443	6/6	0.85	0.12	2.67	21,23,24,25	0
3	GOL	B	1944	6/6	0.90	0.10	2.08	19,21,21,22	0
3	GOL	A	1442	6/6	0.38	0.34	2.01	50,52,52,53	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	A	1441	5/5	0.97	0.20	1.30	24,26,31,31	0
3	GOL	B	1945	6/6	0.84	0.26	1.00	45,45,46,46	0
3	GOL	B	1946	6/6	0.83	0.23	-0.24	42,43,43,44	0
3	GOL	B	1941	6/6	0.72	0.37	-	35,42,44,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.