



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 07:31 am GMT

PDB ID : 4D1O
Title : Structure of human endothelial nitric oxide synthase heme domain with L-Arg bound
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-05-02
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5868 (1.84-1.80)
Clashscore	112137	6856 (1.84-1.80)
Ramachandran outliers	110173	6780 (1.84-1.80)
Sidechain outliers	110143	6780 (1.84-1.80)
RSRZ outliers	101464	5947 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>5%</div> <div>9%</div> </div> </div>
1	B	440	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>7%</div> <div>9%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ARG	B	700	-	-	-	X
4	ACT	B	860	-	-	-	X
5	BTB	A	870	-	-	-	X
6	GOL	A	880	-	-	-	X
6	GOL	B	880	-	-	-	X

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 7442 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	3	0
			3212	2046	563	587	16			
1	B	402	Total	C	N	O	S	0	6	0
			3235	2060	567	591	17			

- Molecule 2 is ARGinine (three-letter code: ARG) (formula: $C_6H_{15}N_4O_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			12	6	4	2		
2	B	1	Total	C	N	O	0	0
			12	6	4	2		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
3	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (three-letter code: BTB) (formula: $C_8H_{19}NO_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		

- Molecule 8 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C₉H₁₅N₅O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			17	9	5	3		
8	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 9 is GADOLINIUM ATOM (three-letter code: GD) (formula: Gd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total	Gd	0	0
			1	1		
9	A	1	Total	Gd	0	0
			1	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	379	Total	O	0	0
			379	379		
10	B	387	Total	O	0	0
			387	387		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.40Å 110.09Å 153.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.39 – 1.82 48.39 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.39-1.82) 99.6 (48.39-1.82)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.82Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.2_1309)	Depositor
R, R_{free}	0.155 , 0.192 0.154 , 0.192	Depositor DCC
R_{free} test set	4704 reflections (4.94%)	DCC
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.372	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7442	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, H4B, BTB, GD, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/3313	0.50	0/4515
1	B	0.36	0/3342	0.50	0/4554
All	All	0.36	0/6655	0.50	0/9069

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3212	0	3114	19	0
1	B	3235	0	3143	18	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
3	A	43	0	30	1	0
3	B	43	0	30	1	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
5	A	28	0	35	4	0
5	B	28	0	34	1	0
6	A	12	0	16	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	6	0	8	0	0
7	A	1	0	0	0	0
8	A	17	0	15	1	0
8	B	17	0	15	1	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	A	379	0	0	5	0
10	B	387	0	0	3	0
All	All	7442	0	6470	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:ARG:NH2	10:A:2180:HOH:O	2.24	0.71
1:A:89:GLN:NE2	1:A:470:SER:H	1.92	0.67
1:A:89:GLN:HE21	1:A:470:SER:H	1.44	0.64
5:A:870:BTB:O3	10:A:2297:HOH:O	2.16	0.61
1:A:128:ARG:NH2	1:A:154:GLU:OE2	2.34	0.56
1:A:167:GLU:OE2	6:A:881:GOL:O2	2.23	0.56
1:A:86:ALA:O	1:B:97:ARG:NH2	2.39	0.56
1:A:382:CYS:HA	5:A:870:BTB:H41	1.88	0.55
1:B:250:ARG:NH1	10:B:2228:HOH:O	2.37	0.54
1:B:386:ASP:OD1	1:B:388:ARG:HD3	2.07	0.54
3:A:500:HEM:HBB2	3:A:500:HEM:HHC	1.92	0.51
1:B:119:ALA:N	1:B:121:GLU:OE2	2.44	0.51
1:A:91:ASP:OD1	10:A:2024:HOH:O	2.19	0.51
1:A:133:GLN:NE2	10:A:2070:HOH:O	2.45	0.49
1:A:150:LEU:O	1:A:154:GLU:HG3	2.12	0.49
1:B:200[A]:ASP:OD1	10:B:2157:HOH:O	2.20	0.48
10:A:2349:HOH:O	1:B:395:LYS:NZ	2.45	0.48
1:A:93:PRO:HB3	1:A:106:PRO:HB3	1.95	0.48
3:B:500:HEM:HHC	3:B:500:HEM:HBB2	1.96	0.46
5:B:871:BTB:H32	5:B:871:BTB:H71	1.60	0.46
1:B:447:TRP:HA	8:B:1481:H4B:N1	2.31	0.46
1:B:290:PRO:HB3	1:B:304:LEU:HD23	1.97	0.45
1:A:453:SER:HB3	1:A:456:LEU:HD12	1.99	0.45
5:A:871:BTB:H72	5:A:871:BTB:H61	1.71	0.45
1:B:121:GLU:CD	1:B:121:GLU:H	2.19	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:364:THR:O	1:B:368:CYS:HB2	2.17	0.44
5:A:871:BTB:O8	5:A:871:BTB:H41	2.19	0.43
1:A:207:MET:HG3	1:A:231:PHE:CE1	2.54	0.43
1:A:447:TRP:HA	8:A:1481:H4B:N1	2.33	0.43
1:B:247:GLN:HB2	1:B:250:ARG:HG2	2.01	0.42
1:B:233:GLN:HB3	1:B:348:PHE:CE2	2.55	0.42
1:B:342:GLU:OE1	1:B:474:ARG:NH1	2.46	0.42
1:B:119:ALA:HB3	1:B:120:PRO:HD3	2.02	0.42
1:A:386:ASP:OD1	1:A:388:ARG:HG2	2.20	0.42
1:B:453:SER:HB3	1:B:456:LEU:HD12	2.02	0.41
1:B:279:TRP:HB2	1:B:302:LEU:HD21	2.01	0.41
1:B:221:ARG:NH1	10:B:2184:HOH:O	2.54	0.40
1:A:183:ARG:HD3	1:A:447:TRP:CD2	2.57	0.40
1:A:178:TRP:CE3	1:A:190:TRP:HA	2.56	0.40
1:B:238:ARG:HD3	1:B:296:PRO:HG2	2.02	0.40
1:A:207:MET:HE2	1:A:207:MET:HB2	1.97	0.40
1:A:470:SER:HA	1:A:471:PRO:C	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/440 (91%)	393 (98%)	6 (2%)	1 (0%)	44	28
1	B	404/440 (92%)	399 (99%)	5 (1%)	0	100	100
All	All	804/880 (91%)	792 (98%)	11 (1%)	1 (0%)	55	38

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	89	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/373 (92%)	337 (98%)	6 (2%)	66	55
1	B	347/373 (93%)	341 (98%)	6 (2%)	66	55
All	All	690/746 (92%)	678 (98%)	12 (2%)	64	55

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	GLN
1	A	98	ARG
1	A	164	GLN
1	A	167	GLU
1	A	207	MET
1	A	216	LYS
1	B	67	LYS
1	B	90	GLN
1	B	98	ARG
1	B	257	GLN
1	B	329	ARG
1	B	388	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 3 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	H4B	A	1481	-	14,18,18	0.84	0	12,26,26	2.36	5 (41%)
3	HEM	A	500	1	28,50,50	2.27	8 (28%)	17,82,82	1.76	3 (17%)
2	ARG	A	700	-	6,11,11	0.42	0	5,13,13	1.03	0
4	ACT	A	860	-	1,3,3	1.24	0	0,3,3	0.00	-
5	BTB	A	870	9	13,13,13	0.55	0	9,16,16	0.90	0
5	BTB	A	871	-	13,13,13	0.51	0	9,16,16	0.69	0
6	GOL	A	880	-	5,5,5	0.43	0	5,5,5	0.15	0
6	GOL	A	881	-	5,5,5	0.31	0	5,5,5	0.38	0
8	H4B	B	1481	-	14,18,18	0.86	1 (7%)	12,26,26	2.24	4 (33%)
3	HEM	B	500	1	28,50,50	2.29	7 (25%)	17,82,82	2.05	5 (29%)
2	ARG	B	700	-	6,11,11	0.44	0	5,13,13	0.94	0
4	ACT	B	860	-	1,3,3	1.09	0	0,3,3	0.00	-
5	BTB	B	870	9	13,13,13	0.50	0	9,16,16	0.55	0
5	BTB	B	871	-	13,13,13	0.55	0	9,16,16	1.08	1 (11%)
6	GOL	B	880	-	5,5,5	0.37	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	H4B	A	1481	-	-	0/8/17/17	0/2/2/2
3	HEM	A	500	1	-	0/6/54/54	0/0/8/8
2	ARG	A	700	-	-	0/7/11/11	0/0/0/0
4	ACT	A	860	-	-	0/0/0/0	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	A	870	9	-	0/21/21/21	0/0/0/0
5	BTB	A	871	-	-	0/21/21/21	0/0/0/0
6	GOL	A	880	-	-	0/4/4/4	0/0/0/0
6	GOL	A	881	-	-	0/4/4/4	0/0/0/0
8	H4B	B	1481	-	-	0/8/17/17	0/2/2/2
3	HEM	B	500	1	-	0/6/54/54	0/0/8/8
2	ARG	B	700	-	-	0/7/11/11	0/0/0/0
4	ACT	B	860	-	-	0/0/0/0	0/0/0/0
5	BTB	B	870	9	-	0/21/21/21	0/0/0/0
5	BTB	B	871	-	-	0/21/21/21	0/0/0/0
6	GOL	B	880	-	-	0/4/4/4	0/0/0/0

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	500	HEM	C3B-C2B	-4.71	1.34	1.40
3	A	500	HEM	C3B-C2B	-4.59	1.34	1.40
3	B	500	HEM	C3C-C2C	-3.65	1.35	1.40
3	A	500	HEM	C3C-C2C	-3.41	1.35	1.40
8	B	1481	H4B	C4-C4A	-2.12	1.38	1.41
3	A	500	HEM	C1D-ND	2.01	1.40	1.36
3	B	500	HEM	C4C-NC	2.04	1.39	1.36
3	A	500	HEM	C1C-NC	2.56	1.39	1.36
3	B	500	HEM	C4D-ND	3.23	1.40	1.36
3	A	500	HEM	C4D-ND	3.30	1.40	1.36
3	B	500	HEM	C3B-CAB	3.86	1.55	1.47
3	A	500	HEM	C3B-CAB	3.95	1.55	1.47
3	A	500	HEM	C3C-CAC	4.00	1.55	1.47
3	B	500	HEM	C3C-CAC	4.09	1.55	1.47
3	A	500	HEM	C3D-C2D	4.91	1.52	1.37
3	B	500	HEM	C3D-C2D	4.99	1.52	1.37

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	500	HEM	CBA-CAA-C2A	-4.93	103.06	112.48
3	A	500	HEM	CBA-CAA-C2A	-4.81	103.29	112.48
3	B	500	HEM	CBD-CAD-C3D	-4.02	104.80	112.47
8	A	1481	H4B	N3-C2-N1	-3.38	119.97	125.45
8	B	1481	H4B	N3-C2-N1	-3.32	120.07	125.45
3	A	500	HEM	CBD-CAD-C3D	-2.99	106.76	112.47
3	B	500	HEM	CMA-C3A-C4A	-2.61	124.46	128.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	871	BTB	O1-C1-C2	-2.46	104.44	111.37
3	A	500	HEM	C1D-C2D-C3D	-2.43	105.31	107.00
3	B	500	HEM	C1D-C2D-C3D	-2.36	105.36	107.00
8	A	1481	H4B	C6-C7-N8	-2.31	107.35	111.01
3	B	500	HEM	C4A-C3A-C2A	2.05	108.42	107.00
8	B	1481	H4B	C2-N1-C8A	2.76	120.73	114.51
8	A	1481	H4B	C2-N1-C8A	2.90	121.04	114.51
8	B	1481	H4B	C4-N3-C2	3.54	121.16	116.06
8	A	1481	H4B	C4-N3-C2	3.64	121.29	116.06
8	B	1481	H4B	C4-C4A-C8A	3.97	118.15	114.56
8	A	1481	H4B	C4-C4A-C8A	3.98	118.17	114.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	1481	H4B	1	0
3	A	500	HEM	1	0
5	A	870	BTB	2	0
5	A	871	BTB	2	0
6	A	881	GOL	1	0
8	B	1481	H4B	1	0
3	B	500	HEM	1	0
5	B	871	BTB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/440 (91%)	-0.06	9 (2%) 62 58	14, 27, 60, 97	0
1	B	402/440 (91%)	-0.19	9 (2%) 62 58	14, 25, 57, 92	0
All	All	803/880 (91%)	-0.13	18 (2%) 62 58	14, 26, 58, 97	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	119	ALA	3.6
1	B	119	ALA	3.5
1	B	257	GLN	3.0
1	B	259	GLY	2.9
1	A	304	LEU	2.9
1	B	67	LYS	2.8
1	A	237	GLY	2.7
1	A	120	PRO	2.6
1	B	122	GLN	2.4
1	B	89	GLN	2.2
1	A	89	GLN	2.2
1	A	121	GLU	2.2
1	A	202	ARG	2.1
1	B	121	GLU	2.1
1	B	388	ARG	2.1
1	A	142	GLY	2.1
1	A	412	LEU	2.1
1	B	120	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	BTB	A	870	14/14	0.91	0.15	5.69	23,70,85,88	0
6	GOL	A	880	6/6	0.91	0.23	5.25	31,41,53,63	0
6	GOL	B	880	6/6	0.94	0.22	4.60	30,39,59,82	0
4	ACT	B	860	4/4	0.87	0.16	3.12	33,44,53,59	0
2	ARG	B	700	12/12	0.96	0.15	2.22	18,22,24,24	0
5	BTB	B	870	14/14	0.93	0.12	1.68	21,25,34,35	0
3	HEM	B	500	43/43	0.99	0.13	1.46	11,15,20,28	0
2	ARG	A	700	12/12	0.97	0.13	1.07	18,22,25,29	0
4	ACT	A	860	4/4	0.92	0.11	0.92	30,40,48,71	0
3	HEM	A	500	43/43	0.98	0.13	0.80	13,18,23,25	0
8	H4B	B	1481	17/17	0.98	0.13	0.66	13,16,19,21	0
8	H4B	A	1481	17/17	0.97	0.11	-0.19	13,18,20,25	0
7	ZN	A	900	1/1	1.00	0.05	-3.65	19,19,19,19	0
9	GD	A	1482	1/1	0.98	0.08	-	53,53,53,53	1
6	GOL	A	881	6/6	0.96	0.29	-	12,39,64,74	0
9	GD	B	1482	1/1	1.00	0.09	-	21,21,21,21	0
5	BTB	A	871	14/14	0.81	0.18	-	32,76,85,87	0
5	BTB	B	871	14/14	0.86	0.25	-	39,64,85,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.