



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:04 pm GMT

PDB ID : 3D36
Title : How to Switch Off a Histidine Kinase: Crystal Structure of Geobacillus stearothermophilus KinB with the Inhibitor Sda
Authors : Bick, M.J.; Lamour, V.; Rajashankar, K.R.; Gordiyenko, Y.; Robinson, C.V.; Darst, S.A.
Deposited on : 2008-05-09
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	8396 (2.04-2.00)
Clashscore	112137	9678 (2.04-2.00)
Ramachandran outliers	110173	9566 (2.04-2.00)
Sidechain outliers	110143	9565 (2.04-2.00)
RSRZ outliers	101464	8490 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	
1	B	244	
2	C	46	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MPD	C	701	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4224 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sporulation kinase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1687	1068	296	313	10			
1	B	221	Total	C	N	O	S	0	0	0
			1720	1088	303	319	10			

- Molecule 2 is a protein called Sporulation kinase inhibitor Sda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	42	Total	C	N	O	S	0	0	0
			356	230	59	66	1			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
4	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			8	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			8	6	2		
5	B	1	Total	C	O	0	0
			8	6	2		
5	C	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	123	Total	O	0	0
			123	123		
6	B	209	Total	O	0	0
			209	209		
6	C	41	Total	O	0	0
			41	41		

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	66.11Å 66.11Å 315.57Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.41 – 2.03 42.41 – 2.03	Depositor EDS
% Data completeness (in resolution range)	98.9 (42.41-2.03) 98.9 (42.41-2.03)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.200 , 0.215 0.282 , 0.290	Depositor DCC
R_{free} test set	2621 reflections (5.01%)	DCC
Wilson B-factor (Å ²)	33.7	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4224	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, MPD, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	1/1710 (0.1%)	0.50	0/2310
1	B	0.42	1/1744 (0.1%)	0.53	0/2355
2	C	0.41	0/361	0.51	0/481
All	All	0.41	2/3815 (0.1%)	0.51	0/5146

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	359	LYS	CE-NZ	8.69	1.70	1.49
1	B	317	CYS	CB-SG	-5.35	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1687	0	1754	10	0
1	B	1720	0	1785	11	0
2	C	356	0	374	5	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	27	0	12	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	27	0	12	0	0
5	B	24	0	42	2	0
5	C	8	0	14	0	0
6	A	123	0	0	1	0
6	B	209	0	0	3	0
6	C	41	0	0	0	0
All	All	4224	0	3993	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:359:LYS:CE	1:A:359:LYS:NZ	1.70	1.49
5:B:702:MPD:H51	5:B:703:MPD:H51	1.70	0.73
1:A:231:GLU:HG2	2:C:15:LYS:HB3	1.82	0.61
1:A:225:PHE:HZ	2:C:1:MET:HE3	1.67	0.60
1:A:365:LEU:HD11	1:A:382:MET:HG3	1.86	0.57
1:B:245:ILE:HG23	6:B:847:HOH:O	2.07	0.55
1:A:292:ASN:O	1:A:293:MET:HB3	2.07	0.54
1:B:383:MET:O	1:B:387:ARG:HG3	2.09	0.52
1:B:248:GLU:OE2	6:B:847:HOH:O	2.21	0.46
1:B:224:GLY:HA3	5:B:700:MPD:H11	1.97	0.46
1:A:223:ARG:HG3	1:B:247:ILE:HD12	1.97	0.46
1:B:273:LYS:HG2	1:B:306:SER:OG	2.16	0.45
1:A:292:ASN:O	1:A:293:MET:CB	2.65	0.45
1:A:261:LEU:O	1:B:212:SER:HB3	2.17	0.44
2:C:30:GLU:O	2:C:34:GLN:HG3	2.17	0.44
2:C:13:TYR:HB2	2:C:29:ILE:HG21	2.00	0.43
1:A:349:ARG:HA	1:A:408:VAL:O	2.18	0.43
1:B:285:ASP:OD1	1:B:288:ARG:NH1	2.52	0.41
2:C:13:TYR:CZ	2:C:17:LYS:HD2	2.55	0.41
1:B:277:LYS:HE2	1:B:281:GLU:OE2	2.20	0.41
1:A:359:LYS:NZ	6:A:616:HOH:O	2.47	0.41
1:B:240:ARG:NH2	6:B:842:HOH:O	2.53	0.41
1:B:279:GLU:OE2	1:B:311:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/244 (87%)	209 (98%)	4 (2%)	0	100	100
1	B	217/244 (89%)	214 (99%)	2 (1%)	1 (0%)	32	25
2	C	40/46 (87%)	40 (100%)	0	0	100	100
All	All	470/534 (88%)	463 (98%)	6 (1%)	1 (0%)	51	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	207	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/206 (89%)	182 (100%)	1 (0%)	91	93
1	B	186/206 (90%)	182 (98%)	4 (2%)	57	58
2	C	41/45 (91%)	41 (100%)	0	100	100
All	All	410/457 (90%)	405 (99%)	5 (1%)	75	79

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	319	LEU
1	B	194	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	219	LEU
1	B	306	SER
1	B	375	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ADP	A	500	3	25,29,29	1.05	1 (4%)	24,45,45	1.67	2 (8%)
4	ADP	B	500	3	25,29,29	1.01	1 (4%)	24,45,45	1.80	3 (12%)
5	MPD	B	700	-	7,7,7	0.25	0	9,10,10	0.23	0
5	MPD	B	702	-	7,7,7	0.27	0	9,10,10	0.30	0
5	MPD	B	703	-	7,7,7	0.23	0	9,10,10	0.26	0
5	MPD	C	701	-	7,7,7	0.26	0	9,10,10	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	500	3	-	0/12/32/32	0/3/3/3
4	ADP	B	500	3	-	0/12/32/32	0/3/3/3
5	MPD	B	700	-	-	0/5/5/5	0/0/0/0
5	MPD	B	702	-	-	0/5/5/5	0/0/0/0
5	MPD	B	703	-	-	0/5/5/5	0/0/0/0
5	MPD	C	701	-	-	0/5/5/5	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	500	ADP	C5-C4	3.08	1.47	1.40
4	A	500	ADP	C5-C4	3.16	1.47	1.40

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	500	ADP	N3-C2-N1	-7.28	122.52	128.86
4	A	500	ADP	N3-C2-N1	-6.61	123.10	128.86
4	A	500	ADP	C4-C5-N7	-2.75	106.75	109.41
4	B	500	ADP	C4-C5-N7	-2.27	107.22	109.41
4	B	500	ADP	C2-N1-C6	2.16	122.56	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	700	MPD	1	0
5	B	702	MPD	1	0
5	B	703	MPD	1	0

5.7 Other polymers

There are no such residues in this entry.









5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/244 (88%)	1.12	54 (24%)  	34, 39, 46, 50	0
1	B	221/244 (90%)	0.71	27 (12%)  	35, 40, 44, 54	0
2	C	42/46 (91%)	0.66	6 (14%)  	36, 39, 48, 54	0
All	All	480/534 (89%)	0.89	87 (18%)  	34, 39, 46, 54	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293	MET	6.1
1	A	303	ALA	5.1
1	B	338	TYR	5.0
1	A	415	ALA	4.9
2	C	42	ILE	4.6
1	A	191	GLY	4.6
1	A	416	SER	4.2
1	A	345	ARG	4.1
1	B	255	ALA	4.1
1	B	257	ILE	3.9
1	A	343	ASN	3.9
1	A	404	LYS	3.8
1	A	331	ASN	3.8
1	B	247	ILE	3.8
1	A	288	ARG	3.8
1	A	338	TYR	3.7
1	A	414	LEU	3.7
1	A	402	ILE	3.7
1	A	305	PHE	3.7
1	A	344	GLY	3.7
1	B	256	ILE	3.6
1	B	250	LEU	3.6
1	A	296	VAL	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	341	ILE	3.4
1	A	282	ARG	3.2
1	B	261	LEU	3.1
1	A	403	HIS	3.1
1	B	251	ASP	3.0
1	A	330	PRO	3.0
1	A	339	VAL	3.0
1	B	416	SER	3.0
1	A	306	SER	2.9
2	C	40	HIS	2.9
1	A	215	ILE	2.9
1	B	206	HIS	2.9
1	B	253	ALA	2.9
1	B	380	LEU	2.8
1	A	326	ILE	2.8
1	A	295	CYS	2.7
1	A	219	LEU	2.7
1	B	207	LEU	2.7
1	B	374	GLY	2.7
2	C	34	GLN	2.7
1	B	215	ILE	2.7
1	A	207	LEU	2.7
1	A	360	GLU	2.6
1	B	211	ILE	2.6
1	A	273	LYS	2.6
1	A	302	LEU	2.6
1	A	274	LEU	2.6
1	A	300	ALA	2.5
1	A	193	HIS	2.5
1	B	258	THR	2.5
1	B	371	THR	2.5
1	A	292	ASN	2.5
1	A	334	THR	2.4
1	A	278	LEU	2.4
1	A	336	GLN	2.4
1	A	332	GLY	2.4
1	A	401	GLU	2.3
2	C	21	LEU	2.3
1	B	260	TYR	2.3
1	B	378	THR	2.2
1	A	304	PRO	2.2
1	B	317	CYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	3	HIS	2.2
1	A	275	ASN	2.2
1	A	328	ALA	2.2
1	A	221	ALA	2.2
1	B	246	ALA	2.2
1	B	218	PRO	2.2
1	A	222	ALA	2.1
1	B	377	GLY	2.1
1	A	308	ILE	2.1
1	A	299	GLN	2.1
1	A	405	GLY	2.1
1	A	355	VAL	2.1
1	A	226	ILE	2.1
1	A	220	THR	2.1
1	A	218	PRO	2.0
2	C	19	LEU	2.0
1	A	213	HIS	2.0
1	B	200	LYS	2.0
1	B	252	ARG	2.0
1	B	384	VAL	2.0
1	A	256	ILE	2.0
1	A	346	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	MPD	C	701	8/8	0.76	0.29	6.40	62,62,63,64	0
5	MPD	B	700	8/8	0.80	0.23	1.98	58,58,59,59	0
5	MPD	B	703	8/8	0.88	0.21	0.88	43,45,47,47	0
3	MG	B	477	1/1	0.76	0.23	0.40	40,40,40,40	0
4	ADP	B	500	27/27	0.97	0.21	0.30	35,37,39,41	0
5	MPD	B	702	8/8	0.90	0.17	-0.50	38,39,42,42	0
4	ADP	A	500	27/27	0.93	0.11	-0.63	35,36,37,37	0
3	MG	A	478	1/1	0.83	0.09	-2.42	41,41,41,41	0

6.5 Other polymers

There are no such residues in this entry.