



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 03:17 am GMT

PDB ID : 2D51  
Title : Pentaketide chromone synthase (M207G mutant)  
Authors : Kohno, T.; Morita, H.  
Deposited on : 2005-10-27  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	<div> <div>5%</div> <div> <div></div> <div>96%</div> <div></div> </div> <div></div> </div>
1	B	406	<div> <div>7%</div> <div> <div></div> <div>94%</div> <div></div> </div> <div></div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6873 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called pentaketide chromone synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	405	Total	C	N	O	S	0	8	0
			3183	2025	537	595	26			
1	B	395	Total	C	N	O	S	0	4	0
			3098	1978	523	572	25			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP Q58VP7
A	2	PRO	-	CLONING ARTIFACT	UNP Q58VP7
A	3	GLY	-	CLONING ARTIFACT	UNP Q58VP7
A	177	CSD	CYS	MODIFIED RESIDUE	UNP Q58VP7
A	210	GLY	MET	ENGINEERED	UNP Q58VP7
B	1	GLY	-	CLONING ARTIFACT	UNP Q58VP7
B	2	PRO	-	CLONING ARTIFACT	UNP Q58VP7
B	3	GLY	-	CLONING ARTIFACT	UNP Q58VP7
B	177	CSD	CYS	MODIFIED RESIDUE	UNP Q58VP7
B	210	GLY	MET	ENGINEERED	UNP Q58VP7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	324	Total	O	0	0
			324	324		
2	B	268	Total	O	0	0
			268	268		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: pentaketide chromone synthase



- Molecule 1: pentaketide chromone synthase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.32Å 89.05Å 70.67Å 90.00° 95.50° 90.00°	Depositor
Resolution (Å)	30.00 – 1.60 29.71 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (30.00-1.60) 97.7 (29.71-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.64 (at 1.60Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.192 , 0.208 0.199 , 0.215	Depositor DCC
$R_{free}$ test set	5939 reflections (5.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.0	Xtriage
Anisotropy	0.574	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 36.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6873	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.60	0/3246	0.70	0/4400
1	B	0.60	0/3160	0.69	0/4284
All	All	0.60	0/6406	0.70	0/8684

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	170	TYR	Sidechain
1	B	170	TYR	Sidechain

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3183	0	3180	5	0
1	B	3098	0	3107	5	0
2	A	324	0	0	0	0
2	B	268	0	0	0	0
All	All	6873	0	6287	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:205:GLU:HG3	1:A:354:VAL:HB	1.82	0.61
1:B:205:GLU:HG3	1:B:354:VAL:HB	1.81	0.60
1:B:143:CYS:SG	1:B:179:ALA:HB1	2.44	0.57
1:A:143:CYS:SG	1:A:179:ALA:HB1	2.48	0.53
1:B:286:MET:HG3	1:B:328:GLN:HE22	1.77	0.47
1:B:386:LEU:HB3	1:B:398:ILE:HB	1.97	0.46
1:A:386:LEU:HB3	1:A:398:ILE:HB	1.99	0.45
1:A:72:ILE:O	1:A:76:THR:HG23	2.20	0.41
1:A:255:THR:HG22	1:A:398:ILE:HG23	2.03	0.41
1:B:341:ARG:O	1:B:345:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	410/406 (101%)	399 (97%)	10 (2%)	1 (0%)	51	27
1	B	396/406 (98%)	387 (98%)	8 (2%)	1 (0%)	44	22
All	All	806/812 (99%)	786 (98%)	18 (2%)	2 (0%)	44	27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	354	VAL
1	B	354	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	345/338 (102%)	341 (99%)	4 (1%)	75	58
1	B	333/338 (98%)	329 (99%)	4 (1%)	75	58
All	All	678/676 (100%)	670 (99%)	8 (1%)	75	58

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	299	VAL
1	A	323	ARG
1	A	366	ARG
1	A	405	LEU
1	B	64	GLU
1	B	215	ASN
1	B	286	MET
1	B	366	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	328	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.



## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSD	A	177	1	4,7,8	1.17	1 (25%)	2,8,10	1.50	0
1	CSD	B	177	1	4,7,8	1.17	1 (25%)	2,8,10	1.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	177	1	-	1/2/6/8	0/0/0/0
1	CSD	B	177	1	-	1/2/6/8	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	177	CSD	CA-C	2.02	1.52	1.50
1	A	177	CSD	CA-C	2.05	1.52	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	177	CSD	CA-CB-SG-OD1
1	A	177	CSD	CA-CB-SG-OD1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	404/406 (99%)	0.26	22 (5%) 26 25	8, 14, 31, 49	0
1	B	394/406 (97%)	0.27	28 (7%) 17 16	8, 13, 38, 55	0
All	All	798/812 (98%)	0.26	50 (6%) 21 20	8, 14, 36, 55	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	THR	9.3
1	B	219	LEU	7.1
1	A	217	THR	7.0
1	A	216	GLU	6.9
1	B	216	GLU	6.2
1	B	215	ASN	5.4
1	B	99	TYR	5.3
1	B	14	MET	4.8
1	A	67	LYS	4.7
1	B	13	LEU	4.6
1	A	63	VAL	4.5
1	A	282	LYS	4.4
1	B	15	GLU	4.3
1	A	219	LEU	4.2
1	B	282	LYS	4.2
1	A	64	GLU	3.8
1	A	405	LEU	3.7
1	B	64	GLU	3.7
1	A	71	HIS	3.6
1	B	63	VAL	3.6
1	B	71	HIS	3.5
1	B	67	LYS	3.5
1	B	243	ILE	3.5
1	B	68	LYS	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	89	PHE	3.2
1	B	12	PRO	3.2
1	B	265	ASP	3.2
1	B	405	LEU	3.2
1	A	286	MET	3.1
1	A	309	GLU	3.1
1	A	243	ILE	3.1
1	A	218	HIS	3.0
1	B	218	HIS	3.0
1	B	61	HIS	2.6
1	B	220	ASP	2.6
1	B	244	GLY	2.6
1	A	65	LEU	2.6
1	B	281	SER	2.5
1	A	70	ASP	2.5
1	A	265	ASP	2.4
1	A	215	ASN	2.4
1	A	354	VAL	2.4
1	A	227	LEU	2.2
1	B	11	LEU	2.2
1	A	1	GLY	2.1
1	A	68	LYS	2.1
1	A	5	SER	2.1
1	B	309	GLU	2.0
1	B	225	ILE	2.0
1	B	280	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	CSD	B	177	8/9	0.97	0.09	-	10,11,16,20	0
1	CSD	A	177	8/9	0.96	0.11	-	9,11,15,21	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.