



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2017 – 12:27 AM EDT

PDB ID : 2D73  
Title : Crystal Structure Analysis of SusB  
Authors : Kitamura, M.; Yao, M.  
Deposited on : unknown  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2696 (1.60-1.60)
Clashscore	112137	2967 (1.60-1.60)
Ramachandran outliers	110173	2887 (1.60-1.60)
Sidechain outliers	110143	2886 (1.60-1.60)
RSRZ outliers	101464	2714 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	738	<div> <div>5%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>
1	B	738	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div>..</div> </div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12992 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-glucosidase SusB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	717	Total	C	N	O	S	0	0	0
			5784	3671	977	1108	28			
1	B	717	Total	C	N	O	S	0	0	0
			5784	3671	977	1108	28			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	EXPRESSION TAG	GB 29341018
A	2	GLY	-	EXPRESSION TAG	GB 29341018
A	3	SER	-	EXPRESSION TAG	GB 29341018
A	4	SER	-	EXPRESSION TAG	GB 29341018
A	5	HIS	-	EXPRESSION TAG	GB 29341018
A	6	HIS	-	EXPRESSION TAG	GB 29341018
A	7	HIS	-	EXPRESSION TAG	GB 29341018
A	8	HIS	-	EXPRESSION TAG	GB 29341018
A	9	HIS	-	EXPRESSION TAG	GB 29341018
A	10	HIS	-	EXPRESSION TAG	GB 29341018
A	11	SER	-	EXPRESSION TAG	GB 29341018
A	12	SER	-	EXPRESSION TAG	GB 29341018
A	13	GLY	-	EXPRESSION TAG	GB 29341018
A	14	LEU	-	EXPRESSION TAG	GB 29341018
A	15	VAL	-	EXPRESSION TAG	GB 29341018
A	16	PRO	-	EXPRESSION TAG	GB 29341018
A	17	ARG	-	EXPRESSION TAG	GB 29341018
A	18	GLY	-	EXPRESSION TAG	GB 29341018
A	19	SER	-	EXPRESSION TAG	GB 29341018
A	20	HIS	-	EXPRESSION TAG	GB 29341018
A	21	MET	-	EXPRESSION TAG	GB 29341018
B	1	MET	-	EXPRESSION TAG	GB 29341018
B	2	GLY	-	EXPRESSION TAG	GB 29341018
B	3	SER	-	EXPRESSION TAG	GB 29341018
B	4	SER	-	EXPRESSION TAG	GB 29341018

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	5	HIS	-	EXPRESSION TAG	GB 29341018
B	6	HIS	-	EXPRESSION TAG	GB 29341018
B	7	HIS	-	EXPRESSION TAG	GB 29341018
B	8	HIS	-	EXPRESSION TAG	GB 29341018
B	9	HIS	-	EXPRESSION TAG	GB 29341018
B	10	HIS	-	EXPRESSION TAG	GB 29341018
B	11	SER	-	EXPRESSION TAG	GB 29341018
B	12	SER	-	EXPRESSION TAG	GB 29341018
B	13	GLY	-	EXPRESSION TAG	GB 29341018
B	14	LEU	-	EXPRESSION TAG	GB 29341018
B	15	VAL	-	EXPRESSION TAG	GB 29341018
B	16	PRO	-	EXPRESSION TAG	GB 29341018
B	17	ARG	-	EXPRESSION TAG	GB 29341018
B	18	GLY	-	EXPRESSION TAG	GB 29341018
B	19	SER	-	EXPRESSION TAG	GB 29341018
B	20	HIS	-	EXPRESSION TAG	GB 29341018
B	21	MET	-	EXPRESSION TAG	GB 29341018

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	699	Total O 699 699	0	0
3	B	723	Total O 723 723	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: alpha-glucosidase SusB



#### • Molecule 1: alpha-glucosidase SusB



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.63Å 112.34Å 102.57Å 90.00° 100.65° 90.00°	Depositor
Resolution (Å)	15.00 – 1.60 45.98 – 1.60	Depositor EDS
% Data completeness (in resolution range)	89.9 (15.00-1.60) 99.8 (45.98-1.60)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.70 (at 1.60Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.177 , 0.198 0.176 , 0.198	Depositor DCC
$R_{free}$ test set	21894 reflections (9.93%)	DCC
Wilson B-factor (Å <sup>2</sup> )	13.8	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 53.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12992	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/5935	0.64	5/8048 (0.1%)
1	B	0.32	1/5935 (0.0%)	0.64	5/8048 (0.1%)
All	All	0.31	1/11870 (0.0%)	0.64	10/16096 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	230	ALA	N-CA	7.43	1.61	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	229	THR	O-C-N	-9.26	107.88	122.70
1	B	229	THR	C-N-CA	7.47	140.38	121.70
1	B	600	ALA	N-CA-C	-6.17	94.33	111.00
1	A	600	ALA	N-CA-C	-6.12	94.46	111.00
1	B	229	THR	CA-C-O	6.02	132.75	120.10
1	B	526	GLU	N-CA-C	-5.66	95.72	111.00
1	A	526	GLU	N-CA-C	-5.59	95.91	111.00
1	B	230	ALA	N-CA-C	-5.47	96.23	111.00
1	A	230	ALA	N-CA-CB	-5.05	103.03	110.10
1	A	259	ASP	N-CA-C	-5.01	97.48	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	230	ALA	Mainchain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5784	0	5552	26	0
1	B	5784	0	5552	56	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	699	0	0	5	0
3	B	723	0	0	7	0
All	All	12992	0	11104	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:453:LYS:HA	1:B:453:LYS:HE2	1.43	0.98
1:B:73:THR:HG21	1:B:77:TRP:HA	1.55	0.88
1:A:730:SER:HA	1:A:733:LYS:HE3	1.72	0.71
1:B:111:GLN:HG2	1:B:119:GLU:HG2	1.74	0.70
1:B:534:GLU:HA	1:B:538:GLY:HA2	1.74	0.68
1:B:73:THR:HG22	1:B:74:ASP:O	1.94	0.68
1:A:534:GLU:HA	1:A:538:GLY:HA2	1.75	0.67
1:A:111:GLN:HG2	1:A:119:GLU:HG2	1.79	0.62
1:B:733:LYS:HE3	1:B:733:LYS:HA	1.82	0.61
1:B:73:THR:CG2	1:B:77:TRP:HA	2.28	0.61
1:B:22:GLN:HE21	1:B:24:LYS:HE2	1.68	0.59
1:B:453:LYS:HA	1:B:453:LYS:CE	2.27	0.58
1:A:57:LYS:HD2	1:A:178:ASN:HD22	1.70	0.56
1:B:206:ARG:HG3	1:B:224:GLN:HG2	1.89	0.55

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:THR:HG21	1:B:77:TRP:CA	2.31	0.55
1:B:604:PRO:O	1:B:608:GLU:HG3	2.05	0.55
1:B:667:PHE:HE2	1:B:709:LEU:HD13	1.72	0.55
1:B:565:GLU:HG2	1:B:609:ARG:NH1	2.20	0.55
1:A:402:GLY:CA	1:B:72:ARG:HH21	2.20	0.54
1:B:23:GLN:N	1:B:23:GLN:NE2	2.56	0.53
1:A:437:HIS:HD2	3:A:1041:HOH:O	1.92	0.52
1:B:181:HIS:HD2	3:B:1345:HOH:O	1.92	0.52
1:B:704:THR:HG23	1:B:737:ARG:HH12	1.74	0.52
1:B:733:LYS:HE3	1:B:733:LYS:CA	2.40	0.52
1:B:327:TYR:CZ	1:B:599:MET:HB2	2.45	0.52
1:B:73:THR:HG23	1:B:78:VAL:N	2.24	0.52
1:B:667:PHE:CE2	1:B:709:LEU:HD13	2.44	0.51
1:B:327:TYR:CE2	1:B:599:MET:HB2	2.45	0.51
1:B:708:LYS:C	1:B:709:LEU:HD12	2.31	0.51
1:A:581:SER:HB2	3:A:1454:HOH:O	2.11	0.50
1:B:581:SER:HB2	3:B:1363:HOH:O	2.12	0.50
1:A:261:LYS:O	1:A:261:LYS:HD3	2.12	0.50
1:B:344:THR:OG1	1:B:346:GLU:HG2	2.12	0.50
1:B:85:LYS:O	1:B:89:LYS:HG3	2.12	0.50
1:B:59:SER:HB3	1:B:174:GLN:HB2	1.93	0.49
1:A:261:LYS:C	1:A:261:LYS:HD3	2.33	0.49
1:B:38:VAL:CG1	1:B:42:GLY:HA2	2.43	0.49
1:A:91:ASN:HB3	1:A:135:MET:HE3	1.96	0.48
1:B:181:HIS:HE1	3:B:1154:HOH:O	1.98	0.47
1:A:59:SER:HB3	1:A:174:GLN:HB2	1.96	0.47
1:A:244:HIS:CG	1:A:245:GLU:H	2.33	0.46
1:A:161:LYS:HD2	3:A:2003:HOH:O	2.16	0.46
1:B:76:ASP:O	1:B:77:TRP:HB2	2.16	0.46
1:B:244:HIS:CG	1:B:245:GLU:H	2.34	0.45
1:A:169:LYS:O	1:A:170:GLU:HG2	2.17	0.45
1:B:667:PHE:O	1:B:705:ASN:HA	2.16	0.45
1:B:605:GLU:HG3	3:B:1595:HOH:O	2.17	0.45
1:A:269:LEU:HB2	1:A:277:LYS:HD3	1.99	0.44
1:B:69:ASP:OD1	1:B:72:ARG:HD3	2.17	0.44
1:B:74:ASP:OD1	1:B:75:PHE:N	2.48	0.44
1:B:69:ASP:OD1	1:B:72:ARG:HB2	2.18	0.44
1:B:437:HIS:HD2	3:B:1053:HOH:O	2.00	0.44
1:A:73:THR:OG1	1:A:78:VAL:N	2.51	0.43
1:B:565:GLU:OE1	1:B:570:LYS:HD2	2.18	0.43
1:B:683:ASP:HB3	1:B:687:ALA:HB3	1.99	0.43

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:73:THR:CG2	1:B:77:TRP:CA	2.95	0.43
1:A:298:ARG:HD2	1:A:629:TYR:O	2.19	0.43
1:B:26:THR:CG2	1:B:30:ASN:HA	2.49	0.43
1:B:444:VAL:HG21	1:B:486:MET:SD	2.59	0.43
1:B:57:LYS:HD2	1:B:178:ASN:HD22	1.83	0.43
1:A:327:TYR:CZ	1:A:599:MET:HB2	2.54	0.42
1:B:85:LYS:HE3	1:B:135:MET:O	2.19	0.42
1:B:565:GLU:CD	1:B:570:LYS:HD2	2.40	0.42
1:B:730:SER:HA	1:B:733:LYS:HD2	2.01	0.42
1:A:111:GLN:CG	1:A:119:GLU:HG2	2.49	0.42
1:A:135:MET:HG2	3:A:1378:HOH:O	2.19	0.42
1:B:326:LYS:HB2	1:B:622:LEU:HD11	2.02	0.42
1:A:328:ILE:HG22	1:A:600:ALA:HB3	2.01	0.42
1:A:26:THR:CG2	1:A:30:ASN:HA	2.50	0.41
1:A:706:LYS:NZ	3:A:1295:HOH:O	2.53	0.41
1:B:38:VAL:HG12	1:B:42:GLY:HA2	2.00	0.41
1:B:328:ILE:HG22	1:B:600:ALA:HB3	2.03	0.41
1:B:424:HIS:HD2	3:B:1247:HOH:O	2.03	0.41
1:A:71:THR:HG23	1:A:71:THR:O	2.21	0.41
1:B:571:MET:HG2	3:B:1511:HOH:O	2.21	0.41
1:B:569:ASN:HA	1:B:572:ASN:O	2.21	0.41
1:A:141:VAL:HB	1:A:156:GLU:HB2	2.03	0.41
1:A:654:CYS:O	1:A:718:GLY:HA3	2.21	0.40
1:B:229:THR:HA	1:B:230:ALA:C	2.42	0.40
1:B:32:LEU:N	1:B:32:LEU:HD12	2.36	0.40
1:B:269:LEU:HB2	1:B:277:LYS:HD3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	715/738 (97%)	688 (96%)	27 (4%)	0	100	100
1	B	715/738 (97%)	691 (97%)	24 (3%)	0	100	100
All	All	1430/1476 (97%)	1379 (96%)	51 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	618/636 (97%)	612 (99%)	6 (1%)	80	65
1	B	618/636 (97%)	608 (98%)	10 (2%)	68	45
All	All	1236/1272 (97%)	1220 (99%)	16 (1%)	73	55

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	70	ASN
1	A	196	ASP
1	A	229	THR
1	A	261	LYS
1	A	511	ARG
1	A	542	TYR
1	B	23	GLN
1	B	87	ASP
1	B	122	ASN
1	B	196	ASP
1	B	229	THR
1	B	424	HIS
1	B	511	ARG
1	B	542	TYR
1	B	565	GLU
1	B	733	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	37	GLN
1	A	70	ASN
1	A	178	ASN
1	A	299	ASN
1	A	437	HIS
1	A	473	ASN
1	A	675	GLN
1	A	715	ASN
1	B	22	GLN
1	B	23	GLN
1	B	31	ASN
1	B	37	GLN
1	B	122	ASN
1	B	178	ASN
1	B	181	HIS
1	B	255	HIS
1	B	285	ASN
1	B	299	ASN
1	B	424	HIS
1	B	437	HIS
1	B	473	ASN
1	B	578	GLN
1	B	628	ASN
1	B	715	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	717/738 (97%)	0.09	34 (4%) 32 31	7, 16, 32, 58	0
1	B	717/738 (97%)	0.12	31 (4%) 36 34	6, 15, 31, 68	0
All	All	1434/1476 (97%)	0.11	65 (4%) 34 32	6, 15, 31, 68	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	83	LEU	11.7
1	A	83	LEU	9.8
1	B	80	ARG	8.6
1	A	728	ASP	7.4
1	B	71	THR	7.4
1	B	70	ASN	7.1
1	B	84	THR	7.1
1	B	78	VAL	7.1
1	B	77	TRP	7.0
1	A	70	ASN	7.0
1	A	80	ARG	6.8
1	B	72	ARG	6.7
1	B	82	ASP	6.3
1	A	78	VAL	6.2
1	A	84	THR	6.0
1	B	69	ASP	5.9
1	B	73	THR	5.9
1	A	77	TRP	5.7
1	B	79	ASP	5.5
1	B	727	LYS	5.5
1	B	74	ASP	5.5
1	B	85	LYS	5.4
1	A	71	THR	5.1
1	A	729	LYS	5.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	81	ARG	4.9
1	A	733	LYS	4.6
1	A	73	THR	4.5
1	B	86	LEU	4.5
1	A	72	ARG	4.2
1	A	85	LYS	4.2
1	A	732	ALA	4.2
1	A	79	ASP	4.1
1	A	86	LEU	4.0
1	A	727	LYS	4.0
1	B	733	LYS	3.9
1	A	82	ASP	3.7
1	B	730	SER	3.6
1	B	75	PHE	3.3
1	A	730	SER	3.3
1	A	74	ASP	3.3
1	A	161	LYS	3.3
1	B	728	ASP	3.0
1	A	69	ASP	3.0
1	A	76	ASP	2.7
1	A	726	VAL	2.7
1	B	658	GLU	2.6
1	B	81	ARG	2.6
1	A	348	THR	2.5
1	B	576	ASN	2.4
1	B	731	GLU	2.4
1	A	75	PHE	2.4
1	B	161	LYS	2.4
1	A	709	LEU	2.3
1	B	685	LYS	2.3
1	A	648	ASP	2.3
1	B	87	ASP	2.3
1	A	316	ALA	2.3
1	A	87	ASP	2.3
1	B	673	GLY	2.3
1	B	648	ASP	2.2
1	B	76	ASP	2.2
1	A	576	ASN	2.1
1	B	709	LEU	2.1
1	A	261	LYS	2.0
1	A	738	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	B	802	1/1	1.00	0.10	-0.41	9,9,9,9	1
2	CA	A	801	1/1	1.00	0.08	-1.76	10,10,10,10	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.