



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:12 pm GMT

PDB ID : 3D8X
Title : Crystal Structure of Saccharomyces cerevisiae NDPPH Dependent Thioredoxin Reductase 1
Authors : Zhang, Z.Y.; Bao, R.; Yu, J.; Chen, Y.X.; Zhou, C.-Z.
Deposited on : 2008-05-26
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	2583 (2.80-2.80)
Clashscore	112137	3033 (2.80-2.80)
Ramachandran outliers	110173	2983 (2.80-2.80)
Sidechain outliers	110143	2985 (2.80-2.80)
RSRZ outliers	101464	2610 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div>2%</div> <div> <div></div> <div>77%</div> <div>16%</div> <div>5%</div> </div> </div>
1	B	326	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>15%</div> <div>.</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5016 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioredoxin reductase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2393	1509	402	470	12			
1	B	318	Total	C	N	O	S	5	0	0
			2393	1509	402	470	12			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	EXPRESSION TAG	UNP P29509
A	-5	GLY	-	EXPRESSION TAG	UNP P29509
A	-4	HIS	-	EXPRESSION TAG	UNP P29509
A	-3	HIS	-	EXPRESSION TAG	UNP P29509
A	-2	HIS	-	EXPRESSION TAG	UNP P29509
A	-1	HIS	-	EXPRESSION TAG	UNP P29509
A	0	HIS	-	EXPRESSION TAG	UNP P29509
A	1	HIS	-	EXPRESSION TAG	UNP P29509
B	-6	MET	-	EXPRESSION TAG	UNP P29509
B	-5	GLY	-	EXPRESSION TAG	UNP P29509
B	-4	HIS	-	EXPRESSION TAG	UNP P29509
B	-3	HIS	-	EXPRESSION TAG	UNP P29509
B	-2	HIS	-	EXPRESSION TAG	UNP P29509
B	-1	HIS	-	EXPRESSION TAG	UNP P29509
B	0	HIS	-	EXPRESSION TAG	UNP P29509
B	1	HIS	-	EXPRESSION TAG	UNP P29509

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			24	6	5	10	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	47	Total 47	O 47	0	0
4	B	53	Total 53	O 53	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thioredoxin reductase 1



• Molecule 1: Thioredoxin reductase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	54.23Å 125.00Å 60.97Å 90.00° 114.16° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80 19.80 – 2.80	Depositor EDS
% Data completeness (in resolution range)	96.6 (20.00-2.80) 96.6 (19.80-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.15	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 2.79Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.226 , 0.234 0.226 , 0.237	Depositor DCC
R_{free} test set	911 reflections (5.45%)	DCC
Wilson B-factor (Å ²)	41.6	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5016	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/2434	0.59	0/3292
1	B	0.47	1/2434 (0.0%)	0.60	0/3292
All	All	0.47	1/4868 (0.0%)	0.60	0/6584

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	227	ARG	CG-CD	5.05	1.64	1.51

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2393	0	2402	53	0
1	B	2393	0	2402	50	0
2	A	53	0	31	1	0
2	B	53	0	31	0	0
3	A	24	0	4	2	0
4	A	47	0	0	4	0
4	B	53	0	0	2	0
All	All	5016	0	4870	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (105) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75:SER:O	1:B:80:THR:HG23	1.35	1.24
1:A:278:THR:HG22	1:A:280:VAL:H	1.15	1.10
1:B:50:THR:CG2	1:B:51:GLU:H	1.65	1.09
1:B:50:THR:HG23	1:B:51:GLU:H	1.12	1.06
1:A:50:THR:HG23	1:A:51:GLU:H	1.19	1.03
1:B:278:THR:HG22	1:B:280:VAL:H	1.20	1.03
1:B:50:THR:HG23	1:B:51:GLU:N	1.76	0.99
1:A:50:THR:HG23	1:A:51:GLU:N	1.78	0.98
1:A:50:THR:CG2	1:A:51:GLU:H	1.79	0.96
1:B:75:SER:O	1:B:80:THR:CG2	2.16	0.93
1:B:92:LEU:H	1:B:260:GLN:HE22	1.13	0.93
1:A:125:ARG:NH1	1:A:131:GLU:OE2	2.06	0.87
1:A:23:ALA:HB2	1:A:80:THR:HG22	1.59	0.85
1:A:75:SER:O	1:A:80:THR:HG23	1.78	0.83
1:B:50:THR:CG2	1:B:51:GLU:N	2.35	0.82
1:A:72:ARG:HH11	1:A:82:ILE:HD12	1.47	0.80
1:A:155:LYS:HB3	1:A:242:SER:OG	1.83	0.79
1:A:7:THR:CG2	1:A:112:THR:HG21	2.15	0.77
1:A:11:SER:OG	1:A:43:GLY:N	2.19	0.75
1:A:19:ALA:HB1	1:A:80:THR:HG21	1.70	0.74
1:B:7:THR:CG2	1:B:112:THR:HG21	2.19	0.73
1:A:72:ARG:NH1	1:A:82:ILE:HD12	2.03	0.72
1:B:11:SER:OG	1:B:43:GLY:N	2.23	0.71
1:B:155:LYS:HB3	1:B:242:SER:OG	1.90	0.71
1:A:50:THR:CG2	1:A:51:GLU:N	2.43	0.70
1:B:92:LEU:H	1:B:260:GLN:NE2	1.88	0.69
1:A:189:LEU:HD23	1:A:196:GLN:HB2	1.75	0.69
1:B:46:LEU:O	1:B:49:THR:HG22	1.93	0.68
1:B:160:ILE:HD12	1:B:244:LEU:HD11	1.74	0.68
1:A:46:LEU:O	1:A:49:THR:HG22	1.94	0.68
1:B:171:GLN:O	1:B:174:THR:HB	1.94	0.68
1:A:44:GLY:O	1:A:47:THR:HB	1.94	0.67
1:B:75:SER:C	1:B:80:THR:HG23	2.12	0.67
1:B:44:GLY:O	1:B:47:THR:HB	1.95	0.66
1:A:7:THR:HG22	1:A:112:THR:HG21	1.77	0.65
1:A:171:GLN:O	1:A:174:THR:HB	1.96	0.65
1:A:160:ILE:HD12	1:A:244:LEU:HD11	1.79	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:278:THR:HG22	1:B:280:VAL:N	2.04	0.61
1:B:7:THR:HG22	1:B:112:THR:HG21	1.81	0.61
1:A:232:LYS:HD2	4:A:1020:HOH:O	2.01	0.60
1:A:278:THR:HG22	1:A:280:VAL:N	2.01	0.60
1:A:278:THR:CG2	1:A:279:SER:N	2.64	0.60
1:B:125:ARG:NH1	1:B:131:GLU:OE2	2.34	0.60
1:B:278:THR:CG2	1:B:279:SER:N	2.65	0.60
1:B:278:THR:HB	1:B:283:PHE:O	2.03	0.58
1:A:278:THR:HB	1:A:283:PHE:O	2.04	0.57
1:B:278:THR:CG2	1:B:280:VAL:H	2.07	0.57
1:A:7:THR:HG21	1:A:112:THR:HG21	1.85	0.56
1:B:29:PRO:HG2	1:B:80:THR:HB	1.87	0.56
1:B:129:PRO:HB2	1:B:215:GLU:HG3	1.86	0.56
1:B:171:GLN:NE2	1:B:202:ASN:HD22	2.03	0.55
1:A:271:THR:HG21	1:A:290:GLN:NE2	2.21	0.55
1:B:23:ALA:HB2	1:B:80:THR:HG22	1.88	0.54
1:B:99:LEU:HD22	1:B:110:VAL:HB	1.89	0.54
1:B:50:THR:HG22	1:B:51:GLU:H	1.65	0.53
1:B:261:VAL:CG2	1:B:278:THR:HG23	2.39	0.52
1:B:261:VAL:HG23	1:B:278:THR:HG23	1.92	0.52
1:A:261:VAL:CG2	1:A:278:THR:HG23	2.40	0.51
1:B:278:THR:HG23	1:B:279:SER:N	2.25	0.51
1:A:129:PRO:HB2	1:A:215:GLU:HG3	1.93	0.51
3:A:481:NDP:H8A	4:A:1029:HOH:O	2.09	0.51
1:B:271:THR:HG21	1:B:290:GLN:NE2	2.24	0.50
1:A:23:ALA:HB2	1:A:80:THR:CG2	2.37	0.49
1:B:160:ILE:CD1	1:B:244:LEU:HD11	2.42	0.49
1:A:174:THR:HG21	1:A:202:ASN:HD21	1.78	0.49
1:A:278:THR:HG23	1:A:279:SER:N	2.28	0.49
1:A:155:LYS:CB	1:A:242:SER:OG	2.59	0.47
1:A:160:ILE:CD1	1:A:244:LEU:HD11	2.44	0.47
1:B:126:MET:CE	1:B:160:ILE:HD13	2.45	0.47
1:A:194:ILE:H	1:A:194:ILE:HG13	1.52	0.47
3:A:481:NDP:H8A	3:A:481:NDP:O2B	2.15	0.46
1:A:261:VAL:HG23	1:A:278:THR:HG23	1.98	0.46
1:B:126:MET:HE3	1:B:128:LEU:HD21	1.98	0.46
1:B:189:LEU:HD12	1:B:196:GLN:HB2	1.97	0.46
1:A:126:MET:CE	1:A:160:ILE:HD13	2.46	0.46
1:B:7:THR:HG21	1:B:112:THR:HG21	1.95	0.46
1:B:278:THR:HG21	1:B:283:PHE:HD1	1.82	0.45
1:A:117:LEU:HD12	1:A:269:ILE:HD12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:ARG:HA	1:A:153:ARG:HD3	1.64	0.44
1:B:72:ARG:HH11	1:B:82:ILE:HD12	1.83	0.43
1:A:145:CYS:SG	2:A:1001:FAD:C4X	3.07	0.43
1:B:126:MET:HE2	1:B:160:ILE:HD13	2.00	0.43
1:A:277:LEU:N	1:A:277:LEU:CD2	2.82	0.43
1:A:277:LEU:N	1:A:277:LEU:HD23	2.34	0.42
1:B:171:GLN:HE22	1:B:202:ASN:HD22	1.67	0.42
1:B:50:THR:O	1:B:63:THR:HA	2.20	0.42
1:A:230:ASN:ND2	1:A:233:LYS:HD2	2.34	0.42
1:B:153:ARG:HD3	1:B:153:ARG:HA	1.65	0.42
1:B:295:ARG:HB2	1:B:295:ARG:HE	1.74	0.42
1:A:171:GLN:NE2	1:A:202:ASN:HD22	2.17	0.41
1:B:194:ILE:H	1:B:194:ILE:HG13	1.43	0.41
1:A:4:ASN:O	1:A:112:THR:HA	2.20	0.41
1:A:217:LYS:HE3	4:A:1017:HOH:O	2.19	0.41
1:B:46:LEU:HB2	4:B:1050:HOH:O	2.20	0.41
1:A:164:ASP:OD1	1:A:164:ASP:N	2.34	0.41
1:A:261:VAL:HG23	1:A:279:SER:HB3	2.02	0.41
1:A:126:MET:HE2	1:A:160:ILE:HG21	2.02	0.41
1:A:271:THR:CB	1:A:290:GLN:HE21	2.34	0.41
1:A:108:GLU:HA	1:A:109:PRO:HD2	1.82	0.41
1:A:261:VAL:CG2	1:A:278:THR:CG2	2.99	0.41
1:A:61:GLY:O	1:A:62:LEU:HB3	2.21	0.41
1:A:270:LYS:HE2	4:A:1030:HOH:O	2.21	0.40
1:B:72:ARG:NH2	4:B:1089:HOH:O	2.50	0.40
1:B:16:HIS:CE1	1:B:72:ARG:HA	2.56	0.40
1:B:95:LYS:HD2	1:B:95:LYS:HA	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/326 (97%)	301 (95%)	15 (5%)	0	100	100
1	B	316/326 (97%)	303 (96%)	13 (4%)	0	100	100
All	All	632/652 (97%)	604 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/260 (97%)	226 (89%)	27 (11%)	8	22
1	B	253/260 (97%)	227 (90%)	26 (10%)	8	24
All	All	506/520 (97%)	453 (90%)	53 (10%)	8	23

All (53) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	46	LEU
1	A	47	THR
1	A	72	ARG
1	A	84	THR
1	A	90	VAL
1	A	108	GLU
1	A	117	LEU
1	A	125	ARG
1	A	128	LEU
1	A	153	ARG
1	A	164	ASP
1	A	183	LEU
1	A	189	LEU
1	A	194	ILE
1	A	196	GLN
1	A	207	ILE
1	A	208	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	214	LEU
1	A	222	LEU
1	A	241	VAL
1	A	265	GLU
1	A	271	THR
1	A	277	LEU
1	A	278	THR
1	A	290	GLN
1	A	292	SER
1	B	45	GLN
1	B	46	LEU
1	B	47	THR
1	B	50	THR
1	B	77	LYS
1	B	80	THR
1	B	84	THR
1	B	90	VAL
1	B	99	LEU
1	B	104	ASN
1	B	117	LEU
1	B	125	ARG
1	B	128	LEU
1	B	164	ASP
1	B	174	THR
1	B	183	LEU
1	B	194	ILE
1	B	196	GLN
1	B	207	ILE
1	B	208	LEU
1	B	241	VAL
1	B	265	GLU
1	B	271	THR
1	B	277	LEU
1	B	278	THR
1	B	290	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	171	GLN
1	A	290	GLN
1	B	171	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	260	GLN
1	B	290	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	FAD	A	1001	-	51,58,58	1.37	7 (13%)	54,89,89	2.01	8 (14%)
3	NDP	A	481	-	18,23,52	0.96	0	18,33,80	2.43	3 (16%)
2	FAD	B	1001	-	51,58,58	1.39	7 (13%)	54,89,89	1.87	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1001	-	-	0/28/50/50	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	A	481	-	-	0/2/5/77	0/2/2/5
2	FAD	B	1001	-	-	0/28/50/50	0/6/6/6

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	FAD	C1'-N10	2.21	1.50	1.48
2	A	1001	FAD	C5X-N5	2.23	1.38	1.35
2	A	1001	FAD	C2A-N1A	2.44	1.38	1.33
2	B	1001	FAD	C5X-N5	2.50	1.39	1.35
2	B	1001	FAD	C2A-N1A	2.66	1.38	1.33
2	B	1001	FAD	C1'-N10	2.78	1.51	1.48
2	A	1001	FAD	C4-N3	3.06	1.38	1.33
2	A	1001	FAD	C10-N1	3.44	1.38	1.33
2	B	1001	FAD	C10-N1	3.53	1.38	1.33
2	B	1001	FAD	C4X-N5	3.55	1.38	1.33
2	A	1001	FAD	C2A-N3A	3.76	1.38	1.32
2	B	1001	FAD	C2A-N3A	3.77	1.38	1.32
2	A	1001	FAD	C4X-N5	4.06	1.39	1.33
2	B	1001	FAD	C4-N3	4.16	1.40	1.33

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	FAD	N3A-C2A-N1A	-9.59	120.51	128.86
2	B	1001	FAD	N3A-C2A-N1A	-9.16	120.88	128.86
3	A	481	NDP	N3A-C2A-N1A	-9.08	120.95	128.86
2	A	1001	FAD	C4X-C4-N3	-2.98	119.24	123.48
2	B	1001	FAD	C4X-C4-N3	-2.86	119.41	123.48
2	B	1001	FAD	C8M-C8-C9	-2.64	113.73	120.34
2	A	1001	FAD	O5'-C5'-C4'	-2.42	102.89	109.36
2	B	1001	FAD	C8M-C8-C7	-2.25	116.00	120.72
2	A	1001	FAD	C8M-C8-C9	-2.23	114.74	120.34
3	A	481	NDP	C4A-C5A-N7A	-2.21	107.27	109.41
2	B	1001	FAD	C4A-C5A-N7A	-2.19	107.29	109.41
2	B	1001	FAD	C5X-C9A-N10	2.25	119.33	117.66
3	A	481	NDP	O2N-PN-O1N	2.29	119.45	110.50
2	A	1001	FAD	C5X-C9A-N10	2.80	119.73	117.66
2	A	1001	FAD	C4X-N5-C5X	2.82	119.74	116.76
2	A	1001	FAD	C1'-N10-C9A	3.49	121.54	118.35
2	B	1001	FAD	C4X-N5-C5X	3.80	120.78	116.76
2	B	1001	FAD	C4-N3-C2	5.37	119.86	115.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	FAD	C4-N3-C2	6.80	121.10	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1001	FAD	1	0
3	A	481	NDP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/326 (97%)	0.01	6 (1%) 67 58	7, 22, 30, 33	4 (1%)
1	B	318/326 (97%)	-0.14	4 (1%) 77 71	7, 23, 30, 34	9 (2%)
All	All	636/652 (97%)	-0.07	10 (1%) 72 65	7, 23, 30, 34	13 (2%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	VAL	3.3
1	A	7	THR	2.9
1	A	115	ILE	2.8
1	B	234	ASN	2.7
1	A	8	ILE	2.4
1	A	117	LEU	2.2
1	A	154	ASN	2.1
1	B	221	LYS	2.1
1	B	197	LYS	2.1
1	B	2	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	NDP	A	481	24/48	0.88	0.22	0.32	5,15,18,18	24
2	FAD	B	1001	53/53	0.96	0.13	-0.80	15,18,21,22	0
2	FAD	A	1001	53/53	0.96	0.13	-1.09	6,16,19,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.