



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 13, 2017 – 11:12 pm GMT

PDB ID : 1DCT
Title : DNA (CYTOSINE-5) METHYLASE FROM HAEIII COVALENTLY BOUND TO DNA
Authors : Reinisch, K.M.; Chen, L.; Verdine, G.L.; Lipscomb, W.N.
Deposited on : 1995-05-17
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric | Whole archive (#Entries) | Similar resolution (#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|---|
| Clashscore | 112137 | 3033 (2.80-2.80) |
| Ramachandran outliers | 110173 | 2983 (2.80-2.80) |
| Sidechain outliers | 110143 | 2985 (2.80-2.80) |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1 | F | 18 | |
| 1 | G | 18 | |
| 2 | M | 18 | |
| 2 | N | 18 | |
| 3 | A | 324 | |
| 3 | B | 324 | |

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

| Mol | Type | Chain | Res | Chirality | Geometry | Clashes | Electron density |
|-----|------|-------|-----|-----------|----------|---------|------------------|
| 1 | C49 | F | 10 | X | - | - | - |
| 1 | C49 | G | 10 | X | - | - | - |

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6908 atoms, of which 268 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*AP*CP*CP*AP*GP*CP*AP*GP*GP*(C49)P*CP*AP*CP*CP*AP*GP*TP*G)-3').

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace | |
|-----|-------|----------|--------------|----------|--------|---------|----------|---------|---------|-------|---|
| | | | Total | C | F | N | O | | | | P |
| 1 | F | 18 | Total 367 | C 174 | F 1 | N 73 | O 102 | P 17 | 0 | 0 | 0 |
| 1 | G | 16 | Total 327 | C 155 | F 1 | N 65 | O 91 | P 15 | 0 | 0 | 0 |

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*CP*AP*CP*TP*GP*GP*TP*GP*GP*(C5M)P*CP*TP*GP*CP*TP*GP*G)-3').


| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace |
|-----|-------|----------|--------------|----------|---------|----------|---------|---------|---------|-------|
| | | | Total | C | N | O | P | | | |
| 2 | M | 18 | Total 368 | C 176 | N 65 | O 110 | P 17 | 0 | 0 | 0 |
| 2 | N | 17 | Total 346 | C 166 | N 60 | O 104 | P 16 | 0 | 0 | 0 |

- Molecule 3 is a protein called PROTEIN (MODIFICATION METHYLASE HAEIII).

| Mol | Chain | Residues | Atoms | | | | | ZeroOcc | AltConf | Trace | |
|-----|-------|----------|---------------|-----------|----------|----------|----------|---------|---------|-------|---|
| | | | Total | C | H | N | O | | | | S |
| 3 | A | 324 | Total 2749 | C 1680 | H 134 | N 455 | O 468 | S 12 | 0 | 0 | 0 |
| 3 | B | 324 | Total 2749 | C 1680 | H 134 | N 455 | O 468 | S 12 | 0 | 0 | 0 |


- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

| Mol | Chain | Residues | Atoms | | ZeroOcc | AltConf |
|-----|-------|----------|------------|---------|---------|---------|
| 4 | B | 1 | Total 1 | Ca 1 | 0 | 0 |
| 4 | A | 1 | Total 1 | Ca 1 | 0 | 0 |



• Molecule 3: PROTEIN (MODIFICATION METHYLASE HAEIII)

Chain B: 48% 44% 7%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

| Property | Value | Source |
|--|--|-----------|
| Space group | P 21 21 21 | Depositor |
| Cell constants a, b, c, α , β , γ | 57.57Å 108.04Å 155.79Å 90.00° 90.00° 90.00° | Depositor |
| Resolution (Å) | 10.00 – 2.80 | Depositor |
| % Data completeness (in resolution range) | (Not available) (10.00-2.80) | Depositor |
| R_{merge} | 0.08 | Depositor |
| R_{sym} | (Not available) | Depositor |
| Refinement program | X-PLOR | Depositor |
| R, R_{free} | 0.226 , 0.326 | Depositor |
| Estimated twinning fraction | No twinning to report. | Xtrriage |
| Total number of atoms | 6908 | wwPDB-VP |
| Average B, all atoms (Å ²) | 24.0 | wwPDB-VP |

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: CA, C49, 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths | | Bond angles | |
|-----|-------|--------------|----------------|-------------|-----------------|
| | | RMSZ | # $ Z > 5$ | RMSZ | # $ Z > 5$ |
| 1 | F | 1.54 | 2/388 (0.5%) | 2.60 | 46/594 (7.7%) |
| 1 | G | 1.57 | 5/343 (1.5%) | 2.38 | 30/525 (5.7%) |
| 2 | M | 1.73 | 5/388 (1.3%) | 2.43 | 42/596 (7.0%) |
| 2 | N | 1.75 | 6/363 (1.7%) | 2.44 | 38/557 (6.8%) |
| 3 | A | 0.52 | 0/2681 | 0.74 | 2/3615 (0.1%) |
| 3 | B | 0.51 | 0/2681 | 0.76 | 3/3615 (0.1%) |
| All | All | 0.89 | 18/6844 (0.3%) | 1.37 | 161/9502 (1.7%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1 | F | 1 | 2 |
| 1 | G | 1 | 0 |
| 2 | M | 0 | 1 |
| 3 | A | 0 | 1 |
| 3 | B | 0 | 1 |
| All | All | 2 | 5 |

The worst 5 of 18 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 2 | N | 8 | DT | C3'-O3' | -8.62 | 1.32 | 1.44 |
| 2 | M | 8 | DT | C3'-O3' | -8.24 | 1.33 | 1.44 |
| 1 | G | 14 | DC | C4'-O4' | -6.53 | 1.38 | 1.45 |
| 1 | F | 9 | DG | C3'-O3' | -6.34 | 1.35 | 1.44 |
| 2 | N | 10 | DG | C4'-O4' | -6.30 | 1.38 | 1.45 |

The worst 5 of 161 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 1 | F | 2 | DC | O4'-C4'-C3' | 10.12 | 112.07 | 106.00 |
| 1 | F | 13 | DC | C1'-O4'-C4' | -9.23 | 100.87 | 110.10 |
| 1 | F | 13 | DC | O4'-C4'-C3' | 9.15 | 111.49 | 106.00 |
| 1 | F | 4 | DA | N1-C2-N3 | -8.98 | 124.81 | 129.30 |
| 1 | G | 4 | DA | N1-C2-N3 | -8.92 | 124.84 | 129.30 |

All (2) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 1 | F | 10 | C49 | C1' |
| 1 | G | 10 | C49 | C1' |

All (5) planarity outliers are listed below:

| Mol | Chain | Res | Type | Group |
|-----|-------|-----|------|-----------|
| 3 | A | 147 | TYR | Sidechain |
| 3 | B | 38 | TYR | Sidechain |
| 1 | F | 1 | DA | Sidechain |
| 1 | F | 17 | DT | Sidechain |
| 2 | M | 17 | DG | Sidechain |

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1 | F | 367 | 0 | 200 | 18 | 0 |
| 1 | G | 327 | 0 | 178 | 10 | 0 |
| 2 | M | 368 | 0 | 207 | 14 | 0 |
| 2 | N | 346 | 0 | 196 | 5 | 0 |
| 3 | A | 2615 | 134 | 2606 | 153 | 0 |
| 3 | B | 2615 | 134 | 2606 | 160 | 0 |
| 4 | A | 1 | 0 | 0 | 0 | 0 |
| 4 | B | 1 | 0 | 0 | 0 | 0 |
| All | All | 6640 | 268 | 5993 | 343 | 0 |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 343 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

| Atom-1 | Atom-2 | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 1:F:4:DA:H2'' | 1:F:5:DG:C8 | 2.04 | 0.93 |
| 3:A:202:ASN:HB3 | 3:A:205:LYS:HB3 | 1.51 | 0.92 |
| 3:B:240:ALA:HB1 | 3:B:302:LYS:HA | 1.52 | 0.90 |
| 3:B:212:GLU:HG3 | 3:B:297:LEU:HD21 | 1.55 | 0.87 |
| 3:B:153:ARG:HG3 | 3:B:237:THR:HG23 | 1.60 | 0.83 |

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Favoured | Allowed | Outliers | Percentiles | |
|-----|-------|---------------|-----------|----------|----------|-------------|----|
| 3 | A | 322/324 (99%) | 273 (85%) | 39 (12%) | 10 (3%) | 5 | 16 |
| 3 | B | 322/324 (99%) | 283 (88%) | 26 (8%) | 13 (4%) | 3 | 11 |
| All | All | 644/648 (99%) | 556 (86%) | 65 (10%) | 23 (4%) | 4 | 13 |

5 of 23 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | A | 9 | GLY |
| 3 | A | 80 | LEU |
| 3 | A | 241 | SER |
| 3 | B | 9 | GLY |
| 3 | B | 186 | TRP |

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed | Rotameric | Outliers | Percentiles | |
|-----|-------|----------------|-----------|----------|-------------|----|
| 3 | A | 281/281 (100%) | 262 (93%) | 19 (7%) | 18 | 47 |
| 3 | B | 281/281 (100%) | 260 (92%) | 21 (8%) | 16 | 41 |
| All | All | 562/562 (100%) | 522 (93%) | 40 (7%) | 17 | 44 |

5 of 40 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | A | 273 | ARG |
| 3 | B | 30 | TYR |
| 3 | B | 274 | ARG |
| 3 | B | 8 | SER |
| 3 | B | 56 | SER |

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 3 | A | 262 | ASN |
| 3 | A | 284 | GLN |
| 3 | B | 211 | HIS |
| 3 | A | 244 | GLN |
| 3 | B | 210 | ASN |

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Type | Chain | Res | Link | Bond lengths | | | Bond angles | | |
|-----|------|-------|-----|------|--------------|------|----------|-------------|------|----------|
| | | | | | Counts | RMSZ | # Z > 2 | Counts | RMSZ | # Z > 2 |
| 1 | C49 | F | 10 | 1,3 | 15,22,24 | 1.24 | 3 (20%) | 17,33,38 | 4.49 | 7 (41%) |
| 1 | C49 | G | 10 | 1,3 | 15,22,24 | 1.27 | 2 (13%) | 17,33,38 | 4.22 | 6 (35%) |
| 2 | 5CM | M | 11 | 1,2 | 14,21,22 | 0.85 | 1 (7%) | 18,30,33 | 0.68 | 0 |
| 2 | 5CM | N | 11 | 1,2 | 14,21,22 | 0.86 | 1 (7%) | 18,30,33 | 0.66 | 0 |

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

| Mol | Type | Chain | Res | Link | Chirals | Torsions | Rings |
|-----|------|-------|-----|------|----------|-----------|---------|
| 1 | C49 | F | 10 | 1,3 | 1/1/8/10 | 0/7/40/46 | 0/2/2/2 |
| 1 | C49 | G | 10 | 1,3 | 1/1/8/10 | 0/7/40/46 | 0/2/2/2 |
| 2 | 5CM | M | 11 | 1,2 | - | 0/3/21/22 | 0/2/2/2 |
| 2 | 5CM | N | 11 | 1,2 | - | 0/3/21/22 | 0/2/2/2 |

The worst 5 of 7 bond length outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(Å) | Ideal(Å) |
|-----|-------|-----|------|---------|-------|-------------|----------|
| 2 | N | 11 | 5CM | C6-C5 | -2.50 | 1.33 | 1.40 |
| 2 | M | 11 | 5CM | C6-C5 | -2.49 | 1.33 | 1.40 |
| 1 | F | 10 | C49 | O3'-C3' | -2.12 | 1.38 | 1.43 |
| 1 | F | 10 | C49 | O4'-C1' | 2.22 | 1.47 | 1.42 |
| 1 | F | 10 | C49 | C4-N4 | 2.24 | 1.32 | 1.27 |

The worst 5 of 13 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms | Z | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-------------|-------|-------------|----------|
| 1 | G | 10 | C49 | C4'-O4'-C1' | -4.98 | 97.21 | 109.42 |
| 1 | F | 10 | C49 | C4'-O4'-C1' | -4.93 | 97.34 | 109.42 |
| 1 | F | 10 | C49 | F-C5-C6 | -3.10 | 104.01 | 107.02 |
| 1 | F | 10 | C49 | C2'-C3'-C4' | -2.84 | 96.68 | 102.73 |
| 1 | G | 10 | C49 | O4'-C1'-C2' | 3.26 | 112.51 | 106.25 |

All (2) chirality outliers are listed below:

| Mol | Chain | Res | Type | Atom |
|-----|-------|-----|------|------|
| 1 | F | 10 | C49 | C1' |
| 1 | G | 10 | C49 | C1' |

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

| Mol | Chain | Res | Type | Clashes | Symm-Clashes |
|-----|-------|-----|------|---------|--------------|
| 1 | F | 10 | C49 | 4 | 0 |
| 1 | G | 10 | C49 | 4 | 0 |

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.