



Full wwPDB NMR Structure Validation Report ⓘ

Feb 12, 2017 – 09:24 pm GMT

PDB ID : 2DC2
Title : Solution Structure of PDZ Domain
Authors : Li, X.; Wu, J.; Shi, Y.
Deposited on : 2005-12-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk28760
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	125131	11601
Ramachandran outliers	121729	10391
Sidechain outliers	121581	10367

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	103	

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:9-A:95 (87)	0.34	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 5, 7, 9, 10, 12, 13, 14, 19, 20
2	6, 11, 16
3	2, 15
4	4, 18
Single-model clusters	8; 17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1341 atoms, of which 683 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called golgi associated PDZ and coiled-coil motif containing isoform b.

Mol	Chain	Residues	Atoms						Trace
1	A	87	Total	C	H	N	O	S	0
			1341	415	683	119	123	1	

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP Q9HD26
A	96	LEU	-	CLONING ARTIFACT	UNP Q9HD26
A	97	GLU	-	CLONING ARTIFACT	UNP Q9HD26
A	98	HIS	-	CLONING ARTIFACT	UNP Q9HD26
A	99	HIS	-	CLONING ARTIFACT	UNP Q9HD26
A	100	HIS	-	CLONING ARTIFACT	UNP Q9HD26
A	101	HIS	-	CLONING ARTIFACT	UNP Q9HD26
A	102	HIS	-	CLONING ARTIFACT	UNP Q9HD26
A	103	HIS	-	CLONING ARTIFACT	UNP Q9HD26

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1 (medoid)

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.2 Score per residue for model 2

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.3 Score per residue for model 3

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.4 Score per residue for model 4

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.5 Score per residue for model 5

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.6 Score per residue for model 6

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.7 Score per residue for model 7

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.8 Score per residue for model 8

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.9 Score per residue for model 9

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.10 Score per residue for model 10

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.11 Score per residue for model 11

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.12 Score per residue for model 12

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.13 Score per residue for model 13

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.14 Score per residue for model 14

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.15 Score per residue for model 15

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b




4.2.16 Score per residue for model 16

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b



4.2.17 Score per residue for model 17

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b



- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b

HIS	MET
HIS	LVS
HIS	SER
HIS	GLN
HIS	GLY
HIS	VAL
	P9
	I10
	R11
	L16
	K17
	E18
	I25
	S26
	I27
	K31
	E32
	P36
	I37
	L38
	I39
	S40
	E41
	I42
	Q46
	L54
	I60
	R69
	D70
	T71
	K72
	H73
	V77
	T78
	I79
	Q83
	R84
	I87
	E88
	V91
	V92
	Y93
	V94
	A95
	LEU
	CU

- Molecule 1: golgi associated PDZ and coiled-coil motif containing isoform b

HIS	MET
HIS	LVS
HIS	SER
HIS	GLN
	GLY
	VAL
	GLY
	P9
	I10
	R11
	K12
	L16
	K17
	E18
	I25
	S26
	I27
	K31
	I37
	L38
	I39
	I42
	Q46
	C51
	L54
	D58
	A59
	I60
	D70
	H73
	K74
	V77
	R84
	E88
	F89
	E90
	V91
	V92
	Y93
	V94
	A95
	LEU
	GLU
	HIS
	HTS

5 Refinement protocol and experimental data overview

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
CNS	refinement	1.1

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	658	683	676	10±3
All	All	13160	13660	13520	199

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:23:LEU:HD12	1:A:25:ILE:HD11	0.81	1.52	17	1
1:A:60:ILE:HG21	1:A:68:LEU:HD13	0.77	1.57	15	3
1:A:39:ILE:HD13	1:A:54:LEU:HD21	0.76	1.57	12	3
1:A:48:ALA:HB1	1:A:54:LEU:HD13	0.75	1.55	17	1
1:A:25:ILE:HD13	1:A:39:ILE:HD12	0.74	1.59	11	3
1:A:68:LEU:HD21	1:A:76:ALA:HB1	0.71	1.61	17	2
1:A:60:ILE:HA	1:A:91:VAL:HG12	0.69	1.64	13	19
1:A:25:ILE:HG22	1:A:42:ILE:HG23	0.69	1.64	20	13
1:A:27:ILE:HG22	1:A:39:ILE:HD13	0.65	1.68	15	6
1:A:63:VAL:HG21	1:A:80:LEU:HD21	0.65	1.66	7	2
1:A:25:ILE:HG21	1:A:54:LEU:CD1	0.64	2.22	6	2
1:A:23:LEU:HD11	1:A:80:LEU:HD13	0.64	1.69	10	2
1:A:10:ILE:HA	1:A:92:VAL:HG12	0.63	1.69	20	7
1:A:23:LEU:HD21	1:A:80:LEU:CD2	0.63	2.24	15	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:68:LEU:HD21	1:A:76:ALA:CB	0.61	2.25	17	5
1:A:60:ILE:HG21	1:A:68:LEU:HD22	0.61	1.72	4	1
1:A:38:LEU:CD2	1:A:59:ALA:HB2	0.60	2.25	20	2
1:A:63:VAL:HG11	1:A:80:LEU:HD23	0.60	1.71	7	2
1:A:39:ILE:HD13	1:A:54:LEU:CD1	0.60	2.27	20	2
1:A:39:ILE:HD13	1:A:54:LEU:CD2	0.59	2.25	12	2
1:A:14:LEU:HD21	1:A:86:GLU:HG3	0.59	1.74	14	1
1:A:73:HIS:O	1:A:77:VAL:HG23	0.59	1.98	9	16
1:A:25:ILE:HG21	1:A:54:LEU:HD12	0.59	1.75	6	4
1:A:60:ILE:CG2	1:A:68:LEU:HD13	0.58	2.28	15	1
1:A:38:LEU:HD23	1:A:59:ALA:HB2	0.58	1.75	17	3
1:A:27:ILE:HD12	1:A:37:ILE:HG21	0.58	1.73	6	7
1:A:25:ILE:HG21	1:A:54:LEU:HD23	0.57	1.75	3	1
1:A:60:ILE:CG2	1:A:68:LEU:HD22	0.57	2.29	4	1
1:A:25:ILE:HG22	1:A:42:ILE:HA	0.57	1.77	3	1
1:A:48:ALA:CB	1:A:54:LEU:HD13	0.56	2.29	17	1
1:A:9:PRO:O	1:A:92:VAL:HG23	0.56	2.00	8	2
1:A:25:ILE:HG21	1:A:54:LEU:HD22	0.55	1.78	17	2
1:A:25:ILE:CG2	1:A:42:ILE:HG23	0.54	2.32	12	5
1:A:80:LEU:HD12	1:A:81:SER:N	0.54	2.17	10	3
1:A:39:ILE:HD13	1:A:54:LEU:HD11	0.54	1.79	20	1
1:A:76:ALA:HA	1:A:79:ILE:HD12	0.53	1.78	14	7
1:A:80:LEU:HD21	1:A:89:PHE:CE2	0.53	2.38	4	1
1:A:9:PRO:O	1:A:92:VAL:HG13	0.53	2.03	11	2
1:A:59:ALA:HB2	1:A:94:VAL:HG21	0.52	1.81	11	2
1:A:71:THR:HG21	1:A:79:ILE:CD1	0.51	2.35	19	1
1:A:25:ILE:CD1	1:A:39:ILE:HD12	0.51	2.35	11	1
1:A:23:LEU:HD12	1:A:25:ILE:CD1	0.51	2.33	17	1
1:A:68:LEU:HD21	1:A:76:ALA:HB2	0.51	1.83	8	3
1:A:25:ILE:N	1:A:25:ILE:HD13	0.50	2.21	17	1
1:A:63:VAL:HG21	1:A:80:LEU:CD2	0.50	2.37	7	1
1:A:42:ILE:HD13	1:A:54:LEU:O	0.49	2.08	5	3
1:A:10:ILE:HD12	1:A:61:LEU:HD12	0.49	1.84	11	1
1:A:42:ILE:HD11	1:A:55:HIS:C	0.48	2.29	5	2
1:A:36:PRO:O	1:A:38:LEU:HD12	0.48	2.09	15	2
1:A:25:ILE:HD12	1:A:54:LEU:HD13	0.48	1.86	16	1
1:A:23:LEU:HD22	1:A:25:ILE:HG13	0.47	1.87	3	2
1:A:25:ILE:HG23	1:A:48:ALA:CB	0.46	2.40	10	2
1:A:25:ILE:HD13	1:A:25:ILE:N	0.46	2.25	16	1
1:A:37:ILE:C	1:A:38:LEU:HD12	0.46	2.30	14	1
1:A:40:SER:O	1:A:56:VAL:HG13	0.46	2.09	17	2

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:39:ILE:HG21	1:A:54:LEU:HD21	0.46	1.86	12	1
1:A:48:ALA:HB1	1:A:54:LEU:HG	0.46	1.86	6	1
1:A:10:ILE:HG23	1:A:92:VAL:HG22	0.45	1.87	19	1
1:A:37:ILE:O	1:A:60:ILE:HD12	0.45	2.10	14	2
1:A:25:ILE:HD13	1:A:39:ILE:HG12	0.45	1.89	7	1
1:A:63:VAL:HG22	1:A:89:PHE:CE1	0.45	2.47	16	1
1:A:68:LEU:HD23	1:A:71:THR:OG1	0.44	2.12	8	1
1:A:77:VAL:HA	1:A:80:LEU:HD12	0.44	1.90	18	1
1:A:27:ILE:HG21	1:A:60:ILE:HD11	0.44	1.90	5	1
1:A:39:ILE:HD12	1:A:58:ASP:CB	0.44	2.42	2	1
1:A:25:ILE:CG2	1:A:54:LEU:HD13	0.43	2.43	16	1
1:A:83:GLN:HB3	1:A:87:ILE:HG23	0.43	1.89	9	1
1:A:63:VAL:HG11	1:A:80:LEU:CD2	0.43	2.43	9	1
1:A:37:ILE:HB	1:A:60:ILE:HD12	0.43	1.88	5	1
1:A:55:HIS:CE1	1:A:93:TYR:CD2	0.43	3.07	18	1
1:A:37:ILE:HD12	1:A:73:HIS:CD2	0.42	2.49	19	1
1:A:27:ILE:HD12	1:A:73:HIS:HE1	0.42	1.73	17	1
1:A:38:LEU:HD23	1:A:59:ALA:CB	0.42	2.44	2	1
1:A:68:LEU:HD11	1:A:76:ALA:HB1	0.42	1.89	3	1
1:A:23:LEU:HD23	1:A:25:ILE:O	0.42	2.14	10	1
1:A:27:ILE:HG22	1:A:39:ILE:HG12	0.42	1.92	4	1
1:A:80:LEU:HD21	1:A:89:PHE:CZ	0.42	2.49	4	1
1:A:25:ILE:HD13	1:A:39:ILE:HD13	0.42	1.91	10	1
1:A:59:ALA:HB2	1:A:94:VAL:CG2	0.42	2.45	3	1
1:A:37:ILE:HD12	1:A:73:HIS:ND1	0.41	2.30	15	1
1:A:13:VAL:HG23	1:A:89:PHE:HB3	0.41	1.91	13	1
1:A:10:ILE:O	1:A:10:ILE:HG23	0.41	2.16	12	1
1:A:83:GLN:HG3	1:A:87:ILE:HG23	0.41	1.92	19	1
1:A:60:ILE:HB	1:A:68:LEU:HD22	0.41	1.91	9	1
1:A:25:ILE:HD13	1:A:39:ILE:CD1	0.41	2.45	10	1
1:A:46:GLN:CB	1:A:47:PRO:HD2	0.41	2.46	18	1
1:A:55:HIS:CE1	1:A:93:TYR:CG	0.40	3.10	4	1
1:A:39:ILE:HD12	1:A:58:ASP:HB3	0.40	1.93	5	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation

was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/103 (83%)	80±1 (94±1%)	5±1 (6±1%)	0±0 (0±0%)	100	100
All	All	1700/2060 (83%)	1606 (94%)	94 (6%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	71/85 (84%)	55±3 (77±4%)	16±3 (23±4%)	3	29
All	All	1420/1700 (84%)	1094 (77%)	326 (23%)	3	29

All 42 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	93	TYR	20
1	A	46	GLN	19
1	A	89	PHE	16
1	A	72	LYS	13
1	A	58	ASP	13
1	A	23	LEU	12
1	A	54	LEU	12
1	A	11	ARG	12
1	A	84	ARG	11
1	A	17	LYS	11
1	A	18	GLU	11
1	A	12	LYS	11
1	A	16	LEU	10
1	A	51	CYS	9
1	A	38	LEU	9
1	A	40	SER	9
1	A	31	LYS	8
1	A	69	ARG	8
1	A	19	ASP	8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	78	THR	7
1	A	80	LEU	7
1	A	68	LEU	7
1	A	70	ASP	7
1	A	26	SER	6
1	A	41	GLU	6
1	A	88	GLU	6
1	A	74	LYS	6
1	A	14	LEU	5
1	A	75	GLU	5
1	A	15	LEU	5
1	A	32	GLU	5
1	A	90	GLU	4
1	A	50	ARG	4
1	A	81	SER	4
1	A	49	ASP	4
1	A	10	ILE	4
1	A	71	THR	3
1	A	25	ILE	3
1	A	67	ASN	2
1	A	21	GLU	2
1	A	86	GLU	1
1	A	64	ASN	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided