



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2017 – 07:03 AM EDT

PDB ID : 3DCD
Title : X-ray structure of the galactose mutarotase related enzyme Q5FKD7 from *Lactobacillus acidophilus* at the resolution 1.9Å. Northeast Structural Genomics consortium target LaR33.
Authors : Kuzin, A.P.; Lew, S.; Vorobiev, S.M.; Seetharaman, J.; Wang, H.; Mao, L.; Foote, E.L.; Xiao, R.; Nair, R.; Baran, M.C.; Acton, T.B.; Rost, B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : unknown
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030345
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030345


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	5047 (1.90-1.90)
Clashscore	112137	5731 (1.90-1.90)
Ramachandran outliers	110173	5669 (1.90-1.90)
Sidechain outliers	110143	5670 (1.90-1.90)
RSRZ outliers	101464	5100 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 1%, orange 1%, orange 87%, yellow 87%, yellow 91%, green 91%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 0% 87% 10% .. </div> </div>
1	B	307	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 0%, red 3%, orange 3%, orange 91%, yellow 91%, yellow 94%, green 94%, green 100%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 3% 91% 7% . </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Galactose mutarotase related enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	300	Total	C	N	O	S	Se	5	3	0
			2474	1573	418	472	1	10			
1	B	300	Total	C	N	O	S	Se	0	0	0
			2459	1562	418	470	1	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	300	LEU	-	expression tag	UNP Q5FKD7
A	301	GLU	-	expression tag	UNP Q5FKD7
A	302	HIS	-	expression tag	UNP Q5FKD7
A	303	HIS	-	expression tag	UNP Q5FKD7
A	304	HIS	-	expression tag	UNP Q5FKD7
A	305	HIS	-	expression tag	UNP Q5FKD7
A	306	HIS	-	expression tag	UNP Q5FKD7
A	307	HIS	-	expression tag	UNP Q5FKD7
B	300	LEU	-	expression tag	UNP Q5FKD7
B	301	GLU	-	expression tag	UNP Q5FKD7
B	302	HIS	-	expression tag	UNP Q5FKD7
B	303	HIS	-	expression tag	UNP Q5FKD7
B	304	HIS	-	expression tag	UNP Q5FKD7
B	305	HIS	-	expression tag	UNP Q5FKD7
B	306	HIS	-	expression tag	UNP Q5FKD7
B	307	HIS	-	expression tag	UNP Q5FKD7

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	244	Total	O	0	0
			244	244		
2	B	243	Total	O	0	0
			243	243		

- Molecule 1: Galactose mutarotase related enzyme



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.27Å 88.27Å 139.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.74 – 1.90 19.57 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.74-1.90) 96.2 (19.57-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.27 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.178 , 0.206 0.177 , 0.205	Depositor DCC
R_{free} test set	2590 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	17.8	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5420	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	2/2545 (0.1%)	0.65	0/3437
1	B	0.54	0/2521	0.64	0/3406
All	All	0.56	2/5066 (0.0%)	0.64	0/6843

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	9[A]	MSE	CG-SE	-6.83	1.72	1.95
1	A	9[B]	MSE	CG-SE	-6.83	1.72	1.95

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2474	0	2344	36	0
1	B	2459	0	2322	15	0
2	A	244	0	0	2	0
2	B	243	0	0	1	0
All	All	5420	0	4666	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MSE:HE3	1:A:2:ASP:O	1.48	1.13
1:A:215:LYS:HD3	1:A:230:TRP:CZ3	1.92	1.04
1:A:128:VAL:HG12	1:A:136:MSE:HE1	1.63	0.79
1:A:168:THR:HG23	1:A:187:ALA:O	1.83	0.77
1:A:54:GLY:H	1:A:73:HIS:HD2	1.33	0.75
1:A:54:GLY:H	1:A:73:HIS:CD2	2.07	0.71
1:A:125:ASN:HD21	1:A:232:ARG:HH22	1.36	0.71
1:A:114:ASN:HD21	1:A:116:ASN:HD22	1.46	0.64
1:B:114:ASN:HD22	1:B:125:ASN:HD22	1.45	0.63
1:A:155:ASN:HD22	1:A:157:GLU:H	1.47	0.62
1:A:215:LYS:CD	1:A:230:TRP:CZ3	2.78	0.60
1:A:7:ASN:ND2	1:A:9[B]:MSE:H	1.99	0.60
1:A:128:VAL:CG1	1:A:136:MSE:HE1	2.30	0.60
1:A:1:MSE:CE	1:A:2:ASP:O	2.36	0.59
1:A:136:MSE:HE2	1:A:277:LEU:HD23	1.84	0.58
1:A:125:ASN:ND2	1:A:232:ARG:HH12	2.01	0.57
1:A:215:LYS:HD3	1:A:230:TRP:CE3	2.39	0.57
1:A:7:ASN:ND2	1:A:10:ILE:H	2.05	0.55
1:B:169:ARG:HH21	1:B:214:ASN:HD22	1.55	0.54
1:B:7:ASN:HB3	1:B:9:MSE:H	1.72	0.54
1:B:7:ASN:ND2	1:B:119:ASN:O	2.41	0.53
1:A:17:HIS:HD2	1:A:80:ASP:OD1	1.92	0.52
1:A:155:ASN:ND2	1:A:157:GLU:H	2.07	0.51
1:A:125:ASN:HD22	1:A:232:ARG:HH12	1.57	0.51
1:A:125:ASN:ND2	1:A:232:ARG:HH22	2.07	0.51
1:A:7:ASN:HD22	1:A:9[B]:MSE:H	1.58	0.50
1:A:128:VAL:HG12	1:A:136:MSE:CE	2.39	0.50
1:A:7:ASN:C	1:A:7:ASN:HD22	2.15	0.50
1:A:17:HIS:HE1	2:A:548:HOH:O	1.95	0.49
1:A:169:ARG:HH21	1:A:214:ASN:HD22	1.59	0.49
1:A:128:VAL:CG1	1:A:136:MSE:CE	2.91	0.48
1:B:207:TYR:CE2	1:B:239:ILE:HD13	2.48	0.48
1:B:143:HIS:HA	1:B:253:GLU:HG2	1.95	0.48
1:B:207:TYR:HE2	1:B:239:ILE:HD13	1.79	0.47
1:A:215:LYS:HG2	1:A:230:TRP:CE3	2.48	0.47
1:B:169:ARG:HH21	1:B:214:ASN:ND2	2.11	0.47
1:A:169:ARG:HH21	1:A:214:ASN:ND2	2.14	0.46
1:B:94:LEU:HD21	1:B:96:LYS:HE3	1.98	0.45
1:A:49:LEU:HD22	1:A:73:HIS:CG	2.51	0.45
1:A:226:HIS:CE1	1:A:228:ASN:HD22	2.34	0.45
1:A:73:HIS:HE1	2:A:344:HOH:O	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:164:HIS:HA	1:B:165:PRO:C	2.38	0.45
1:B:239:ILE:N	1:B:239:ILE:HD12	2.32	0.44
1:A:114:ASN:ND2	1:A:116:ASN:HD22	2.12	0.43
1:A:235:PRO:HD2	1:A:255:TRP:O	2.19	0.43
1:B:72:GLN:NE2	2:B:403:HOH:O	2.51	0.43
1:A:254:PRO:HG2	1:A:287:PHE:CG	2.54	0.43
1:B:35:TRP:O	1:B:45:HIS:HD2	2.03	0.41
1:B:17:HIS:HD2	1:B:80:ASP:OD1	2.03	0.40
1:A:10:ILE:HD11	1:A:117:LEU:HD22	2.02	0.40
1:B:175:LYS:HE3	1:B:175:LYS:HB2	1.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/307 (98%)	291 (97%)	10 (3%)	0	100	100
1	B	298/307 (97%)	287 (96%)	11 (4%)	0	100	100
All	All	599/614 (98%)	578 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/267 (102%)	268 (99%)	3 (1%)	78	77
1	B	268/267 (100%)	268 (100%)	0	100	100
All	All	539/534 (101%)	536 (99%)	3 (1%)	89	90

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	212	ASN
1	A	300	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	17	HIS
1	A	45	HIS
1	A	73	HIS
1	A	114	ASN
1	A	119	ASN
1	A	125	ASN
1	A	147	ASN
1	A	155	ASN
1	A	182	ASN
1	A	212	ASN
1	A	214	ASN
1	A	228	ASN
1	A	276	HIS
1	B	17	HIS
1	B	45	HIS
1	B	69	HIS
1	B	72	GLN
1	B	114	ASN
1	B	116	ASN
1	B	147	ASN
1	B	214	ASN
1	B	276	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/307 (95%)	-0.07	3 (1%) 82 84	12, 18, 26, 33	0
1	B	292/307 (95%)	-0.02	9 (3%) 49 53	13, 18, 26, 33	0
All	All	584/614 (95%)	-0.04	12 (2%) 64 67	12, 18, 26, 33	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300	LEU	7.8
1	B	300	LEU	7.4
1	A	152	HIS	3.3
1	A	230	TRP	3.1
1	B	151	ASP	3.0
1	B	152	HIS	2.6
1	B	153	GLY	2.5
1	B	299	LYS	2.4
1	B	298	VAL	2.4
1	B	164	HIS	2.3
1	B	230	TRP	2.2
1	B	133	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.