



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 2, 2017 – 08:28 PM EDT

PDB ID : 2DEP  
Title : Crystal Structure of xylanase B from Clostridium stercorarium F9  
Authors : Fushinobu, S.; Nishimoto, M.; Miyanaga, A.; Kitaoka, M.; Hayashi, K.  
Deposited on : unknown  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20030345  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20030345


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	4827 (1.80-1.80)
Clashscore	112137	5742 (1.80-1.80)
Ramachandran outliers	110173	5676 (1.80-1.80)
Sidechain outliers	110143	5675 (1.80-1.80)
RSRZ outliers	101464	4906 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 1%, green 85%, grey 9%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>1%</span> <span>85%</span> <span>9%</span> </div> </div>
1	B	356	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 11%, green 83%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> <span>1%</span> <span>83%</span> <span>11%</span> <span>5%</span> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thermostable cellobiohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	0	0	0
			2778	1786	463	522	7			
1	B	339	Total	C	N	O	S	0	0	0
			2770	1782	462	519	7			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	MET	-	INITIATING METHIONINE	UNP P40942
A	131	ASP	THR	SEE REMARK 999	UNP P40942
A	132	TRP	GLY	SEE REMARK 999	UNP P40942
A	134	PHE	SER	SEE REMARK 999	UNP P40942
A	388	LEU	-	EXPRESSION TAG	UNP P40942
A	389	GLU	-	EXPRESSION TAG	UNP P40942
A	390	HIS	-	EXPRESSION TAG	UNP P40942
A	391	HIS	-	EXPRESSION TAG	UNP P40942
A	392	HIS	-	EXPRESSION TAG	UNP P40942
A	393	HIS	-	EXPRESSION TAG	UNP P40942
A	394	HIS	-	EXPRESSION TAG	UNP P40942
A	395	HIS	-	EXPRESSION TAG	UNP P40942
B	40	MET	-	INITIATING METHIONINE	UNP P40942
B	131	ASP	THR	SEE REMARK 999	UNP P40942
B	132	TRP	GLY	SEE REMARK 999	UNP P40942
B	134	PHE	SER	SEE REMARK 999	UNP P40942
B	388	LEU	-	EXPRESSION TAG	UNP P40942
B	389	GLU	-	EXPRESSION TAG	UNP P40942
B	390	HIS	-	EXPRESSION TAG	UNP P40942
B	391	HIS	-	EXPRESSION TAG	UNP P40942
B	392	HIS	-	EXPRESSION TAG	UNP P40942
B	393	HIS	-	EXPRESSION TAG	UNP P40942
B	394	HIS	-	EXPRESSION TAG	UNP P40942
B	395	HIS	-	EXPRESSION TAG	UNP P40942


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	365	Total 365	O 365	0	0
2	B	343	Total 343	O 343	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Thermostable cellobiohydrolase



- Molecule 1: Thermostable cellobiohydrolase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	64.76 Å 96.60 Å 138.44 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.14 – 1.80 50.14 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.14-1.80) 99.7 (50.14-1.80)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.66 (at 1.79 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.184 , 0.210 0.185 , 0.182	Depositor DCC
$R_{free}$ test set	4060 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtriage
Anisotropy	0.313	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6256	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/2852	0.57	0/3877
1	B	0.31	0/2844	0.58	0/3866
All	All	0.31	0/5696	0.58	0/7743

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2778	0	2707	30	0
1	B	2770	0	2703	41	0
2	A	365	0	0	5	0
2	B	343	0	0	6	0
All	All	6256	0	5410	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:361:ASN:HD22	1:B:362:ALA:H	1.20	0.88

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:93:GLN:HE21	1:B:97:GLY:H	1.19	0.88
1:A:93:GLN:HE21	1:A:97:GLY:H	1.22	0.84
1:A:361:ASN:HD22	1:A:362:ALA:H	1.24	0.84
1:A:272:VAL:HG21	1:A:325:GLU:HG2	1.64	0.79
1:B:247:LYS:O	1:B:251:GLU:HG3	1.90	0.71
1:B:302:ASN:HD22	1:B:302:ASN:H	1.42	0.67
1:B:93:GLN:NE2	1:B:97:GLY:H	1.94	0.66
1:B:146:THR:HG23	1:B:147:ASP:H	1.60	0.66
1:A:93:GLN:NE2	1:A:97:GLY:H	1.96	0.63
1:A:167:ARG:NH1	2:A:671:HOH:O	2.32	0.63
1:B:137:LYS:HE3	1:B:137:LYS:H	1.64	0.62
1:A:378:ALA:O	1:A:382:PRO:HB3	2.00	0.61
1:B:137:LYS:H	1:B:137:LYS:CE	2.14	0.60
1:B:146:THR:O	1:B:147:ASP:C	2.39	0.60
1:A:214:ARG:O	1:A:218:GLU:HG3	2.03	0.59
1:A:298:ILE:H	1:A:361:ASN:HD21	1.51	0.58
1:B:146:THR:HG23	1:B:147:ASP:N	2.18	0.57
1:B:361:ASN:HD22	1:B:362:ALA:N	1.98	0.56
1:A:252:LYS:NZ	1:A:252:LYS:HB3	2.22	0.55
1:A:136:ASP:OD2	1:A:140:LYS:HG3	2.07	0.54
1:A:262:GLN:HE21	1:A:264:HIS:HD2	1.54	0.54
1:B:302:ASN:HD22	1:B:302:ASN:N	2.00	0.54
1:A:302:ASN:HB2	1:B:139:GLY:HA3	1.90	0.54
1:B:90:ALA:HA	1:B:130:PRO:HG3	1.90	0.54
1:A:310:SER:OG	1:A:311:ILE:N	2.41	0.54
1:A:310:SER:O	1:A:311:ILE:HB	2.08	0.53
1:A:361:ASN:HD22	1:A:362:ALA:N	2.02	0.53
1:A:358:LYS:HD3	2:A:750:HOH:O	2.08	0.52
1:A:262:GLN:HE21	1:A:264:HIS:CD2	2.27	0.52
1:B:302:ASN:H	1:B:302:ASN:ND2	2.07	0.52
1:A:138:GLU:HB2	1:A:140:LYS:HE3	1.91	0.52
1:A:223:ASP:HB2	2:A:731:HOH:O	2.10	0.51
1:B:237:VAL:HG13	1:B:238:LYS:N	2.26	0.51
1:B:298:ILE:H	1:B:361:ASN:HD21	1.57	0.51
1:B:121:HIS:HE1	2:B:545:HOH:O	1.94	0.51
1:B:160:GLN:HE21	1:B:164:ASN:ND2	2.08	0.51
1:B:293:GLU:HG2	1:B:343:TRP:CE3	2.47	0.49
1:B:223:ASP:HB2	2:B:517:HOH:O	2.12	0.49
1:A:94:PRO:HG2	2:A:463:HOH:O	2.12	0.49
1:B:381:ASP:N	1:B:382:PRO:HD3	2.26	0.48
1:A:140:LYS:NZ	2:A:535:HOH:O	2.47	0.47

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:197:ASN:ND2	1:B:202:GLN:HG3	2.30	0.47
1:B:245:LEU:HD13	1:B:245:LEU:C	2.35	0.47
1:B:160:GLN:HE21	1:B:164:ASN:HD21	1.63	0.46
1:B:334:LYS:HD2	1:B:335:ASP:CG	2.37	0.45
1:A:138:GLU:HB2	1:A:140:LYS:CE	2.47	0.45
1:A:245:LEU:O	1:A:249:LEU:HG	2.17	0.44
1:A:246:VAL:HG11	1:A:287:LEU:HD11	1.99	0.44
1:B:167:ARG:HH21	1:B:167:ARG:HG2	1.82	0.44
1:B:382:PRO:HG2	2:B:611:HOH:O	2.17	0.44
1:A:353:ASN:O	1:A:358:LYS:HD2	2.18	0.43
1:A:231:TYR:HB3	1:A:262:GLN:NE2	2.33	0.43
1:A:60:ALA:HA	1:A:82:VAL:HG13	2.00	0.43
1:A:309:ASP:CG	1:A:310:SER:H	2.21	0.43
1:B:382:PRO:HG3	2:B:476:HOH:O	2.18	0.43
1:B:293:GLU:HG2	1:B:343:TRP:CZ3	2.54	0.43
1:A:310:SER:O	1:A:311:ILE:CB	2.66	0.43
1:B:362:ALA:N	1:B:363:PRO:HD3	2.34	0.43
1:B:272:VAL:HG21	1:B:325:GLU:HG2	2.01	0.42
1:B:334:LYS:HD2	1:B:335:ASP:OD1	2.20	0.42
1:B:382:PRO:CG	2:B:476:HOH:O	2.67	0.42
1:B:49:GLU:OE2	1:B:52:ARG:HD3	2.19	0.42
1:A:362:ALA:N	1:A:363:PRO:HD3	2.35	0.42
1:B:81:LEU:HD12	1:B:111:ALA:HB2	2.03	0.41
1:B:380:VAL:C	1:B:382:PRO:HD3	2.40	0.41
1:B:302:ASN:ND2	1:B:302:ASN:N	2.67	0.41
1:B:147:ASP:O	1:B:149:GLN:N	2.54	0.41
1:B:199:PRO:HD2	2:B:630:HOH:O	2.20	0.41
1:B:44:ILE:HA	1:B:45:PRO:HD3	1.92	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/356 (95%)	328 (97%)	8 (2%)	2 (1%)	28	13
1	B	337/356 (95%)	322 (96%)	14 (4%)	1 (0%)	44	29
All	All	675/712 (95%)	650 (96%)	22 (3%)	3 (0%)	38	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	310	SER
1	A	311	ILE
1	B	148	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/314 (95%)	295 (99%)	4 (1%)	73	66
1	B	298/314 (95%)	290 (97%)	8 (3%)	50	35
All	All	597/628 (95%)	585 (98%)	12 (2%)	60	48

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	PHE
1	A	140	LYS
1	A	316	LEU
1	A	361	ASN
1	B	55	PHE
1	B	137	LYS
1	B	149	GLN
1	B	197	ASN
1	B	302	ASN
1	B	316	LEU
1	B	334	LYS
1	B	361	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	GLN
1	A	109	GLN
1	A	127	ASN
1	A	190	ASN
1	A	262	GLN
1	A	361	ASN
1	B	93	GLN
1	B	121	HIS
1	B	127	ASN
1	B	149	GLN
1	B	154	ASN
1	B	164	ASN
1	B	190	ASN
1	B	197	ASN
1	B	302	ASN
1	B	324	GLN
1	B	361	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	340/356 (95%)	-0.18	3 (0%) 84 82	11, 18, 31, 45	0
1	B	339/356 (95%)	-0.13	5 (1%) 74 70	10, 16, 32, 52	0
All	All	679/712 (95%)	-0.15	8 (1%) 79 77	10, 17, 31, 52	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	382	PRO	6.3
1	B	146	THR	5.5
1	B	147	ASP	4.6
1	B	148	PRO	4.2
1	A	309	ASP	3.5
1	B	149	GLN	3.4
1	A	310	SER	3.0
1	A	69	GLN	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.