



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 12:05 pm GMT

PDB ID : 4DH2
Title : Crystal structure of Coh-OlpC(Cthe_0452)-Doc435(Cthe_0435) complex: A novel type I Cohesin-Dockerin complex from Clostridium thermocellum ATTC 27405
Authors : Alves, V.D.; Carvalho, A.L.; Najmudin, S.H.; Bras, J.; Prates, J.A.M.; Fontes, C.M.G.A.
Deposited on : 2012-01-27
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1762 (1.76-1.76)
Clashscore	112137	1889 (1.76-1.76)
Ramachandran outliers	110173	1868 (1.76-1.76)
Sidechain outliers	110143	1868 (1.76-1.76)
RSRZ outliers	101464	1770 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>13%</div> <div>7%</div> </div> </div>
1	C	170	<div> <div></div> <div> <div>86%</div> <div>6%</div> <div>8%</div> </div> </div>
2	B	82	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>•</div> <div>12%</div> </div> </div>
2	D	82	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>•</div> <div>15%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	B	103	-	-	-	X
4	CA	B	101	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7829 atoms, of which 3648 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellulosome anchoring protein cohesin region.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	158	Total	C	H	N	O	S	0	4	0
			2539	825	1268	192	253	1			
1	C	156	Total	C	H	N	O	S	0	4	0
			2522	814	1268	190	249	1			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP A3DCL1
A	2	ALA	-	EXPRESSION TAG	UNP A3DCL1
A	3	SER	-	EXPRESSION TAG	UNP A3DCL1
A	163	LEU	-	EXPRESSION TAG	UNP A3DCL1
A	164	GLU	-	EXPRESSION TAG	UNP A3DCL1
A	165	HIS	-	EXPRESSION TAG	UNP A3DCL1
A	166	HIS	-	EXPRESSION TAG	UNP A3DCL1
A	167	HIS	-	EXPRESSION TAG	UNP A3DCL1
A	168	HIS	-	EXPRESSION TAG	UNP A3DCL1
A	169	HIS	-	EXPRESSION TAG	UNP A3DCL1
A	170	HIS	-	EXPRESSION TAG	UNP A3DCL1
C	1	MET	-	INITIATING METHIONINE	UNP A3DCL1
C	2	ALA	-	EXPRESSION TAG	UNP A3DCL1
C	3	SER	-	EXPRESSION TAG	UNP A3DCL1
C	163	LEU	-	EXPRESSION TAG	UNP A3DCL1
C	164	GLU	-	EXPRESSION TAG	UNP A3DCL1
C	165	HIS	-	EXPRESSION TAG	UNP A3DCL1
C	166	HIS	-	EXPRESSION TAG	UNP A3DCL1
C	167	HIS	-	EXPRESSION TAG	UNP A3DCL1
C	168	HIS	-	EXPRESSION TAG	UNP A3DCL1
C	169	HIS	-	EXPRESSION TAG	UNP A3DCL1
C	170	HIS	-	EXPRESSION TAG	UNP A3DCL1


- Molecule 2 is a protein called Dockerin type 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	72	Total	C	H	N	O	S	0	0	0
			1142	371	560	97	110	4			
2	D	70	Total	C	H	N	O	S	0	0	0
			1116	359	552	95	106	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MET	-	INITIATING METHIONINE	UNP A3DCJ4
D	1	MET	-	INITIATING METHIONINE	UNP A3DCJ4

- Molecule 3 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total	Ca	0	0
			2	2		
4	D	2	Total	Ca	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	163	Total	O	0	0
			163	163		
5	B	65	Total	O	0	0
			65	65		
5	C	177	Total	O	0	0
			177	177		
5	D	56	Total	O	0	0
			56	56		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cellulosome anchoring protein cohesin region




- Molecule 1: Cellulosome anchoring protein cohesin region



- Molecule 2: Dockerin type 1



- Molecule 2: Dockerin type 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.76Å 90.76Å 135.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.30 – 1.75 43.02 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.9 (39.30-1.75) 96.6 (43.02-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.75Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7.2_865)	Depositor
R, R_{free}	0.181 , 0.215 0.183 , 0.216	Depositor DCC
R_{free} test set	3323 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	22.4	Xtriage
Anisotropy	0.154	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 51.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7829	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.53 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.3041e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1306	0.69	0/1763
1	C	0.56	0/1289	0.69	0/1740
2	B	0.52	0/593	0.62	0/805
2	D	0.46	0/574	0.56	0/779
All	All	0.53	0/3762	0.66	0/5087

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1271	1268	1283	18	0
1	C	1254	1268	1266	8	0
2	B	582	560	565	2	0
2	D	564	552	551	2	0
3	A	15	0	0	1	0
3	B	10	0	0	0	0
3	C	20	0	0	0	0
4	B	2	0	0	0	0
4	D	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	163	0	0	3	0
5	B	65	0	0	0	0
5	C	177	0	0	0	0
5	D	56	0	0	2	0
All	All	4181	3648	3665	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:201:SO4:O4	5:A:455:HOH:O	1.99	0.80
2:D:62:ARG:NH2	5:D:256:HOH:O	2.19	0.74
1:A:80:TYR:OH	5:A:392:HOH:O	2.06	0.72
1:A:112:LYS:NZ	5:A:317:HOH:O	2.36	0.57
1:A:163:LEU:O	1:A:164:GLU:HB2	2.03	0.57
1:C:53:LEU:HD21	1:C:136:PHE:CE1	2.42	0.55
1:A:53:LEU:HD21	1:A:136:PHE:CE1	2.42	0.54
2:D:69:ARG:NH2	5:D:251:HOH:O	2.39	0.53
1:A:37:LYS:NZ	1:A:39[B]:GLU:OE2	2.41	0.53
1:A:53:LEU:CD2	1:A:136:PHE:CE1	2.93	0.52
1:A:8:HIS:HD2	1:C:155:GLU:OE2	1.95	0.49
1:A:53:LEU:HD23	1:A:136:PHE:CD1	2.49	0.48
1:A:36:LEU:HD13	1:A:53:LEU:HD11	1.97	0.47
1:A:26:ALA:HB3	1:A:117:ILE:CD1	2.45	0.46
2:B:21:TYR:CZ	2:B:25:LYS:HE3	2.51	0.46
1:A:24:VAL:HG13	1:A:158:VAL:HG13	1.97	0.45
1:A:37:LYS:CE	1:A:39[B]:GLU:OE2	2.66	0.43
1:C:53:LEU:CD2	1:C:136:PHE:CE1	3.02	0.43
1:A:8:HIS:CE1	1:A:139:SER:HB3	2.52	0.43
1:A:13:ALA:HB3	1:A:150:LEU:HG	2.00	0.43
1:A:94:ASP:OD2	2:B:25:LYS:HD2	2.19	0.43
1:C:72:THR:O	1:C:73:ASP:C	2.57	0.42
1:C:13:ALA:HB3	1:C:150:LEU:HG	2.01	0.42
1:A:62:VAL:HG21	1:A:89:ILE:HD11	2.02	0.41
1:C:53:LEU:CD2	1:C:136:PHE:CD1	3.03	0.41
1:C:17:LEU:HD12	1:C:17:LEU:C	2.41	0.41
1:A:37:LYS:HE2	1:A:39[B]:GLU:OE2	2.20	0.41
1:C:64:LYS:HE3	1:C:66[B]:GLU:CG	2.51	0.41
1:A:72:THR:O	1:A:73:ASP:C	2.59	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/170 (94%)	158 (99%)	2 (1%)	0	100	100
1	C	158/170 (93%)	157 (99%)	1 (1%)	0	100	100
2	B	70/82 (85%)	68 (97%)	2 (3%)	0	100	100
2	D	68/82 (83%)	66 (97%)	2 (3%)	0	100	100
All	All	456/504 (90%)	449 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/150 (96%)	144 (100%)	0	100	100
1	C	142/150 (95%)	142 (100%)	0	100	100
2	B	64/73 (88%)	64 (100%)	0	100	100
2	D	62/73 (85%)	62 (100%)	0	100	100
All	All	412/446 (92%)	412 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
2	B	58	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	A	201	-	4,4,4	0.40	0	6,6,6	0.28	0
3	SO4	A	202	-	4,4,4	0.16	0	6,6,6	0.44	0
3	SO4	A	203	-	4,4,4	0.15	0	6,6,6	0.20	0
3	SO4	B	103	-	4,4,4	0.15	0	6,6,6	0.32	0
3	SO4	B	104	-	4,4,4	0.17	0	6,6,6	0.13	0
3	SO4	C	201	-	4,4,4	0.16	0	6,6,6	0.30	0
3	SO4	C	202	-	4,4,4	0.17	0	6,6,6	0.23	0
3	SO4	C	203	-	4,4,4	0.18	0	6,6,6	0.18	0
3	SO4	C	204	-	4,4,4	0.19	0	6,6,6	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means

no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	201	-	-	0/0/0/0	0/0/0/0
3	SO4	A	202	-	-	0/0/0/0	0/0/0/0
3	SO4	A	203	-	-	0/0/0/0	0/0/0/0
3	SO4	B	103	-	-	0/0/0/0	0/0/0/0
3	SO4	B	104	-	-	0/0/0/0	0/0/0/0
3	SO4	C	201	-	-	0/0/0/0	0/0/0/0
3	SO4	C	202	-	-	0/0/0/0	0/0/0/0
3	SO4	C	203	-	-	0/0/0/0	0/0/0/0
3	SO4	C	204	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	201	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	158/170 (92%)	-0.07	4 (2%) 58 64	15, 27, 53, 116	0
1	C	156/170 (91%)	0.01	0 100 100	17, 26, 43, 59	0
2	B	72/82 (87%)	-0.12	1 (1%) 75 83	16, 26, 59, 85	0
2	D	70/82 (85%)	0.23	4 (5%) 24 31	21, 32, 59, 90	0
All	All	456/504 (90%)	-0.00	9 (1%) 65 73	15, 27, 54, 116	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	7	ILE	3.7
2	D	12	ALA	3.7
2	D	74	ASN	2.8
1	A	127	SER	2.7
2	B	76	TYR	2.5
1	A	128	THR	2.3
2	D	38	ASP	2.3
1	A	123	LYS	2.3
2	D	35	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	B	103	5/5	0.93	0.18	4.62	42,49,60,69	1
4	CA	B	101	1/1	0.99	0.12	2.29	20,20,20,20	0
3	SO4	C	204	5/5	0.94	0.18	0.89	45,82,87,89	0
3	SO4	A	202	5/5	0.98	0.11	0.38	52,58,72,72	0
4	CA	D	102	1/1	0.98	0.08	-0.36	27,27,27,27	0
4	CA	D	101	1/1	0.98	0.07	-0.67	29,29,29,29	0
4	CA	B	102	1/1	0.99	0.06	-0.76	21,21,21,21	0
3	SO4	A	201	5/5	0.98	0.07	-0.91	25,34,40,48	0
3	SO4	C	201	5/5	0.98	0.08	-1.95	34,39,55,59	0
3	SO4	C	202	5/5	0.91	0.15	-	65,84,86,91	0
3	SO4	C	203	5/5	0.95	0.23	-	55,78,83,87	0
3	SO4	B	104	5/5	0.95	0.14	-	69,72,73,74	0
3	SO4	A	203	5/5	0.95	0.12	-	59,74,77,82	0

6.5 Other polymers [i](#)

There are no such residues in this entry.