



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2017 – 09:11 PM EDT

PDB ID : 5DHU  
Title : Crystal structure of NAD kinase 1 from *Listeria monocytogenes* in complex with a novel inhibitor  
Authors : Gelin, M.; Paoletti, J.; Assairi, L.; Huteau, V.; Pochet, S.; Labesse, G.  
Deposited on : unknown  
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20029824  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20029824


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1570 (2.36-2.32)
Clashscore	112137	1673 (2.36-2.32)
Ramachandran outliers	110173	1654 (2.36-2.32)
Sidechain outliers	110143	1655 (2.36-2.32)
RSRZ outliers	101464	1576 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div>3%</div> <div>89%</div> <div>7%</div> <div>•</div> </div>
1	B	272	<div> <div>6%</div> <div>78%</div> <div>18%</div> <div>•</div> </div>
1	C	272	<div> <div>8%</div> <div>87%</div> <div>6%</div> <div>7%</div> </div>
1	D	272	<div> <div>7%</div> <div>80%</div> <div>10%</div> <div>10%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	301	-	-	-	X
4	CIT	B	301	-	-	-	X

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8425 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	1	0
			2065	1323	345	388	9			
1	B	262	Total	C	N	O	S	0	2	0
			2107	1352	354	392	9			
1	C	254	Total	C	N	O	S	0	0	0
			1981	1271	332	369	9			
1	D	244	Total	C	N	O	S	0	0	0
			1894	1211	323	352	8			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	LEU	-	expression tag	UNP Q8Y8D7
A	266	GLU	-	expression tag	UNP Q8Y8D7
A	267	HIS	-	expression tag	UNP Q8Y8D7
A	268	HIS	-	expression tag	UNP Q8Y8D7
A	269	HIS	-	expression tag	UNP Q8Y8D7
A	270	HIS	-	expression tag	UNP Q8Y8D7
A	271	HIS	-	expression tag	UNP Q8Y8D7
A	272	HIS	-	expression tag	UNP Q8Y8D7
B	265	LEU	-	expression tag	UNP Q8Y8D7
B	266	GLU	-	expression tag	UNP Q8Y8D7
B	267	HIS	-	expression tag	UNP Q8Y8D7
B	268	HIS	-	expression tag	UNP Q8Y8D7
B	269	HIS	-	expression tag	UNP Q8Y8D7
B	270	HIS	-	expression tag	UNP Q8Y8D7
B	271	HIS	-	expression tag	UNP Q8Y8D7
B	272	HIS	-	expression tag	UNP Q8Y8D7
C	265	LEU	-	expression tag	UNP Q8Y8D7
C	266	GLU	-	expression tag	UNP Q8Y8D7
C	267	HIS	-	expression tag	UNP Q8Y8D7
C	268	HIS	-	expression tag	UNP Q8Y8D7
C	269	HIS	-	expression tag	UNP Q8Y8D7

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	270	HIS	-	expression tag	UNP Q8Y8D7
C	271	HIS	-	expression tag	UNP Q8Y8D7
C	272	HIS	-	expression tag	UNP Q8Y8D7
D	265	LEU	-	expression tag	UNP Q8Y8D7
D	266	GLU	-	expression tag	UNP Q8Y8D7
D	267	HIS	-	expression tag	UNP Q8Y8D7
D	268	HIS	-	expression tag	UNP Q8Y8D7
D	269	HIS	-	expression tag	UNP Q8Y8D7
D	270	HIS	-	expression tag	UNP Q8Y8D7
D	271	HIS	-	expression tag	UNP Q8Y8D7
D	272	HIS	-	expression tag	UNP Q8Y8D7

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 5'-azido-5'-deoxy-8-[(2-{[2-(1H-indol-3-yl)ethyl]amino}-2-oxoethyl)sulfanyl]adenosine (three-letter code: 5A8) (formula: C<sub>22</sub>H<sub>24</sub>N<sub>10</sub>O<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			37	22	10	4	1		
3	B	1	Total	C	N	O	S	0	0
			37	22	10	4	1		
3	C	1	Total	C	N	O	S	0	0
			37	22	10	4	1		
3	D	1	Total	C	N	O	S	0	0
			37	22	10	4	1		

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C<sub>6</sub>H<sub>8</sub>O<sub>7</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			13	6	7		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	74	Total	O	0	0
			74	74		
5	B	34	Total	O	0	0
			34	34		
5	C	53	Total	O	0	0
			53	53		
5	D	50	Total	O	0	0
			50	50		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD kinase 1




- Molecule 1: NAD kinase 1




- Molecule 1: NAD kinase 1



- Molecule 1: NAD kinase 1







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.92Å 119.34Å 67.75Å 90.00° 100.37° 90.00°	Depositor
Resolution (Å)	47.45 – 2.33 59.67 – 2.33	Depositor EDS
% Data completeness (in resolution range)	92.2 (47.45-2.33) 92.1 (59.67-2.33)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.30 (at 2.34Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.225 , 0.259 0.220 , 0.260	Depositor DCC
$R_{free}$ test set	1276 reflections (3.10%)	DCC
Wilson B-factor (Å <sup>2</sup> )	47.6	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8425	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CIT, 5A8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.21	0/2120	0.35	0/2869
1	B	0.21	0/2160	0.34	0/2917
1	C	0.21	0/2029	0.35	0/2747
1	D	0.21	0/1939	0.34	0/2624
All	All	0.21	0/8248	0.35	0/11157

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	1988	10	0
1	B	2107	0	2058	28	0
1	C	1981	0	1885	9	0
1	D	1894	0	1803	12	0
2	A	6	0	8	0	0
3	A	37	0	0	0	0
3	B	37	0	0	0	0
3	C	37	0	0	1	0
3	D	37	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	13	0	5	3	0
5	A	74	0	0	1	0
5	B	34	0	0	0	0
5	C	53	0	0	0	0
5	D	50	0	0	2	0
All	All	8425	0	7747	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ILE:HD13	1:A:90:LEU:HD22	1.74	0.69
4:B:301:CIT:O4	4:B:301:CIT:O7	2.13	0.67
1:B:38:ILE:HD13	1:B:90:LEU:HD22	1.76	0.66
1:A:256:ARG:NH2	5:A:402:HOH:O	2.31	0.63
1:A:125:THR:OG1	1:A:220:SER:OG	2.19	0.59
1:B:106:THR:HB	1:B:235:ARG:HB2	1.86	0.58
1:C:125:THR:OG1	1:C:220:SER:OG	2.21	0.56
1:D:256:ARG:NH2	5:D:403:HOH:O	2.38	0.56
1:B:125:THR:OG1	1:B:220:SER:OG	2.25	0.54
1:D:160:THR:HG21	1:D:172:MET:HG2	1.88	0.54
1:B:182:THR:HG22	1:B:198:PRO:HB3	1.90	0.53
1:C:24:PHE:HB3	1:C:29:MET:HB2	1.91	0.53
1:B:249[A]:ARG:HD3	1:B:250:SER:N	2.24	0.53
1:C:5:ILE:HG12	1:C:40:ILE:HB	1.90	0.52
1:C:77:ASP:N	1:C:77:ASP:OD2	2.43	0.51
1:B:138:VAL:HB	1:B:208:SER:HB3	1.93	0.51
1:C:223:HIS:O	1:C:223:HIS:ND1	2.44	0.50
1:C:172:MET:HE1	1:C:178:ALA:HB3	1.95	0.48
1:A:197:SER:OG	1:B:165[B]:LYS:NZ	2.44	0.48
1:B:209:LEU:HB2	1:B:234:ILE:HB	1.95	0.47
1:B:78:TRP:CD2	1:B:86:LEU:HD21	2.51	0.46
1:D:2:LYS:N	5:D:405:HOH:O	2.48	0.46
1:C:123:GLU:OE2	3:C:301:5A8:OAQ	2.34	0.45
1:D:125:THR:OG1	1:D:220:SER:OG	2.21	0.45
1:A:119:LEU:HD11	1:A:241:LYS:HD2	1.99	0.44
1:B:210:GLN:HA	1:B:211:PRO:HD3	1.84	0.44
1:B:78:TRP:CE2	1:B:86:LEU:HD21	2.52	0.44
1:D:68:HIS:O	1:D:80:PRO:HD3	2.18	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:GLN:HA	1:A:211:PRO:HD3	1.83	0.43
1:B:95:TYR:HB3	1:B:248:PHE:CE1	2.53	0.43
1:D:210:GLN:HA	1:D:211:PRO:HD3	1.85	0.43
1:A:5:ILE:HG12	1:A:40:ILE:HB	2.01	0.43
1:B:260:SER:O	1:D:189:ASN:HB2	2.19	0.43
1:B:253:PHE:H	4:B:301:CIT:H42	1.83	0.43
1:A:133:PHE:CD2	1:A:151:GLY:HA2	2.53	0.43
1:B:89:LEU:HB3	1:B:248:PHE:CE1	2.54	0.43
1:D:133:PHE:CD2	1:D:151:GLY:HA2	2.54	0.43
1:B:133:PHE:CD2	1:B:151:GLY:HA2	2.54	0.42
1:B:128:SER:OG	1:B:131:GLY:O	2.24	0.42
4:B:301:CIT:O7	4:B:301:CIT:O2	2.33	0.42
1:C:133:PHE:CD2	1:C:151:GLY:HA2	2.55	0.42
1:B:120:ALA:HB2	1:B:221:VAL:HG23	2.01	0.42
1:D:5:ILE:HD12	1:D:21:ILE:HD12	2.02	0.42
3:D:301:5A8:NAY	3:D:301:5A8:CBE	2.77	0.42
1:B:49:LEU:HD23	1:B:121:LEU:HD23	2.00	0.42
1:D:49:LEU:HD23	1:D:121:LEU:HD23	2.00	0.42
1:A:146:ARG:HG2	1:A:192:TYR:HD1	1.85	0.41
1:B:173:HIS:HA	1:B:174:PRO:HD3	1.87	0.41
1:B:24:PHE:CD1	1:B:29:MET:HG3	2.55	0.41
1:D:182:THR:HG22	1:D:198:PRO:HB3	2.01	0.41
1:B:266:GLU:HG3	1:B:267:HIS:ND1	2.36	0.41
1:A:77:ASP:OD1	1:A:77:ASP:N	2.50	0.41
1:B:119:LEU:HD11	1:B:241:LYS:HD2	2.03	0.41
1:B:101:PRO:HG3	1:B:177:GLU:HG2	2.03	0.41
1:D:146:ARG:HG2	1:D:192:TYR:HD1	1.85	0.41
1:B:160:THR:HA	1:B:164:ASN:HB3	2.04	0.40
1:B:264:ASP:OD1	1:B:267:HIS:ND1	2.54	0.40
1:B:77:ASP:OD1	1:B:77:ASP:N	2.54	0.40
1:C:210:GLN:HA	1:C:211:PRO:HD3	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/272 (96%)	246 (95%)	14 (5%)	0	100	100
1	B	258/272 (95%)	245 (95%)	13 (5%)	0	100	100
1	C	246/272 (90%)	231 (94%)	15 (6%)	0	100	100
1	D	236/272 (87%)	220 (93%)	16 (7%)	0	100	100
All	All	1000/1088 (92%)	942 (94%)	58 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/237 (93%)	217 (99%)	3 (1%)	71	82
1	B	228/237 (96%)	221 (97%)	7 (3%)	45	56
1	C	208/237 (88%)	205 (99%)	3 (1%)	71	82
1	D	199/237 (84%)	193 (97%)	6 (3%)	46	57
All	All	855/948 (90%)	836 (98%)	19 (2%)	59	70

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	LEU
1	A	163	TYR
1	A	251	PHE
1	B	123	GLU
1	B	163	TYR
1	B	223	HIS
1	B	226	ILE
1	B	249[A]	ARG
1	B	249[B]	ARG
1	B	251	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	123	GLU
1	C	163	TYR
1	C	251	PHE
1	D	29	MET
1	D	71	HIS
1	D	79	ARG
1	D	123	GLU
1	D	163	TYR
1	D	251	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	301	-	5,5,5	0.35	0	5,5,5	0.23	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	5A8	A	302	-	37,41,41	2.33	13 (35%)	32,58,58	1.96	5 (15%)
4	CIT	B	301	-	3,12,12	1.34	0	3,17,17	2.00	2 (66%)
3	5A8	B	302	-	37,41,41	2.24	16 (43%)	32,58,58	2.37	6 (18%)
3	5A8	C	301	-	37,41,41	2.36	15 (40%)	32,58,58	2.28	5 (15%)
3	5A8	D	301	-	37,41,41	1.85	9 (24%)	32,58,58	2.09	6 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	301	-	-	0/4/4/4	0/0/0/0
3	5A8	A	302	-	-	0/13/35/35	0/5/5/5
4	CIT	B	301	-	-	0/6/16/16	0/0/0/0
3	5A8	B	302	-	-	0/13/35/35	0/5/5/5
3	5A8	C	301	-	-	0/13/35/35	0/5/5/5
3	5A8	D	301	-	-	0/13/35/35	0/5/5/5

All (53) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	5A8	C5-N7	-6.61	1.30	1.38
3	A	302	5A8	C8-N9	-6.42	1.29	1.36
3	C	301	5A8	C5-N7	-6.29	1.30	1.38
3	B	302	5A8	C5-N7	-6.15	1.30	1.38
3	C	301	5A8	C8-N9	-5.80	1.30	1.36
3	D	301	5A8	C5-N7	-5.76	1.31	1.38
3	B	302	5A8	C8-N9	-5.17	1.31	1.36
3	C	301	5A8	OAM-CAN	-3.84	1.36	1.45
3	C	301	5A8	C8-SAA	-3.70	1.65	1.75
3	C	301	5A8	C4-N3	-3.68	1.30	1.35
3	D	301	5A8	C8-N9	-3.66	1.32	1.36
3	B	302	5A8	OAM-CAN	-3.51	1.37	1.45
3	A	302	5A8	C4-N3	-3.41	1.30	1.35
3	B	302	5A8	OAM-CAL	-3.29	1.36	1.41
3	A	302	5A8	C8-SAA	-3.27	1.66	1.75
3	A	302	5A8	CBH-CBG	-3.25	1.36	1.41
3	C	301	5A8	C2-N1	-3.08	1.28	1.33
3	C	301	5A8	CBE-NBD	-3.02	1.30	1.36
3	B	302	5A8	C4-N3	-2.97	1.31	1.35

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	302	5A8	CBG-NBD	-2.95	1.29	1.38
3	A	302	5A8	CBE-NBD	-2.81	1.30	1.36
3	C	301	5A8	CBF-CBG	-2.75	1.35	1.42
3	C	301	5A8	CBG-NBD	-2.73	1.30	1.38
3	B	302	5A8	CBE-NBD	-2.69	1.31	1.36
3	C	301	5A8	CBK-CBF	-2.68	1.36	1.42
3	A	302	5A8	CBK-CBF	-2.67	1.36	1.42
3	A	302	5A8	CBE-CBC	-2.61	1.30	1.37
3	B	302	5A8	CBK-CBF	-2.58	1.36	1.42
3	C	301	5A8	C5-C4	-2.56	1.34	1.40
3	B	302	5A8	CAP-CAL	-2.55	1.49	1.53
3	C	301	5A8	CBE-CBC	-2.55	1.30	1.37
3	A	302	5A8	CBF-CBG	-2.54	1.35	1.42
3	A	302	5A8	OAM-CAN	-2.50	1.39	1.45
3	B	302	5A8	CBF-CBG	-2.50	1.35	1.42
3	B	302	5A8	CBG-NBD	-2.45	1.31	1.38
3	B	302	5A8	CBE-CBC	-2.43	1.30	1.37
3	B	302	5A8	CBH-CBG	-2.42	1.37	1.41
3	D	301	5A8	C4-N3	-2.39	1.32	1.35
3	A	302	5A8	C5-C4	-2.29	1.35	1.40
3	B	302	5A8	C5-C4	-2.27	1.35	1.40
3	C	301	5A8	CBH-CBG	-2.25	1.37	1.41
3	D	301	5A8	CBG-NBD	-2.19	1.31	1.38
3	C	301	5A8	OAM-CAL	-2.12	1.38	1.41
3	D	301	5A8	CBE-NBD	-2.10	1.32	1.36
3	B	302	5A8	CAO-CAP	-2.06	1.48	1.53
3	B	302	5A8	C2-N1	-2.04	1.30	1.33
3	D	301	5A8	CBH-CBG	-2.02	1.38	1.41
3	D	301	5A8	CBK-CBF	-2.02	1.38	1.42
3	A	302	5A8	NAU-NAT	2.10	1.28	1.23
3	B	302	5A8	NAU-NAT	2.39	1.29	1.23
3	D	301	5A8	CAS-CAN	2.39	1.55	1.51
3	C	301	5A8	NAU-NAT	2.81	1.30	1.23
3	D	301	5A8	NAU-NAT	4.52	1.35	1.23

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	301	5A8	N3-C2-N1	-9.92	120.22	128.86
3	B	302	5A8	N3-C2-N1	-9.21	120.84	128.86
3	D	301	5A8	N3-C2-N1	-7.35	122.46	128.86
3	A	302	5A8	N3-C2-N1	-7.32	122.48	128.86

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	5A8	CAN-OAM-CAL	-6.02	103.36	109.77
3	D	301	5A8	CAN-OAM-CAL	-5.37	104.05	109.77
3	A	302	5A8	OAM-CAL-N9	-4.28	103.11	108.18
3	C	301	5A8	CAN-OAM-CAL	-4.26	105.23	109.77
3	A	302	5A8	CAN-OAM-CAL	-4.13	105.37	109.77
3	A	302	5A8	C4-C5-N7	-2.87	106.78	109.47
3	D	301	5A8	OAM-CAN-CAO	-2.72	99.76	105.17
4	B	301	CIT	C3-C2-C1	-2.63	110.84	114.95
3	B	302	5A8	OAR-CAO-CAP	-2.57	103.59	111.83
3	B	302	5A8	CAL-N9-C4	-2.43	122.45	126.71
3	B	302	5A8	OAQ-CAP-CAO	-2.38	104.20	111.83
4	B	301	CIT	C3-C4-C5	-2.18	111.55	114.95
3	D	301	5A8	CBA-CAZ-NAY	-2.11	106.91	111.91
3	D	301	5A8	C4-C5-N7	-2.06	107.54	109.47
3	C	301	5A8	CBJ-CBI-CBH	-2.00	117.59	120.45
3	C	301	5A8	C2-N1-C6	2.08	122.40	118.77
3	A	302	5A8	CAS-NAT-NAU	2.38	119.39	115.30
3	B	302	5A8	CAS-NAT-NAU	4.00	122.19	115.30
3	D	301	5A8	CAS-NAT-NAU	4.42	122.90	115.30
3	C	301	5A8	CAS-NAT-NAU	4.55	123.12	115.30

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	301	CIT	3	0
3	C	301	5A8	1	0
3	D	301	5A8	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	263/272 (96%)	0.41	7 (2%) 55 64	27, 52, 90, 134	46 (17%)
1	B	262/272 (96%)	0.64	15 (5%) 24 35	31, 70, 106, 131	52 (19%)
1	C	254/272 (93%)	0.70	22 (8%) 11 16	33, 70, 121, 136	40 (15%)
1	D	244/272 (89%)	0.74	19 (7%) 14 20	35, 72, 122, 156	37 (15%)
All	All	1023/1088 (94%)	0.62	63 (6%) 21 30	27, 66, 117, 156	175 (17%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	248	PHE	6.5
1	A	264	ASP	5.3
1	C	40	ILE	4.9
1	A	111	ILE	4.4
1	B	20	MET	4.4
1	C	3	TYR	4.3
1	C	212	VAL	4.2
1	C	19	ASN	4.2
1	D	119	LEU	4.2
1	C	65	ILE	3.9
1	C	243	ILE	3.9
1	B	217	PHE	3.8
1	D	247	ARG	3.8
1	D	34	VAL	3.5
1	B	18	LEU	3.5
1	B	87	VAL	3.5
1	C	24	PHE	3.4
1	C	76	ALA	3.4
1	C	248	PHE	3.3
1	A	78	TRP	3.2
1	A	263	GLU	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	248	PHE	3.1
1	B	216	ASP	3.0
1	B	230	ASP	2.9
1	D	217	PHE	2.9
1	D	70	GLY	2.9
1	D	83	ALA	2.8
1	D	71	HIS	2.8
1	D	84	ASP	2.8
1	A	110	GLY	2.8
1	D	130	GLY	2.7
1	C	49	LEU	2.7
1	D	15	LEU	2.7
1	C	94	GLU	2.7
1	B	229	ARG	2.6
1	B	34	VAL	2.6
1	B	187	ILE	2.6
1	C	81	ALA	2.4
1	D	227	LEU	2.4
1	D	117	THR	2.4
1	C	6	THR	2.4
1	B	95	TYR	2.4
1	C	31	TYR	2.3
1	D	20	MET	2.3
1	C	66	GLY	2.2
1	A	71	HIS	2.2
1	C	78	TRP	2.2
1	B	156	THR	2.2
1	D	226	ILE	2.2
1	C	15	LEU	2.2
1	C	59	LEU	2.2
1	B	96	GLN	2.2
1	C	84	ASP	2.2
1	D	32	ASP	2.1
1	C	13	SER	2.1
1	D	78	TRP	2.1
1	B	109	TYR	2.1
1	B	94	GLU	2.1
1	B	6	THR	2.1
1	C	70	GLY	2.1
1	C	116	ALA	2.1
1	D	16	LEU	2.0
1	D	61	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	CIT	B	301	13/13	0.80	0.34	7.89	57,60,65,65	13
2	GOL	A	301	6/6	0.84	0.32	6.03	81,81,81,82	0
3	5A8	D	301	37/37	0.83	0.24	1.47	51,60,68,77	4
3	5A8	B	302	37/37	0.89	0.15	-0.14	50,57,64,67	5
3	5A8	A	302	37/37	0.93	0.15	-0.27	33,44,55,57	5
3	5A8	C	301	37/37	0.92	0.15	-0.51	38,44,59,59	5

## 6.5 Other polymers [i](#)

There are no such residues in this entry.