



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 02:02 pm GMT

PDB ID : 5DKG
Title : Crystal Structure of the ER-alpha Ligand-binding Domain in complex with a t-butyl-substituted, methyl, triaryl-ethylene derivative 4,4'-[2-(4-tert-butylphenyl)prop-1-ene-1,1-diyl]diphenol
Authors : Nwachukwu, J.C.; Srinivasan, S.; Zheng, Y.; Wang, S.; Min, J.; Dong, C.; Liao, Z.; Cavett, V.; Nowak, J.; Houtman, R.; Carlson, K.E.; Josan, J.S.; Elemento, O.; Katzenellenbogen, J.A.; Zhou, H.B.; Nettles, K.W.
Deposited on : 2015-09-03
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	1170 (2.16-2.16)
Clashscore	112137	1278 (2.16-2.16)
Ramachandran outliers	110173	1256 (2.16-2.16)
Sidechain outliers	110143	1255 (2.16-2.16)
RSRZ outliers	101464	1175 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	<div> <div>2%</div> <div> <div></div> <div>80%</div> <div>9%</div> <div>10%</div> </div> </div>
1	B	257	<div> <div>4%</div> <div> <div></div> <div>78%</div> <div>14%</div> <div>7%</div> </div> </div>
2	C	14	<div> <div></div> <div> <div>57%</div> <div>7%</div> <div>7%</div> <div>29%</div> </div> </div>
2	D	14	<div> <div></div> <div> <div>50%</div> <div>14%</div> <div>7%</div> <div>29%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4117 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estrogen receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	2	0
			1831	1173	312	327	19			
1	B	240	Total	C	N	O	S	0	1	0
			1870	1191	321	340	18			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	537	SER	TYR	engineered mutation	UNP P03372
B	537	SER	TYR	engineered mutation	UNP P03372

- Molecule 2 is a protein called Nuclear receptor coactivator 2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	0	0	0
			89	57	19	13			
2	D	10	Total	C	N	O	0	0	0
			84	54	17	13			

- Molecule 3 is 4,4'-[2-(4-tert-butylphenyl)prop-1-ene-1,1-diyl]diphenol (three-letter code: 5C7) (formula: C₂₅H₂₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			27	25	2		
3	B	1	Total	C	O	0	0
			27	25	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	90	Total	O	0	0
			90	90		
4	B	94	Total	O	0	0
			94	94		
4	C	1	Total	O	0	0
			1	1		
4	D	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Estrogen receptor



- Molecule 1: Estrogen receptor



- Molecule 2: Nuclear receptor coactivator 2



- Molecule 2: Nuclear receptor coactivator 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.34Å 82.08Å 58.62Å 90.00° 110.83° 90.00°	Depositor
Resolution (Å)	45.57 – 2.15 45.57 – 2.15	Depositor EDS
% Data completeness (in resolution range)	93.3 (45.57-2.15) 93.1 (45.57-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.34 (at 2.14Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.194 , 0.233 0.195 , 0.232	Depositor DCC
R_{free} test set	1875 reflections (7.37%)	DCC
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.451	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.029 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4117	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5C7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.21	0/1869	0.37	0/2528
1	B	0.25	0/1902	0.41	0/2576
2	C	0.22	0/90	0.63	0/119
2	D	0.24	0/84	0.54	0/111
All	All	0.23	0/3945	0.40	0/5334

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	464	SER	Peptide
1	B	466	LEU	Peptide
1	B	467	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1831	0	1845	25	0
1	B	1870	0	1872	39	0
2	C	89	0	95	1	0
2	D	84	0	90	2	0
3	A	27	0	0	0	0
3	B	27	0	0	0	0
4	A	90	0	0	10	0
4	B	94	0	0	5	0
4	C	1	0	0	0	0
4	D	4	0	0	0	0
All	All	4117	0	3902	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:464:SER:N	1:B:468:SER:HG	0.93	1.41
1:B:464:SER:N	1:B:468:SER:OG	1.76	1.16
1:B:464:SER:HB2	1:B:465:THR:CG2	1.98	0.94
1:A:513[B]:HIS:ND1	4:A:701:HOH:O	1.99	0.91
1:B:464:SER:HB2	1:B:465:THR:HB	1.63	0.80
1:B:465:THR:HG22	1:B:468:SER:HA	1.63	0.79
1:B:464:SER:HB2	1:B:465:THR:CB	2.13	0.77
1:B:464:SER:CB	1:B:465:THR:HB	2.15	0.76
4:A:781:HOH:O	1:B:381:CYS:SG	2.43	0.75
1:A:336:PRO:O	4:A:702:HOH:O	2.04	0.73
1:A:513[A]:HIS:ND1	4:A:701:HOH:O	2.23	0.71
1:A:339:GLU:HG3	1:A:418:VAL:HG22	1.73	0.69
1:A:413:ASN:ND2	4:A:705:HOH:O	2.25	0.69
1:B:464:SER:HB2	1:B:465:THR:HG22	1.74	0.68
1:A:394:ARG:NH1	1:A:404:PHE:O	2.26	0.68
1:B:464:SER:N	1:B:468:SER:CB	2.58	0.66
1:A:548:ARG:O	4:A:703:HOH:O	2.12	0.66
1:A:473:ASP:OD2	1:A:477:ARG:NH1	2.28	0.66
1:B:527:SER:O	4:B:701:HOH:O	2.14	0.66
2:D:687:HIS:O	2:D:688:LYS:HB2	1.97	0.65
1:B:426:ASP:OD2	4:B:702:HOH:O	2.14	0.64
1:B:465:THR:CG2	1:B:468:SER:HA	2.28	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:513[B]:HIS:CE1	1:B:515:ARG:HH22	2.18	0.61
1:A:359:ASN:ND2	4:A:706:HOH:O	2.29	0.58
2:C:691:HIS:O	2:C:695:GLN:HG2	2.04	0.56
1:B:307:ALA:HA	1:B:310:LEU:HD12	1.87	0.56
1:B:460:THR:HG22	4:B:751:HOH:O	2.06	0.55
1:B:526:TYR:O	1:B:548:ARG:NH2	2.41	0.54
1:A:530:CYS:O	4:A:704:HOH:O	2.18	0.53
1:B:498:GLN:HA	1:B:501:HIS:CE1	2.43	0.53
1:A:498:GLN:HA	1:A:501:HIS:CE1	2.43	0.53
1:A:424:ILE:HA	1:A:427:MET:HE3	1.91	0.52
1:B:377:HIS:NE2	1:B:460:THR:HG23	2.25	0.52
1:A:396:MET:O	1:A:436[B]:ARG:HD3	2.11	0.51
1:B:529:LYS:O	1:B:532:ASN:N	2.44	0.50
1:B:321:ASP:OD2	1:B:363:ARG:NH2	2.46	0.49
1:A:394:ARG:NH2	4:A:713:HOH:O	2.45	0.49
1:B:539:LEU:O	1:B:543:MET:HG2	2.13	0.48
1:B:533:VAL:HG23	1:B:534:VAL:N	2.27	0.48
1:B:305:SER:N	4:B:716:HOH:O	2.46	0.48
1:A:424:ILE:HD12	1:A:427:MET:HE1	1.96	0.47
1:A:508:LEU:O	1:A:511:LEU:HB2	2.14	0.47
1:A:304:ASN:ND2	4:A:710:HOH:O	2.42	0.47
1:A:377:HIS:NE2	1:A:460:THR:HB	2.30	0.46
1:B:487:ILE:HD11	1:B:504:LEU:HD22	1.97	0.46
1:B:464:SER:CB	1:B:465:THR:CB	2.85	0.46
1:A:513[B]:HIS:CE1	1:B:459:TYR:HD2	2.33	0.46
1:B:523:GLU:HA	1:B:526:TYR:CZ	2.51	0.46
1:B:526:TYR:HB2	1:B:548:ARG:NH2	2.30	0.45
1:B:501:HIS:HB3	4:B:779:HOH:O	2.15	0.45
1:B:536:LEU:HB2	1:B:541:LEU:HD13	1.98	0.45
1:B:398:HIS:HB3	1:B:401:LYS:HB2	1.98	0.45
1:A:377:HIS:CE1	1:A:460:THR:HB	2.52	0.44
1:B:347:THR:HG23	1:B:540:LEU:HD23	1.99	0.43
1:A:513[B]:HIS:ND1	1:B:459:TYR:HD2	2.18	0.41
1:B:465:THR:OG1	1:B:466:LEU:N	2.53	0.41
1:B:376:VAL:HG13	2:D:690:LEU:HD23	2.03	0.41
1:A:513[B]:HIS:CE1	1:B:515:ARG:NH2	2.86	0.41
1:B:413:ASN:OD1	1:B:416:LYS:NZ	2.54	0.41
1:B:464:SER:N	1:B:465:THR:HG22	2.36	0.41
1:A:396:MET:O	1:A:436[A]:ARG:HD2	2.22	0.40
1:A:528:MET:O	1:A:530:CYS:N	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/257 (88%)	222 (98%)	4 (2%)	1 (0%)	38	32
1	B	235/257 (91%)	230 (98%)	3 (1%)	2 (1%)	20	12
2	C	8/14 (57%)	7 (88%)	0	1 (12%)	0	0
2	D	8/14 (57%)	7 (88%)	0	1 (12%)	0	0
All	All	478/542 (88%)	466 (98%)	7 (2%)	5 (1%)	17	11

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	305	SER
2	C	691	HIS
2	D	688	LYS
1	B	465	THR
1	B	533	VAL

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/232 (88%)	201 (99%)	2 (1%)	80	84
1	B	205/232 (88%)	205 (100%)	0	100	100
2	C	10/14 (71%)	10 (100%)	0	100	100
2	D	9/14 (64%)	8 (89%)	1 (11%)	7	3
All	All	427/492 (87%)	424 (99%)	3 (1%)	87	91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	413	ASN
1	A	460	THR
2	D	688	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	304	ASN
1	A	524	HIS
1	B	455	ASN
2	C	691	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	5C7	A	601	-	29,29,29	3.17	15 (51%)	41,42,42	0.64	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	5C7	B	601	-	29,29,29	3.13	15 (51%)	41,42,42	0.62	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5C7	A	601	-	-	0/22/22/22	0/3/3/3
3	5C7	B	601	-	-	0/22/22/22	0/3/3/3

All (30) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	5C7	C24-C22	-2.70	1.44	1.53
3	A	601	5C7	C24-C22	-2.68	1.44	1.53
3	B	601	5C7	C20-C19	-2.67	1.34	1.38
3	A	601	5C7	C20-C19	-2.57	1.34	1.38
3	B	601	5C7	O15-C14	-2.39	1.31	1.37
3	A	601	5C7	C01-C02	-2.38	1.46	1.50
3	B	601	5C7	C01-C02	-2.36	1.46	1.50
3	B	601	5C7	C10-C09	-2.33	1.34	1.38
3	A	601	5C7	O15-C14	-2.32	1.31	1.37
3	B	601	5C7	C13-C14	-2.25	1.34	1.38
3	A	601	5C7	C13-C14	-2.25	1.34	1.38
3	A	601	5C7	C10-C09	-2.21	1.34	1.38
3	A	601	5C7	C17-C16	2.79	1.43	1.38
3	A	601	5C7	C18-C02	2.88	1.54	1.49
3	B	601	5C7	C17-C16	2.88	1.43	1.38
3	B	601	5C7	C18-C02	2.90	1.54	1.49
3	A	601	5C7	C19-C18	2.96	1.44	1.39
3	B	601	5C7	C11-C03	2.96	1.54	1.49
3	B	601	5C7	C19-C18	2.98	1.44	1.39
3	A	601	5C7	C11-C03	2.99	1.54	1.49
3	B	601	5C7	C12-C13	3.52	1.45	1.38
3	A	601	5C7	C12-C13	3.55	1.45	1.38
3	A	601	5C7	C22-C21	3.64	1.61	1.53
3	B	601	5C7	C22-C21	3.67	1.61	1.53
3	B	601	5C7	C04-C03	3.80	1.55	1.49
3	A	601	5C7	C04-C03	4.00	1.55	1.49
3	B	601	5C7	C20-C21	5.31	1.47	1.39
3	A	601	5C7	C20-C21	5.32	1.47	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	601	5C7	C03-C02	11.43	1.42	1.35
3	A	601	5C7	C03-C02	11.71	1.42	1.35

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	231/257 (89%)	-0.13	5 (2%) 62 69	28, 45, 75, 107	0
1	B	240/257 (93%)	-0.06	9 (3%) 41 48	28, 47, 88, 120	0
2	C	10/14 (71%)	0.46	0 100 100	51, 67, 80, 92	0
2	D	10/14 (71%)	-0.07	0 100 100	40, 60, 80, 82	0
All	All	491/542 (90%)	-0.08	14 (2%) 52 60	28, 47, 82, 120	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	549	LEU	7.1
1	B	535	PRO	3.5
1	A	530	CYS	3.5
1	A	336	PRO	2.8
1	A	333	PRO	2.5
1	B	467	LYS	2.3
1	A	305	SER	2.3
1	B	342	MET	2.3
1	B	468	SER	2.3
1	B	532	ASN	2.2
1	B	526	TYR	2.1
1	B	459	TYR	2.1
1	B	464	SER	2.1
1	A	461	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	5C7	B	601	27/27	0.95	0.12	-0.08	29,36,43,44	0
3	5C7	A	601	27/27	0.97	0.12	-0.22	30,35,41,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.