



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:34 am GMT

PDB ID : 4DL3
Title : Human DNA polymerase eta inserting dCMPNPP opposite GG template (GG0b).
Authors : Zhao, Y.; Biertumpfel, C.; Gregory, M.; Hua, Y.; Hanaoka, F.; Yang, W.
Deposited on : 2012-02-05
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix) : 1.9-1692
EDS : trunk28620
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : recalc28949


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	100719	4243 (2.10-2.10)
Clashscore	112137	4788 (2.10-2.10)
Ramachandran outliers	110173	4740 (2.10-2.10)
Sidechain outliers	110143	4741 (2.10-2.10)
RSRZ outliers	101464	4275 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	<div> <div>10%</div> <div>68%</div> <div>27%</div> <div>• •</div> </div>
2	T	12	<div> <div>25%</div> <div>50%</div> <div>50%</div> </div>
3	P	9	<div> <div>89%</div> <div>11%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	504	-	-	-	X
5	GOL	A	505	-	-	-	X

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4138 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	4	0
			3380	2120	606	628	26			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q9Y253
A	-1	PRO	-	EXPRESSION TAG	UNP Q9Y253
A	0	HIS	-	EXPRESSION TAG	UNP Q9Y253

- Molecule 2 is a DNA chain called DNA (5'-D(*TP*AP*CP*GP*GP*TP*CP*AP*CP*AP*CP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	12	Total	C	N	O	P	0	2	1
			265	125	49	78	13			


- Molecule 3 is a DNA chain called DNA (5'-D(*TP*AP*GP*TP*GP*TP*GP*AP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	9	Total	C	N	O	P	0	0	1
			168	79	32	49	8			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total 6	C 3	O 3	0	0
5	A	1	Total 6	C 3	O 3	0	0
5	A	1	Total 6	C 3	O 3	0	0

- Molecule 6 is 2'-DEOXY-5'-O-[(R)-HYDROXY{[(R)-HYDROXY(PHOSPHONOOXY) PHOSPHORYL]AMINO}PHOSPHORYL]CYTIDINE (three-letter code: 0KX) (formula: $C_9H_{17}N_4O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	P	1	Total	C	N	O	P	0	0
			28	9	4	12	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	245	Total	O	0	0
			245	245		
7	T	16	Total	O	0	0
			16	16		
7	P	16	Total	O	0	0
			16	16		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase eta



- Molecule 2: DNA (5'-D(*TP*AP*CP*GP*GP*TP*CP*AP*CP*AP*CP*T)-3')



- Molecule 3: DNA (5'-D(*TP*AP*GP*TP*GP*TP*GP*AP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	97.38Å 97.38Å 81.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.10 19.47 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-2.10) 98.6 (19.47-2.10)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.09Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7.1_743)	Depositor
R, R_{free}	0.204 , 0.217 0.198 , 0.213	Depositor DCC
R_{free} test set	1951 reflections (7.74%)	DCC
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.048 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4138	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OKX, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3451	0.43	0/4657
2	T	0.51	0/296	0.87	0/455
3	P	0.53	0/188	0.97	0/290
All	All	0.30	0/3935	0.52	0/5402

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3380	0	3429	125	0
2	T	265	0	146	20	0
3	P	168	0	90	1	0
4	A	2	0	0	0	0
5	A	18	0	24	3	0
6	P	28	0	15	3	0
7	A	245	0	0	4	0
7	P	16	0	0	0	0
7	T	16	0	0	0	0
All	All	4138	0	3704	136	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (136) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:3[B]:DC:H4'	2:T:4[B]:DG:OP1	1.38	1.10
2:T:5:DG:H2''	2:T:6:DT:H5''	1.40	1.01
1:A:61:ARG:CG	2:T:3[B]:DC:H41	1.73	0.99
1:A:61:ARG:HG2	2:T:3[B]:DC:H41	1.27	0.96
1:A:61:ARG:HG3	2:T:3[B]:DC:N4	1.90	0.86
1:A:61:ARG:CG	2:T:3[B]:DC:N4	2.41	0.84
2:T:5:DG:C2'	2:T:6:DT:H5''	2.09	0.81
1:A:136:PRO:HA	1:A:180:ILE:HG21	1.63	0.79
1:A:132:LEU:O	1:A:132:LEU:HD12	1.84	0.78
1:A:133:GLN:OE1	1:A:133:GLN:HA	1.82	0.78
1:A:371:ARG:HA	1:A:380:SER:HB3	1.64	0.78
1:A:62:SER:HB2	2:T:3[B]:DC:C5	2.20	0.77
1:A:371:ARG:HG2	1:A:420:THR:OG1	1.84	0.77
1:A:160:THR:HG22	1:A:169:GLN:NE2	2.00	0.76
1:A:378:LEU:HD22	1:A:378:LEU:H	1.50	0.76
1:A:62:SER:HA	2:T:3[B]:DC:C4	2.22	0.75
1:A:376:LYS:HD3	1:A:376:LYS:O	1.90	0.71
1:A:261:LYS:NZ	3:P:6:DT:OP1	2.22	0.70
1:A:146:TYR:OH	1:A:163:LYS:HD3	1.93	0.68
1:A:369:SER:HB3	1:A:423:PHE:HB3	1.76	0.67
1:A:128:ARG:HH22	1:A:199:GLU:CD	1.97	0.66
1:A:209:GLY:O	5:A:504:GOL:H32	1.95	0.65
1:A:375:ASP:OD2	1:A:377:ARG:NH2	2.29	0.65
1:A:382:ARG:CG	1:A:382:ARG:HH11	2.10	0.64
1:A:122:THR:O	1:A:126:GLN:HG2	1.96	0.64
1:A:132:LEU:HB2	1:A:135:GLN:HB3	1.80	0.62
1:A:409:SER:HB3	1:A:416:SER:OG	1.99	0.62
1:A:40:LYS:HD3	1:A:79:GLN:HE21	1.63	0.62
2:T:3[B]:DC:C4'	2:T:4[B]:DG:OP1	2.30	0.62
1:A:40:LYS:HD3	1:A:79:GLN:NE2	2.15	0.62
1:A:367:VAL:CG1	1:A:382:ARG:HD2	2.30	0.62
1:A:160:THR:CG2	1:A:169:GLN:NE2	2.64	0.61
1:A:372:VAL:HB	1:A:417:PRO:HG2	1.82	0.61
1:A:371:ARG:HD3	1:A:377:ARG:O	2.01	0.61
1:A:128:ARG:NH2	1:A:199:GLU:OE2	2.32	0.61
1:A:371:ARG:CD	1:A:377:ARG:O	2.48	0.61
1:A:61:ARG:O	1:A:62:SER:HB3	2.01	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:7:DC:H2''	2:T:8:DA:H5'	1.83	0.60
1:A:322:SER:HB3	1:A:423:PHE:CD1	2.37	0.60
1:A:382:ARG:HH11	1:A:382:ARG:HG3	1.64	0.59
2:T:5:DG:H2''	2:T:6:DT:C5'	2.23	0.56
1:A:24:ARG:HB2	5:A:504:GOL:H31	1.87	0.56
1:A:317:LYS:HG3	2:T:8:DA:OP1	2.05	0.56
1:A:126:GLN:O	1:A:130:GLN:HG3	2.06	0.55
1:A:383:ARG:HB3	1:A:402:VAL:CG1	2.37	0.54
1:A:367:VAL:HG13	1:A:382:ARG:HD2	1.88	0.54
1:A:62:SER:CA	2:T:3[B]:DC:C4	2.91	0.54
1:A:371:ARG:NH1	1:A:375:ASP:O	2.42	0.53
1:A:383:ARG:CB	1:A:402:VAL:CG1	2.86	0.53
1:A:14:MET:HG2	1:A:115:ASP:O	2.09	0.52
1:A:198:GLU:HG2	1:A:235:GLN:NE2	2.25	0.52
1:A:230:ASN:O	1:A:234:ARG:HB2	2.09	0.52
1:A:-2:GLY:N	1:A:-1:PRO:HD2	2.25	0.52
1:A:277:GLU:O	1:A:280:GLN:HG2	2.08	0.52
1:A:413:THR:O	1:A:413:THR:CG2	2.58	0.51
1:A:61:ARG:HG2	2:T:3[B]:DC:N4	2.10	0.51
1:A:2:ALA:HB1	7:A:777:HOH:O	2.09	0.51
1:A:339:TRP:O	1:A:343:GLN:HG2	2.10	0.51
1:A:382:ARG:NH1	1:A:382:ARG:CG	2.70	0.51
1:A:290:PHE:O	1:A:294:ASN:HB2	2.11	0.51
1:A:62:SER:N	2:T:3[B]:DC:N4	2.58	0.50
1:A:266:VAL:HA	1:A:290:PHE:CZ	2.47	0.50
1:A:378:LEU:N	1:A:378:LEU:HD13	2.27	0.49
1:A:181:ASP:OD1	1:A:182:ASN:N	2.45	0.49
1:A:411:ILE:O	1:A:412:GLN:HB3	2.12	0.49
1:A:365:GLN:OE1	1:A:428:LYS:HE2	2.12	0.49
1:A:141:LEU:HA	7:A:808:HOH:O	2.13	0.49
1:A:147:ILE:HG21	1:A:150:LEU:HD12	1.93	0.49
1:A:342:LEU:O	1:A:346:GLN:HG3	2.13	0.49
1:A:11:LEU:HD13	1:A:11:LEU:C	2.34	0.49
1:A:152:GLN:HG2	1:A:153:GLY:N	2.26	0.49
1:A:110:GLU:HB2	1:A:311:LYS:O	2.12	0.49
1:A:370:ILE:HG22	1:A:422:LEU:HD23	1.95	0.49
6:P:10:0KX:H7	6:P:10:0KX:O2B	2.13	0.48
1:A:376:LYS:C	1:A:376:LYS:HD3	2.33	0.48
1:A:413:THR:O	1:A:413:THR:HG23	2.14	0.48
1:A:368:VAL:HG21	1:A:399:ALA:HA	1.96	0.48
1:A:61:ARG:O	1:A:62:SER:CB	2.62	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:LEU:O	1:A:266:VAL:HG23	2.15	0.47
1:A:397:HIS:C	1:A:397:HIS:HD1	2.17	0.47
1:A:51:SER:O	1:A:55:ARG:HG3	2.14	0.47
1:A:381:LEU:HD11	1:A:383:ARG:NE	2.30	0.47
1:A:5:GLN:HB3	1:A:186:PRO:HB3	1.97	0.47
1:A:201:ARG:HD3	1:A:235:GLN:OE1	2.16	0.46
1:A:160:THR:HG22	1:A:169:GLN:HE22	1.75	0.46
1:A:356:ARG:NH1	7:A:761:HOH:O	2.43	0.46
1:A:16:CYS:HA	6:P:10:0KX:O1B	2.15	0.46
1:A:96:SER:O	1:A:100:MET:HG2	2.16	0.46
1:A:39:TYR:O	1:A:45:GLY:HA2	2.15	0.46
1:A:400:PHE:O	1:A:404:LYS:HB3	2.16	0.46
1:A:198:GLU:HG2	1:A:235:GLN:HE22	1.81	0.46
1:A:363:ALA:HB2	1:A:429:PHE:CE1	2.51	0.46
1:A:108:VAL:HG11	1:A:312:PRO:HB3	1.97	0.45
1:A:308:ASP:OD2	1:A:309:PRO:HD2	2.16	0.45
1:A:28:HIS:ND1	5:A:505:GOL:O3	2.46	0.45
1:A:62:SER:CB	2:T:3[B]:DC:C5	2.95	0.45
1:A:381:LEU:HD22	1:A:406:CYS:SG	2.57	0.44
1:A:133:GLN:C	1:A:135:GLN:N	2.70	0.44
1:A:404:LYS:HA	1:A:415:TRP:HH2	1.82	0.44
1:A:129:LEU:O	1:A:132:LEU:O	2.36	0.44
1:A:133:GLN:OE1	1:A:133:GLN:CA	2.61	0.44
1:A:106:PHE:CG	1:A:200[A]:MET:HG2	2.52	0.43
1:A:122:THR:HG23	7:A:757:HOH:O	2.18	0.43
1:A:381:LEU:HD11	1:A:383:ARG:CG	2.49	0.43
6:P:10:0KX:O5'	6:P:10:0KX:H15	2.18	0.43
1:A:136:PRO:CA	1:A:180:ILE:HG21	2.42	0.43
1:A:381:LEU:CD1	1:A:383:ARG:CZ	2.97	0.43
1:A:325:PHE:CD2	1:A:331:LEU:HD21	2.54	0.43
2:T:7:DC:H2'	2:T:8:DA:C8	2.54	0.43
1:A:180:ILE:H	1:A:180:ILE:HG13	1.43	0.43
1:A:178:LEU:HA	1:A:178:LEU:HD23	1.91	0.42
1:A:381:LEU:HD12	1:A:383:ARG:CZ	2.49	0.42
1:A:142:LEU:HB3	1:A:145:THR:HB	2.00	0.42
1:A:15:ASP:O	1:A:16:CYS:C	2.57	0.42
1:A:255:ILE:CD1	1:A:275:MET:HG3	2.49	0.42
1:A:383:ARG:CB	1:A:402:VAL:HG11	2.49	0.42
1:A:185:SER:HA	1:A:186:PRO:HD3	1.89	0.42
2:T:7:DC:H2''	2:T:8:DA:C5'	2.50	0.42
2:T:3[B]:DC:H1'	2:T:4[B]:DG:C8	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:356:ARG:HD3	1:A:362:VAL:HG12	2.01	0.41
1:A:372:VAL:HG13	1:A:380:SER:CB	2.51	0.41
1:A:256:ARG:O	1:A:257:SER:CB	2.68	0.41
1:A:237:LEU:HD23	1:A:237:LEU:C	2.41	0.41
1:A:322:SER:HB3	1:A:423:PHE:HD1	1.84	0.41
1:A:152:GLN:CG	1:A:153:GLY:N	2.83	0.41
1:A:162:GLN:HA	1:A:162:GLN:OE1	2.21	0.41
1:A:383:ARG:HB2	1:A:402:VAL:CG1	2.51	0.41
1:A:47:ILE:HD11	1:A:68:ALA:CB	2.51	0.41
1:A:334:ARG:HG3	1:A:400:PHE:CZ	2.56	0.41
1:A:133:GLN:O	1:A:134:GLY:C	2.59	0.40
1:A:132:LEU:CD1	1:A:132:LEU:O	2.63	0.40
1:A:182:ASN:OD1	1:A:184:THR:OG1	2.30	0.40
1:A:321[B]:CYS:SG	1:A:344:LEU:HD22	2.62	0.40
1:A:352:LEU:HB3	1:A:390:TYR:CE1	2.57	0.40
1:A:412:GLN:O	1:A:412:GLN:CG	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	430/435 (99%)	423 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	372/372 (100%)	351 (94%)	21 (6%)	25	21

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	SER
1	A	81	ARG
1	A	115	ASP
1	A	133	GLN
1	A	163	LYS
1	A	177	SER
1	A	180	ILE
1	A	183	LEU
1	A	372	VAL
1	A	376	LYS
1	A	378	LEU
1	A	380	SER
1	A	381	LEU
1	A	382	ARG
1	A	397	HIS
1	A	404	LYS
1	A	408	THR
1	A	409	SER
1	A	411	ILE
1	A	412	GLN
1	A	413	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
1	A	169	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	503	-	5,5,5	0.35	0	5,5,5	0.24	0
5	GOL	A	504	-	5,5,5	0.12	0	5,5,5	0.29	0
5	GOL	A	505	-	5,5,5	0.15	0	5,5,5	0.28	0
6	OKX	P	10	4	26,29,29	3.45	9 (34%)	31,45,45	1.58	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	503	-	-	0/4/4/4	0/0/0/0
5	GOL	A	504	-	-	0/4/4/4	0/0/0/0
5	GOL	A	505	-	-	0/4/4/4	0/0/0/0
6	OKX	P	10	4	-	0/14/34/34	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	P	10	OKX	O3'-C3'	-4.65	1.33	1.43
6	P	10	OKX	PG-O3G	-2.67	1.43	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	P	10	0KX	C5'-C4'	-2.52	1.43	1.51
6	P	10	0KX	C2-N3	-2.51	1.33	1.38
6	P	10	0KX	PB-O1B	-2.40	1.50	1.56
6	P	10	0KX	PG-O3B	-2.30	1.56	1.60
6	P	10	0KX	C2'-C3'	-2.14	1.47	1.52
6	P	10	0KX	C3'-C4'	-2.10	1.47	1.53
6	P	10	0KX	PA-O1A	15.09	1.63	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	P	10	0KX	C5-C4-N3	-2.93	118.19	121.68
6	P	10	0KX	C2'-C1'-N1	-2.35	108.68	114.23
6	P	10	0KX	O1A-PA-N3A	-2.26	108.41	111.79
6	P	10	0KX	N4-C4-N3	2.51	120.88	116.64
6	P	10	0KX	O4'-C1'-N1	5.66	117.32	107.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	504	GOL	2	0
5	A	505	GOL	1	0
6	P	10	0KX	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.









5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	430/435 (98%)	0.44	42 (9%)  	14, 24, 52, 71	0
2	T	12/12 (100%)	0.65	3 (25%)  	23, 30, 58, 70	0
3	P	9/9 (100%)	-0.38	0  	22, 28, 37, 44	0
All	All	451/456 (98%)	0.43	45 (9%)  	14, 24, 52, 71	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	412	GLN	7.6
1	A	411	ILE	6.7
1	A	1	MET	5.9
1	A	410	GLY	5.3
1	A	0	HIS	5.2
1	A	131	LYS	4.9
2	T	3[A]	DC	4.8
1	A	328	LYS	4.5
1	A	181	ASP	4.3
1	A	133	GLN	4.1
1	A	-1	PRO	4.0
1	A	153	GLY	3.9
1	A	413	THR	3.8
1	A	408	THR	3.7
1	A	179	GLN	3.6
1	A	393	HIS	3.6
1	A	-2	GLY	3.5
1	A	161	VAL	3.4
1	A	380	SER	3.3
2	T	1	DT	3.3
1	A	376	LYS	3.3
1	A	378	LEU	3.3
1	A	432	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	183	LEU	3.2
1	A	182	ASN	3.2
1	A	134	GLY	3.2
1	A	397	HIS	3.2
1	A	135	GLN	3.1
1	A	2	ALA	3.0
2	T	2	DA	2.9
1	A	417	PRO	2.6
1	A	327	GLY	2.5
1	A	374	GLY	2.5
1	A	36	VAL	2.5
1	A	431	ALA	2.5
1	A	127	GLU	2.4
1	A	332	ALA	2.3
1	A	132	LEU	2.3
1	A	409	SER	2.2
1	A	373	GLN	2.2
1	A	42	TRP	2.2
1	A	190	LEU	2.1
1	A	430	SER	2.1
1	A	35	ALA	2.0
1	A	162	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GOL	A	505	6/6	0.87	0.30	6.67	29,31,33,35	0
5	GOL	A	504	6/6	0.84	0.22	3.66	24,27,31,32	0
5	GOL	A	503	6/6	0.91	0.15	1.96	17,20,22,23	0
4	MG	A	502	1/1	0.83	0.11	-0.45	19,19,19,19	0
6	OKX	P	10	28/28	0.97	0.09	-0.75	14,16,20,21	0
4	MG	A	501	1/1	0.97	0.05	-4.88	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.