



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 05:43 am GMT

PDB ID : 5DL6  
Title : Crystal structure of Acinetobacter baumannii OccAB2  
Authors : Zahn, M.; Basle, A.; van den Berg, B.  
Deposited on : 2015-09-04  
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1586 (2.90-2.90)
Clashscore	112137	1807 (2.90-2.90)
Ramachandran outliers	110173	1768 (2.90-2.90)
Sidechain outliers	110143	1770 (2.90-2.90)
RSRZ outliers	101464	1596 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	413	<div> <div></div> <div>77%</div> <div>15%</div> <div>• •</div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative porin for vanillate trafficking (VanP).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	0	0
			3170	2016	536	610	8			


There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-14	ALA	-	expression tag	UNP A0A0E4HLL9
A	-13	ASN	-	expression tag	UNP A0A0E4HLL9
A	-12	VAL	-	expression tag	UNP A0A0E4HLL9
A	-11	ARG	-	expression tag	UNP A0A0E4HLL9
A	-10	LEU	-	expression tag	UNP A0A0E4HLL9
A	-9	GLN	-	expression tag	UNP A0A0E4HLL9
A	-8	HIS	-	expression tag	UNP A0A0E4HLL9
A	-7	HIS	-	expression tag	UNP A0A0E4HLL9
A	-6	HIS	-	expression tag	UNP A0A0E4HLL9
A	-5	HIS	-	expression tag	UNP A0A0E4HLL9
A	-4	HIS	-	expression tag	UNP A0A0E4HLL9
A	-3	HIS	-	expression tag	UNP A0A0E4HLL9
A	-2	HIS	-	expression tag	UNP A0A0E4HLL9
A	-1	LEU	-	expression tag	UNP A0A0E4HLL9
A	0	GLU	-	expression tag	UNP A0A0E4HLL9

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative porin for vanillate trafficking (VanP)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 6 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	150.17Å 150.17Å 105.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	55.32 – 2.90 48.77 – 2.90	Depositor EDS
% Data completeness (in resolution range)	97.4 (55.32-2.90) 97.5 (48.77-2.90)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.08 (at 2.91Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, $R_{free}$	0.237 , 0.284 0.245 , 0.284	Depositor DCC
$R_{free}$ test set	746 reflections (5.01%)	DCC
Wilson B-factor (Å <sup>2</sup> )	63.4	Xtriage
Anisotropy	0.450	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 39.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3170	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.99	8/3250 (0.2%)	1.10	10/4400 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	238	ASP	CB-CG	9.41	1.71	1.51
1	A	330	ASN	CB-CG	6.98	1.67	1.51
1	A	238	ASP	N-CA	6.47	1.59	1.46
1	A	328	GLY	N-CA	5.49	1.54	1.46
1	A	290	ASP	CB-CG	5.41	1.63	1.51
1	A	92	GLU	CB-CG	5.38	1.62	1.52
1	A	92	GLU	CA-CB	5.16	1.65	1.53
1	A	238	ASP	CA-CB	5.13	1.65	1.53

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	ASP	CB-CG-OD1	8.16	125.64	118.30
1	A	238	ASP	OD1-CG-OD2	-6.71	110.56	123.30
1	A	20	ARG	NE-CZ-NH1	6.70	123.65	120.30
1	A	367	ARG	NE-CZ-NH2	-6.68	116.96	120.30
1	A	157	ASP	CB-CG-OD1	6.04	123.73	118.30
1	A	236	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	A	19	ASP	CB-CG-OD1	5.59	123.33	118.30
1	A	238	ASP	CB-CG-OD2	5.57	123.31	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	107	ARG	NE-CZ-NH2	-5.42	117.59	120.30
1	A	92	GLU	OE1-CD-OE2	-5.13	117.15	123.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	289	ILE	Peptide
1	A	354	VAL	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3170	0	3027	29	1
All	All	3170	0	3027	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:TRP:HB2	1:A:293:PRO:CD	2.29	0.63
1:A:18:PHE:CZ	1:A:20:ARG:HD2	2.35	0.62
1:A:120:ALA:HB1	1:A:128:GLN:O	2.01	0.60
1:A:64:LYS:CD	1:A:67:ALA:HB3	2.31	0.60
1:A:92:GLU:OE2	1:A:158:SER:O	2.20	0.59
1:A:330:ASN:CG	1:A:340:ASN:OD1	2.42	0.57
1:A:64:LYS:HD2	1:A:67:ALA:HB3	1.91	0.53
1:A:265:ILE:HB	1:A:307:ARG:HB3	1.90	0.52
1:A:202:ASP:OD2	1:A:238:ASP:CG	2.48	0.52
1:A:292:TRP:HB2	1:A:293:PRO:HD2	1.93	0.51
1:A:292:TRP:CB	1:A:293:PRO:CD	2.89	0.50
1:A:178:THR:HB	1:A:206:LEU:HD22	1.94	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:LEU:HD13	1:A:291:THR:OG1	2.14	0.48
1:A:75:GLY:O	1:A:157:ASP:HB3	2.14	0.47
1:A:173:PHE:CE1	1:A:246:VAL:HG22	2.50	0.47
1:A:202:ASP:OD2	1:A:238:ASP:OD1	2.33	0.46
1:A:92:GLU:OE2	1:A:155:HIS:O	2.34	0.46
1:A:172:ARG:HD3	1:A:276:ALA:CB	2.46	0.45
1:A:83:THR:O	1:A:84:ASN:HB2	2.18	0.44
1:A:115:ASN:HB2	1:A:116:PRO:CD	2.48	0.44
1:A:109:GLY:O	1:A:131:ARG:HA	2.17	0.44
1:A:121:SER:OG	1:A:284:GLU:OE1	2.30	0.43
1:A:261:ASN:HB2	1:A:310:TYR:CE1	2.54	0.43
1:A:292:TRP:CH2	1:A:367:ARG:HG2	2.54	0.43
1:A:64:LYS:HD3	1:A:67:ALA:HB3	2.00	0.43
1:A:2:SER:O	1:A:3:PHE:C	2.57	0.42
1:A:92:GLU:OE2	1:A:154:ASN:OD1	2.37	0.41
1:A:358:TRP:CZ3	1:A:359:LEU:HD21	2.56	0.40
1:A:92:GLU:CD	1:A:154:ASN:OD1	2.59	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:85:GLU:OE2	1:A:330:ASN:ND2[6_555]	2.01	0.19

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	391/413 (95%)	357 (91%)	30 (8%)	4 (1%)	18 51

All (4) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	292	TRP
1	A	239	GLY
1	A	242	LYS
1	A	170	ASN

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/348 (95%)	303 (91%)	29 (9%)	12	34

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	GLU
1	A	4	ILE
1	A	13	THR
1	A	21	ASP
1	A	22	TYR
1	A	28	TYR
1	A	55	LEU
1	A	60	THR
1	A	128	GLN
1	A	131	ARG
1	A	138	LYS
1	A	139	ASP
1	A	159	THR
1	A	169	VAL
1	A	195	LYS
1	A	202	ASP
1	A	206	LEU
1	A	208	ASN
1	A	210	THR
1	A	220	ILE
1	A	235	SER
1	A	236	ARG
1	A	285	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	292	TRP
1	A	295	GLU
1	A	340	ASN
1	A	344	ARG
1	A	355	GLN
1	A	390	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	395/413 (95%)	-0.30	3 (0%) 86 85	41, 59, 86, 109	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	TYR	3.6
1	A	28	TYR	2.1
1	A	188	TYR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.