



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2017 – 08:18 am GMT

PDB ID : 5DLG  
Title : Crystal Structure of Human DNA Polymerase Eta Inserting dGMPNPP Opposite O4-Methylhydine  
Authors : Patra, A.; OFlaherty, D.K.; Egli, M.  
Deposited on : 2015-09-04  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.2 (RC1), CSD as538be (2017)  
Xtriage (Phenix) : 1.9-1692  
EDS : trunk28620  
Percentile statistics : 20161228.v01 (using entries in the PDB archive December 28th 2016)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	1522 (2.38-2.34)
Clashscore	112137	1626 (2.38-2.34)
Ramachandran outliers	110173	1605 (2.38-2.34)
Sidechain outliers	110143	1606 (2.38-2.34)
RSRZ outliers	101464	1528 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	<div> <div>6%</div> <div> <div></div> <div>83%</div> <div>15%</div> <div>..</div> </div> </div>
2	P	8	<div> <div>100%</div> </div>
3	T	12	<div> <div>8%</div> <div> <div></div> <div>58%</div> <div>33%</div> <div>8%</div> </div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 4014 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	430	Total	C	N	O	S	0	0	0
			3366	2110	606	627	23			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q9Y253
A	-1	PRO	-	expression tag	UNP Q9Y253
A	0	HIS	-	expression tag	UNP Q9Y253

- Molecule 2 is a DNA chain called DNA (5'-D(\*AP\*GP\*CP\*GP\*TP\*CP\*AP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	8	Total	C	N	O	P	0	0	0
			161	78	30	46	7			

- Molecule 3 is a DNA chain called DNA (5'-D(\*CP\*AP\*TP\*(5DB)P\*AP\*TP\*GP\*AP\*CP\*GP\*CP\*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	11	Total	C	N	O	P	0	0	0
			226	109	39	67	11			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is 2'-deoxy-5'-O-[(R)-hydroxy{[(R)-hydroxy(phosphonooxy)phosphoryl]amino}phosphoryl]guanosine (three-letter code: XG4) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	196	Total	O	0	0
			196	196		
7	P	10	Total	O	0	0
			10	10		
7	T	16	Total	O	0	0
			16	16		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase eta



- Molecule 2: DNA (5'-D(\*AP\*GP\*CP\*GP\*TP\*CP\*AP\*T)-3')



There are no outlier residues recorded for this chain.

- Molecule 3: DNA (5'-D(\*CP\*AP\*TP\*(5DB)P\*AP\*TP\*GP\*AP\*CP\*GP\*CP\*T)-3')



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.66Å 98.66Å 82.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.90 – 2.35 49.33 – 2.35	Depositor EDS
% Data completeness (in resolution range)	97.9 (37.90-2.35) 98.0 (49.33-2.35)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	0.13	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.172 , 0.247 0.172 , 0.247	Depositor DCC
$R_{free}$ test set	879 reflections (4.73%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.9	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.046 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4014	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5DB, GOL, XG4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.43	0/3425	0.56	1/4623 (0.0%)
2	P	0.82	0/180	0.91	0/276
3	T	0.87	0/228	0.98	0/347
All	All	0.49	0/3833	0.62	1/5246 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	115	ASP	CB-CG-OD1	7.10	124.69	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3366	0	3405	42	1
2	P	161	0	92	0	0
3	T	226	0	114	3	0
4	A	6	0	8	1	0
5	A	2	0	0	0	0
6	A	31	0	13	2	0
7	A	196	0	0	6	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	P	10	0	0	0	0
7	T	16	0	0	0	0
All	All	4014	0	3632	43	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:160:THR:HG22	1:A:162:GLN:H	1.44	0.81
1:A:369:SER:HB2	1:A:423:PHE:HB3	1.75	0.67
1:A:292:GLU:OE2	7:A:601:HOH:O	2.12	0.67
1:A:389:ARG:NH2	7:A:605:HOH:O	2.28	0.63
1:A:322:SER:HB3	1:A:423:PHE:CD1	2.37	0.60
1:A:375:ASP:OD2	1:A:377:ARG:NH1	2.36	0.59
1:A:269:ILE:HG22	1:A:270:LEU:HD23	1.84	0.58
1:A:323:LYS:NZ	7:A:602:HOH:O	2.19	0.57
1:A:132:LEU:O	1:A:134:GLY:N	2.39	0.54
1:A:160:THR:HG21	1:A:165:GLY:HA3	1.91	0.52
1:A:6:ASP:OD1	1:A:6:ASP:N	2.44	0.51
1:A:369:SER:OG	1:A:382:ARG:HG3	2.12	0.50
1:A:405:ASN:ND2	7:A:611:HOH:O	2.43	0.49
1:A:84:ARG:NH1	1:A:350:GLU:OE1	2.35	0.48
1:A:179:GLN:NE2	7:A:603:HOH:O	2.47	0.48
1:A:354:LYS:HE3	1:A:358:ASP:OD2	2.14	0.48
1:A:40:LYS:HD2	1:A:40:LYS:HA	1.52	0.47
1:A:334:ARG:HD3	1:A:413:THR:HA	1.96	0.47
1:A:404:LYS:HA	1:A:415:TRP:HH2	1.80	0.46
1:A:317:LYS:NZ	3:T:9:DC:OP2	2.36	0.46
1:A:321:CYS:O	1:A:423:PHE:HD1	1.99	0.46
1:A:256:ARG:HD2	7:A:710:HOH:O	2.14	0.46
1:A:62:SER:OG	1:A:62:SER:O	2.30	0.46
1:A:352:LEU:HB3	1:A:390:TYR:CE2	2.52	0.45
1:A:334:ARG:HG2	1:A:335:GLU:N	2.32	0.45
1:A:16:CYS:HA	6:A:504:XG4:O2B	2.16	0.44
1:A:14:MET:HG2	1:A:115:ASP:O	2.16	0.44
1:A:324:ASN:OD1	1:A:421:MET:HG3	2.17	0.44
1:A:51:SER:O	1:A:55:ARG:HG3	2.18	0.44
1:A:325:PHE:CD1	1:A:331:LEU:HD21	2.54	0.43
1:A:128:ARG:HH12	1:A:199:GLU:CD	2.21	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:96:SER:O	1:A:100:MET:HG2	2.19	0.43
3:T:5:DA:H2''	3:T:6:DT:O5'	2.19	0.43
1:A:61:ARG:NE	6:A:504:XG4:N7	2.68	0.42
1:A:193:GLY:O	1:A:197:VAL:HG23	2.20	0.42
1:A:290:PHE:O	1:A:294:ASN:HB2	2.19	0.42
1:A:372:VAL:HG13	1:A:417:PRO:HB2	2.01	0.42
1:A:230:ASN:O	1:A:234:ARG:HB2	2.20	0.41
1:A:320:GLY:N	3:T:7:DG:OP2	2.54	0.41
1:A:404:LYS:HA	1:A:415:TRP:CH2	2.55	0.41
1:A:115:ASP:C	1:A:115:ASP:OD1	2.58	0.41
1:A:276:GLY:HA3	4:A:501:GOL:O2	2.21	0.41
1:A:371:ARG:HA	1:A:380:SER:HB3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:LYS:NZ	1:A:432:SER:OXT[4_545]	2.16	0.04

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/435 (98%)	411 (96%)	12 (3%)	3 (1%)	25	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	133	GLN
1	A	134	GLY
1	A	327	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	368/372 (99%)	359 (98%)	9 (2%)	54 67

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	40	LYS
1	A	115	ASP
1	A	138	SER
1	A	152	GLN
1	A	163	LYS
1	A	306	GLU
1	A	321	CYS
1	A	409	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	5DB	T	4	3	14,22,23	0.93	1 (7%)	20,31,34	2.23	5 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5DB	T	4	3	-	0/5/23/24	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	T	4	5DB	C4-N3	2.33	1.35	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	4	5DB	C5-C4-N3	-5.12	119.59	125.24
3	T	4	5DB	C5-C6-N1	-2.34	119.61	122.15
3	T	4	5DB	C1-O4-C4	3.24	120.55	117.30
3	T	4	5DB	O4-C4-C5	3.37	120.77	115.24
3	T	4	5DB	C2-N3-C4	6.02	120.43	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GOL	A	501	-	5,5,5	0.38	0	5,5,5	0.24	0
6	XG4	A	504	5	29,33,33	1.82	5 (17%)	34,52,52	2.02	9 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	501	-	-	0/4/4/4	0/0/0/0
6	XG4	A	504	5	-	0/14/34/34	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	504	XG4	PA-O1A	2.03	1.48	1.46
6	A	504	XG4	C5-C4	2.94	1.47	1.40
6	A	504	XG4	C6-C5	4.08	1.49	1.41
6	A	504	XG4	PA-N3A	4.79	1.76	1.63
6	A	504	XG4	PB-N3A	4.83	1.76	1.63

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	504	XG4	C6-C5-C4	-4.95	115.93	120.84
6	A	504	XG4	N3-C2-N1	-3.95	121.70	127.46
6	A	504	XG4	C5-C6-N1	-3.37	118.69	123.48
6	A	504	XG4	C4-C5-N7	-3.09	106.42	109.41
6	A	504	XG4	PG-O3B-PB	-2.50	123.57	132.38
6	A	504	XG4	C2'-C1'-N9	-2.15	109.16	114.23
6	A	504	XG4	O4'-C1'-N9	2.91	112.68	107.78
6	A	504	XG4	C6-N1-C2	4.62	122.70	116.06
6	A	504	XG4	C2-N3-C4	5.18	121.21	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	501	GOL	1	0
6	A	504	XG4	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	430/435 (98%)	-0.08	24 (5%) 25 36	20, 31, 60, 80	0
2	P	8/8 (100%)	-0.51	0 100 100	32, 37, 40, 41	0
3	T	10/12 (83%)	-0.15	1 (10%) 8 12	34, 36, 54, 71	0
All	All	448/455 (98%)	-0.09	25 (5%) 25 36	20, 31, 60, 80	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	133	GLN	5.5
1	A	378	LEU	5.2
1	A	181	ASP	4.6
1	A	154	PRO	4.1
3	T	2	DA	3.8
1	A	412	GLN	3.7
1	A	328	LYS	3.5
1	A	182	ASN	3.4
1	A	377	ARG	3.4
1	A	372	VAL	3.3
1	A	134	GLY	3.2
1	A	327	GLY	3.2
1	A	132	LEU	3.2
1	A	376	LYS	3.0
1	A	180	ILE	3.0
1	A	130	GLN	2.8
1	A	417	PRO	2.5
1	A	410	GLY	2.5
1	A	1	MET	2.4
1	A	183	LEU	2.3
1	A	408	THR	2.2
1	A	179	GLN	2.2
1	A	42	TRP	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	256	ARG	2.1
1	A	153	GLY	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	5DB	T	4	21/22	0.93	0.15	-	31,43,47,49	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	XG4	A	504	31/31	0.97	0.11	-0.42	20,26,34,40	0
4	GOL	A	501	6/6	0.97	0.08	-1.56	24,26,28,29	0
5	MG	A	502	1/1	0.97	0.07	-1.76	24,24,24,24	0
5	MG	A	503	1/1	0.92	0.07	-1.77	33,33,33,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.