



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 23, 2018 – 11:36 PM EST

PDB ID : 1DML  
Title : CRYSTAL STRUCTURE OF HERPES SIMPLEX UL42 BOUND TO THE C-TERMINUS OF HSV POL  
Authors : Zuccola, H.J.; Filman, D.J.; Coen, D.M.; Hogle, J.M.  
Deposited on : 1999-12-14  
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20030736
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20030736


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2259 (2.70-2.70)
Clashscore	112137	2590 (2.70-2.70)
Ramachandran outliers	110173	2550 (2.70-2.70)
Sidechain outliers	110143	2550 (2.70-2.70)
RSRZ outliers	101464	2275 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	<div> <div>3%</div> <div>50%</div> <div>29%</div> <div>• •</div> <div>16%</div> </div>
1	C	319	<div> <div>4%</div> <div>51%</div> <div>29%</div> <div>5%</div> <div>15%</div> </div>
1	E	319	<div> <div>3%</div> <div>52%</div> <div>30%</div> <div>•</div> <div>14%</div> </div>
1	G	319	<div> <div>6%</div> <div>51%</div> <div>31%</div> <div>•</div> <div>14%</div> </div>
2	B	36	<div> <div>3%</div> <div>56%</div> <div>39%</div> <div>6%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	36	<div><div></div><div>3%</div><div>53%</div><div>42%</div><div>6%</div></div>
2	F	36	<div><div></div><div>3%</div><div>44%</div><div>47%</div><div>8%</div></div>
2	H	36	<div><div></div><div>8%</div><div>42%</div><div>50%</div><div>8%</div></div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9414 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA POLYMERASE PROCESSIVITY FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2059	1309	362	381	7			
1	C	272	Total	C	N	O	S	0	0	0
			2087	1325	368	387	7			
1	E	275	Total	C	N	O	S	0	0	0
			2108	1336	370	395	7			
1	G	275	Total	C	N	O	S	0	0	0
			2108	1336	370	395	7			


- Molecule 2 is a protein called DNA POLYMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	36	Total	C	N	O	S	0	0	0
			263	159	53	50	1			
2	D	36	Total	C	N	O	S	0	0	0
			263	159	53	50	1			
2	F	36	Total	C	N	O	S	0	0	0
			263	159	53	50	1			
2	H	36	Total	C	N	O	S	0	0	0
			263	159	53	50	1			


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR




#### • Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR




#### • Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR







• Molecule 1: DNA POLYMERASE PROCESSIVITY FACTOR




• Molecule 2: DNA POLYMERASE




• Molecule 2: DNA POLYMERASE



• Molecule 2: DNA POLYMERASE



• Molecule 2: DNA POLYMERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.32Å 100.07Å 129.53Å 90.00° 100.62° 90.00°	Depositor
Resolution (Å)	12.00 – 2.70 29.87 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.6 (12.00-2.70) 99.9 (29.87-2.50)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.21 (at 2.51Å)	Xtriage
Refinement program	CNS 0.5	Depositor
R, $R_{free}$	0.230 , 0.281 0.190 , 0.224	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	30.9	Xtriage
Anisotropy	0.174	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 44.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9414	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 56.37 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.7625e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/2097	0.71	2/2851 (0.1%)
1	C	0.41	0/2125	0.70	2/2889 (0.1%)
1	E	0.39	0/2146	0.65	0/2918
1	G	0.38	0/2146	0.64	0/2918
2	B	0.41	0/265	0.56	0/354
2	D	0.40	0/265	0.56	0/354
2	F	0.38	0/265	0.52	0/354
2	H	0.39	0/265	0.55	0/354
All	All	0.40	0/9574	0.66	4/12992 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	155	GLU	N-CA-C	-6.85	92.49	111.00
1	C	155	GLU	N-CA-C	-6.85	92.50	111.00
1	A	136	LEU	CA-CB-CG	5.30	127.49	115.30
1	C	136	LEU	CA-CB-CG	5.27	127.43	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2059	0	2104	100	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2087	0	2130	96	0
1	E	2108	0	2150	104	0
1	G	2108	0	2150	104	0
2	B	263	0	256	18	0
2	D	263	0	256	20	0
2	F	263	0	256	25	0
2	H	263	0	256	22	0
All	All	9414	0	9558	444	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 444 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:121:ARG:HH11	1:G:143:THR:HG22	1.18	1.06
1:A:267:VAL:HG11	1:A:315:LEU:HD13	1.34	1.05
1:G:171:GLN:HA	2:H:1233:THR:HG21	1.39	1.04
1:C:149:ALA:HB3	1:C:152:LEU:HD22	1.41	0.99
1:E:286:GLY:HA2	1:E:304:THR:HG23	1.44	0.99

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	263/319 (82%)	237 (90%)	18 (7%)	8 (3%)	5	12
1	C	268/319 (84%)	240 (90%)	16 (6%)	12 (4%)	3	6
1	E	271/319 (85%)	243 (90%)	21 (8%)	7 (3%)	6	15
1	G	271/319 (85%)	238 (88%)	28 (10%)	5 (2%)	10	25

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	34/36 (94%)	33 (97%)	0	1 (3%)	5	13
2	D	34/36 (94%)	33 (97%)	0	1 (3%)	5	13
2	F	34/36 (94%)	31 (91%)	2 (6%)	1 (3%)	5	13
2	H	34/36 (94%)	30 (88%)	3 (9%)	1 (3%)	5	13
All	All	1209/1420 (85%)	1085 (90%)	88 (7%)	36 (3%)	5	12

5 of 36 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	152	LEU
1	A	223	ALA
1	A	270	ASP
1	A	272	CYS
2	B	1216	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/266 (86%)	214 (94%)	14 (6%)	22	47
1	C	229/266 (86%)	216 (94%)	13 (6%)	24	51
1	E	235/266 (88%)	224 (95%)	11 (5%)	30	60
1	G	235/266 (88%)	222 (94%)	13 (6%)	25	52
2	B	22/22 (100%)	20 (91%)	2 (9%)	11	25
2	D	22/22 (100%)	20 (91%)	2 (9%)	11	25
2	F	22/22 (100%)	19 (86%)	3 (14%)	4	10
2	H	22/22 (100%)	19 (86%)	3 (14%)	4	10
All	All	1015/1152 (88%)	954 (94%)	61 (6%)	22	48

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	204	LEU
1	E	71	THR
1	G	287	THR
1	C	263	ARG
2	D	1234	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	70	ASN
1	E	207	ASN
1	G	171	GLN
1	E	36	GLN
1	G	207	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	267/319 (83%)	-0.19	11 (4%) 38 36	5, 19, 64, 85	0
1	C	272/319 (85%)	-0.08	13 (4%) 31 29	5, 23, 67, 87	0
1	E	275/319 (86%)	-0.02	10 (3%) 43 42	8, 29, 69, 90	0
1	G	275/319 (86%)	0.15	19 (6%) 18 16	8, 33, 76, 91	0
2	B	36/36 (100%)	-0.28	1 (2%) 53 54	9, 23, 45, 69	0
2	D	36/36 (100%)	-0.15	1 (2%) 53 54	11, 23, 48, 71	0
2	F	36/36 (100%)	0.28	1 (2%) 53 54	11, 46, 68, 82	0
2	H	36/36 (100%)	0.43	3 (8%) 12 9	15, 48, 71, 86	0
All	All	1233/1420 (86%)	-0.02	59 (4%) 31 29	5, 27, 69, 91	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	144	THR	6.8
1	E	146	ASP	6.2
1	G	146	ASP	6.1
1	C	152	LEU	5.9
2	H	1200	ASP	5.7

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.