



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 14, 2017 – 04:58 pm GMT

PDB ID : 4DM2  
Title : Contribution of disulfide bond toward thermostability in hyperthermostable endocellulase  
Authors : Kim, H.-W.; Ishikawa, K.  
Deposited on : 2012-02-06  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.2 (RC1), CSD as538be (2017)
Xtriage (Phenix)	:	1.9-1692
EDS	:	trunk28620
Percentile statistics	:	20161228.v01 (using entries in the PDB archive December 28th 2016)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	recalc28949


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	100719	2004 (1.96-1.96)
Clashscore	112137	2136 (1.96-1.96)
Ramachandran outliers	110173	2117 (1.96-1.96)
Sidechain outliers	110143	2117 (1.96-1.96)
RSRZ outliers	101464	2018 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	377	<div> <div>3%</div> <div> <div></div> <div>89%</div> <div>10%</div> </div> <div>•</div> </div>
1	B	377	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>12%</div> </div> <div>•</div> </div>
1	C	377	<div> <div>2%</div> <div> <div></div> <div>85%</div> <div>13%</div> </div> <div>•</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	B	501	-	-	-	X
2	GOL	B	502	-	-	X	X
2	GOL	B	503	-	-	X	X
2	GOL	B	505	-	-	X	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 458aa long hypothetical endo-1,4-beta-glucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	377	Total	C	N	O	S	0	0	0
			3067	2000	500	556	11			
1	B	377	Total	C	N	O	S	0	0	0
			3067	2000	500	556	11			
1	C	377	Total	C	N	O	S	0	0	0
			3067	2000	500	556	11			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	CYS	PRO	CONFLICT	UNP O58925
A	289	LYS	ARG	CONFLICT	UNP O58925
B	74	CYS	PRO	CONFLICT	UNP O58925
B	289	LYS	ARG	CONFLICT	UNP O58925
C	74	CYS	PRO	CONFLICT	UNP O58925
C	289	LYS	ARG	CONFLICT	UNP O58925

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	116	Total	O	0	0
			116	116		
3	B	111	Total	O	0	0
			111	111		
3	C	106	Total	O	0	0
			106	106		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 458aa long hypothetical endo-1,4-beta-glucanase



- Molecule 1: 458aa long hypothetical endo-1,4-beta-glucanase



- Molecule 1: 458aa long hypothetical endo-1,4-beta-glucanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	160.24Å 58.67Å 138.54Å 90.00° 108.93° 90.00°	Depositor
Resolution (Å)	34.53 – 1.95 34.54 – 1.95	Depositor EDS
% Data completeness (in resolution range)	98.9 (34.53-1.95) 98.9 (34.54-1.95)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.39 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.166 , 0.207 0.166 , 0.207	Depositor DCC
$R_{free}$ test set	4450 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.4	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 48.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9570	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	1.24	7/3177 (0.2%)	1.01	7/4336 (0.2%)
1	B	1.21	7/3177 (0.2%)	1.08	12/4336 (0.3%)
1	C	1.13	9/3177 (0.3%)	0.97	9/4336 (0.2%)
All	All	1.20	23/9531 (0.2%)	1.02	28/13008 (0.2%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	108	GLU	CD-OE2	9.08	1.35	1.25
1	A	365	TRP	CD2-CE2	6.81	1.49	1.41
1	B	86	TRP	CD2-CE2	6.51	1.49	1.41
1	A	229	TRP	CD2-CE2	6.13	1.48	1.41
1	B	234	GLU	CD-OE1	6.12	1.32	1.25
1	C	394	TRP	CD2-CE2	5.90	1.48	1.41
1	C	191	TRP	CD2-CE2	5.89	1.48	1.41
1	A	220	TRP	CD2-CE2	5.88	1.48	1.41
1	C	82	TRP	CD2-CE2	5.76	1.48	1.41
1	B	273	TRP	CD2-CE2	5.61	1.48	1.41
1	C	86	TRP	CD2-CE2	5.52	1.48	1.41
1	A	108	GLU	CD-OE1	5.39	1.31	1.25
1	A	86	TRP	CD2-CE2	5.36	1.47	1.41
1	A	272	TRP	CD2-CE2	5.30	1.47	1.41
1	C	323	TRP	CD2-CE2	5.29	1.47	1.41
1	C	166	TRP	CD2-CE2	5.28	1.47	1.41
1	C	180	TRP	CD2-CE2	5.23	1.47	1.41
1	B	220	TRP	CD2-CE2	5.15	1.47	1.41
1	B	379	TRP	CD2-CE2	5.12	1.47	1.41
1	C	220	TRP	CD2-CE2	5.05	1.47	1.41
1	C	267	TRP	CD2-CE2	5.02	1.47	1.41
1	B	365	TRP	CD2-CE2	5.01	1.47	1.41
1	B	368	GLU	CD-OE1	-5.00	1.20	1.25



All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	44	ARG	NE-CZ-NH1	-14.18	113.21	120.30
1	B	44	ARG	NE-CZ-NH2	9.38	124.99	120.30
1	A	331	LYS	CD-CE-NZ	8.63	131.55	111.70
1	C	400	ASP	CB-CG-OD1	7.80	125.32	118.30
1	B	156	ARG	NE-CZ-NH1	-7.40	116.60	120.30
1	B	133	LEU	CB-CG-CD1	-7.24	98.69	111.00
1	A	400	ASP	CB-CG-OD1	7.05	124.65	118.30
1	B	393	ASP	CB-CG-OD2	6.49	124.14	118.30
1	B	400	ASP	CB-CG-OD1	6.46	124.11	118.30
1	B	364	ASP	CB-CG-OD1	6.36	124.02	118.30
1	B	357	ILE	CG1-CB-CG2	-6.26	97.62	111.40
1	A	156	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	C	393	ASP	CB-CG-OD2	5.86	123.57	118.30
1	C	128	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	A	46	ASP	CB-CG-OD1	5.74	123.47	118.30
1	B	131	ASP	CB-CG-OD1	5.73	123.45	118.30
1	C	145	ASP	CB-CG-OD2	-5.60	113.26	118.30
1	B	361	LYS	CD-CE-NZ	-5.59	98.84	111.70
1	A	46	ASP	CB-CG-OD2	-5.54	113.31	118.30
1	C	382	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	352	ASP	CB-CG-OD1	5.41	123.17	118.30
1	A	393	ASP	CB-CG-OD2	5.37	123.13	118.30
1	C	373	ASP	CB-CG-OD1	5.33	123.10	118.30
1	C	186	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	B	186	ARG	NE-CZ-NH2	-5.21	117.70	120.30
1	C	186	ARG	NE-CZ-NH2	-5.14	117.73	120.30
1	B	299	TYR	CB-CG-CD1	-5.07	117.96	121.00
1	C	145	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3067	0	2906	23	0
1	B	3067	0	2906	32	1
1	C	3067	0	2906	38	0
2	A	6	0	8	0	0
2	B	30	0	37	11	0
3	A	116	0	0	2	2
3	B	111	0	0	1	2
3	C	106	0	0	9	0
All	All	9570	0	8763	94	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:266:LYS:H	1:C:266:LYS:CD	1.65	1.09
1:C:128:ARG:HD3	3:C:521:HOH:O	1.51	1.07
1:C:52:ASN:HB2	3:C:501:HOH:O	1.59	1.01
1:C:266:LYS:HD3	1:C:266:LYS:N	1.71	1.01
1:C:266:LYS:HD3	1:C:266:LYS:H	0.86	1.00
1:B:385:ASP:OD2	2:B:503:GOL:H12	1.67	0.95
1:A:67:ASN:HD21	1:A:102:ARG:HH11	1.24	0.86
1:C:67:ASN:HD21	1:C:102:ARG:HH11	1.19	0.85
1:C:128:ARG:CD	3:C:521:HOH:O	2.20	0.79
1:B:238:LYS:O	1:B:242:LYS:HD2	1.83	0.78
1:B:67:ASN:HD21	1:B:102:ARG:HH11	1.31	0.78
1:C:270:ASN:HD22	1:C:306:GLN:HE22	1.34	0.75
1:B:102:ARG:HH22	1:B:200:ASN:HD22	1.36	0.74
1:B:44:ARG:NH1	1:B:58:GLU:OE1	2.22	0.72
1:B:346:LYS:NZ	2:B:502:GOL:H11	2.04	0.71
1:C:67:ASN:ND2	1:C:102:ARG:HH11	1.90	0.69
1:A:102:ARG:HH22	1:A:200:ASN:HD22	1.38	0.69
1:A:67:ASN:ND2	1:A:102:ARG:HH11	1.89	0.69
1:B:331:LYS:HE3	1:B:370:LYS:O	1.92	0.69
1:A:160:THR:OG1	1:A:161:HIS:HD2	1.78	0.66
1:B:86:TRP:H	1:B:124:ASN:HD21	1.44	0.65
1:C:102:ARG:HH22	1:C:200:ASN:HD22	1.46	0.63
1:B:346:LYS:HZ2	2:B:502:GOL:H11	1.64	0.61
1:C:266:LYS:N	1:C:266:LYS:CD	2.39	0.61
1:C:406:LYS:HA	1:C:409:MET:CE	2.31	0.60
1:B:346:LYS:HZ2	2:B:502:GOL:C1	2.14	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:67:ASN:ND2	1:B:102:ARG:HH11	1.98	0.60
1:C:52:ASN:CB	3:C:501:HOH:O	2.31	0.60
1:C:86:TRP:HD1	1:C:124:ASN:HD22	1.50	0.59
1:C:270:ASN:HD22	1:C:306:GLN:NE2	1.99	0.59
1:A:102:ARG:HH22	1:A:200:ASN:ND2	2.00	0.58
1:B:86:TRP:HD1	1:B:124:ASN:HD22	1.50	0.58
1:A:86:TRP:H	1:A:124:ASN:HD21	1.51	0.58
1:A:49:TYR:OH	1:C:54:THR:HG21	2.02	0.58
1:B:86:TRP:H	1:B:124:ASN:ND2	2.01	0.57
1:B:102:ARG:HH22	1:B:200:ASN:ND2	1.99	0.57
1:C:51:ILE:HG12	1:C:58:GLU:HG3	1.85	0.57
1:B:384:GLY:CA	2:B:505:GOL:H31	2.36	0.55
1:C:86:TRP:H	1:C:124:ASN:HD21	1.52	0.55
1:A:67:ASN:HD21	1:A:102:ARG:HD3	1.72	0.55
2:B:503:GOL:O1	2:B:505:GOL:H11	2.08	0.54
1:C:263:SER:HB3	1:C:269:TYR:OH	2.09	0.52
1:B:67:ASN:HD21	1:B:102:ARG:HD3	1.73	0.52
1:C:123:LYS:HE3	3:C:514:HOH:O	2.11	0.51
1:C:52:ASN:CG	3:C:501:HOH:O	2.48	0.51
1:C:102:ARG:HH22	1:C:200:ASN:ND2	2.07	0.51
1:B:302:ASP:HA	2:B:502:GOL:H31	1.94	0.50
1:B:156:ARG:HB3	1:B:158:GLY:O	2.12	0.50
1:C:123:LYS:CE	3:C:514:HOH:O	2.60	0.49
1:C:62:HIS:HA	1:C:373:ASP:OD1	2.12	0.49
1:C:406:LYS:HA	1:C:409:MET:HE2	1.93	0.49
1:B:86:TRP:N	1:B:124:ASN:HD21	2.09	0.49
1:A:86:TRP:HD1	1:A:124:ASN:HD22	1.60	0.49
2:B:503:GOL:O1	2:B:505:GOL:C1	2.61	0.48
1:A:86:TRP:H	1:A:124:ASN:ND2	2.10	0.48
1:C:406:LYS:HA	1:C:409:MET:HE3	1.95	0.48
1:B:385:ASP:CG	2:B:503:GOL:H12	2.34	0.48
1:C:306:GLN:HE21	1:C:307:PRO:HD2	1.79	0.47
1:A:34:GLN:NE2	3:A:703:HOH:O	2.46	0.47
1:A:253:GLY:O	1:A:274:GLY:HA2	2.15	0.47
1:B:384:GLY:HA3	2:B:505:GOL:H31	1.97	0.47
1:B:367:ILE:HG13	1:B:408:LEU:CD1	2.45	0.46
1:C:220:TRP:CE2	1:C:230:ASN:HB3	2.51	0.46
1:A:220:TRP:CE2	1:A:230:ASN:HB3	2.51	0.46
1:B:265:TYR:CE2	1:B:267:TRP:HB2	2.51	0.46
1:C:76:HIS:HE1	3:C:538:HOH:O	1.98	0.46
1:A:381:PRO:HD2	1:A:394:TRP:CZ2	2.51	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:368:GLU:HG3	3:B:676:HOH:O	2.15	0.46
1:B:346:LYS:HZ1	2:B:502:GOL:H11	1.81	0.46
1:C:253:GLY:O	1:C:274:GLY:HA2	2.16	0.45
1:C:86:TRP:H	1:C:124:ASN:ND2	2.13	0.45
1:B:380:ASN:HB3	1:B:382:ASP:OD1	2.17	0.45
1:A:76:HIS:HE1	1:A:108:GLU:OE1	2.00	0.45
1:A:197:ASP:HA	1:A:250:PHE:HB2	1.98	0.45
1:C:311:PRO:C	1:C:313:LYS:H	2.20	0.44
1:C:89:MET:CG	1:C:379:TRP:HE1	2.30	0.44
1:B:208:PRO:HD3	1:B:256:PHE:CE2	2.52	0.44
1:B:220:TRP:CE2	1:B:230:ASN:HB3	2.53	0.43
1:B:200:ASN:ND2	1:B:297:HIS:HE1	2.17	0.43
1:C:52:ASN:ND2	3:C:501:HOH:O	2.51	0.43
1:A:295:SER:HA	1:A:339:VAL:O	2.19	0.42
1:B:323:TRP:CH2	1:B:362:LEU:HA	2.55	0.42
1:A:331:LYS:HE2	1:A:370:LYS:O	2.20	0.42
1:C:201:GLU:HB3	1:C:273:TRP:HB3	2.01	0.42
1:C:295:SER:HA	1:C:339:VAL:O	2.20	0.42
1:A:155:HIS:HA	1:A:200:ASN:HB3	2.02	0.41
1:C:52:ASN:C	1:C:52:ASN:OD1	2.59	0.41
1:A:160:THR:OG1	1:A:161:HIS:CD2	2.67	0.41
1:B:295:SER:HA	1:B:339:VAL:O	2.22	0.40
1:A:380:ASN:HB3	1:A:382:ASP:OD1	2.21	0.40
1:A:86:TRP:CE2	1:A:87:GLU:HG3	2.56	0.40
1:A:34:GLN:HG3	3:A:601:HOH:O	2.21	0.40
1:B:102:ARG:NH2	1:B:200:ASN:HD22	2.11	0.40
1:C:377:TRP:HA	1:C:378:SER:HA	1.85	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:612:HOH:O	3:A:677:HOH:O[4_556]	1.37	0.83
3:B:627:HOH:O	3:B:676:HOH:O[4_555]	1.71	0.49
3:A:603:HOH:O	3:A:684:HOH:O[4_556]	1.78	0.42
1:B:368:GLU:OE1	3:B:627:HOH:O[4_545]	2.12	0.08

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/377 (100%)	364 (97%)	11 (3%)	0	100	100
1	B	375/377 (100%)	368 (98%)	7 (2%)	0	100	100
1	C	375/377 (100%)	368 (98%)	6 (2%)	1 (0%)	44	33
All	All	1125/1131 (100%)	1100 (98%)	24 (2%)	1 (0%)	55	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	312	ALA

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	324/324 (100%)	317 (98%)	7 (2%)	57	49
1	B	324/324 (100%)	317 (98%)	7 (2%)	57	49
1	C	324/324 (100%)	315 (97%)	9 (3%)	49	37
All	All	972/972 (100%)	949 (98%)	23 (2%)	54	44

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	LYS
1	A	117	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	153	ASP
1	A	154	TYR
1	A	331	LYS
1	A	408	LEU
1	A	410	ASP
1	B	34	GLN
1	B	57	GLU
1	B	108	GLU
1	B	153	ASP
1	B	154	TYR
1	B	242	LYS
1	B	408	LEU
1	C	35	THR
1	C	54	THR
1	C	153	ASP
1	C	154	TYR
1	C	207	SER
1	C	266	LYS
1	C	289	LYS
1	C	320	PRO
1	C	410	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	67	ASN
1	A	76	HIS
1	A	124	ASN
1	A	161	HIS
1	A	200	ASN
1	A	359	GLN
1	B	67	ASN
1	B	76	HIS
1	B	124	ASN
1	B	200	ASN
1	B	286	ASN
1	C	67	ASN
1	C	76	HIS
1	C	124	ASN
1	C	200	ASN
1	C	306	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	359	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	501	-	5,5,5	1.04	0	5,5,5	1.39	0
2	GOL	B	501	-	5,5,5	0.34	0	5,5,5	0.62	0
2	GOL	B	502	-	5,5,5	0.53	0	5,5,5	1.70	2 (40%)
2	GOL	B	503	-	5,5,5	1.03	0	5,5,5	2.58	3 (60%)
2	GOL	B	504	-	5,5,5	0.91	0	5,5,5	0.76	0
2	GOL	B	505	-	5,5,5	1.65	1 (20%)	5,5,5	4.11	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	501	-	-	0/4/4/4	0/0/0/0
2	GOL	B	501	-	-	0/4/4/4	0/0/0/0
2	GOL	B	502	-	-	0/4/4/4	0/0/0/0
2	GOL	B	503	-	-	0/4/4/4	0/0/0/0
2	GOL	B	504	-	-	0/4/4/4	0/0/0/0
2	GOL	B	505	-	-	0/4/4/4	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	505	GOL	C1-C2	-2.45	1.43	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	505	GOL	O1-C1-C2	-8.51	67.21	110.07
2	B	503	GOL	O2-C2-C1	-4.10	89.47	108.84
2	B	503	GOL	O3-C3-C2	-3.48	92.56	110.07
2	B	502	GOL	O1-C1-C2	2.06	120.46	110.07
2	B	503	GOL	C3-C2-C1	2.07	119.74	111.52
2	B	505	GOL	O2-C2-C3	2.57	120.98	108.84
2	B	502	GOL	C3-C2-C1	3.10	123.84	111.52

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	502	GOL	5	0
2	B	503	GOL	4	0
2	B	505	GOL	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	377/377 (100%)	-0.27	12 (3%) 48 59	4, 11, 25, 58	8 (2%)
1	B	377/377 (100%)	-0.27	13 (3%) 46 56	6, 12, 23, 50	10 (2%)
1	C	377/377 (100%)	-0.10	7 (1%) 67 76	6, 15, 29, 40	9 (2%)
All	All	1131/1131 (100%)	-0.21	32 (2%) 53 64	4, 13, 27, 58	27 (2%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	311	PRO	5.3
1	C	136	MET	5.2
1	A	136	MET	4.9
1	A	410	ASP	4.7
1	B	136	MET	4.6
1	A	162	ILE	4.3
1	A	286	ASN	4.3
1	A	234	GLU	3.5
1	A	311	PRO	3.4
1	A	161	HIS	3.3
1	B	234	GLU	3.2
1	B	160	THR	3.1
1	B	161	HIS	3.1
1	B	169	GLU	2.7
1	B	162	ILE	2.6
1	B	286	ASN	2.6
1	A	310	GLY	2.5
1	C	54	THR	2.5
1	C	234	GLU	2.4
1	B	368	GLU	2.4
1	C	170	ASP	2.4
1	B	410	ASP	2.4
1	A	312	ALA	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	312	ALA	2.3
1	C	286	ASN	2.2
1	B	311	PRO	2.2
1	B	170	ASP	2.2
1	A	122	SER	2.1
1	A	160	THR	2.1
1	A	368	GLU	2.1
1	B	310	GLY	2.0
1	C	151	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	505	6/6	0.91	0.20	15.89	12,14,22,34	0
2	GOL	B	502	6/6	0.91	0.22	7.19	19,26,28,36	0
2	GOL	B	503	6/6	0.84	0.19	4.84	17,24,25,26	0
2	GOL	B	501	6/6	0.94	0.11	2.50	19,29,33,36	0
2	GOL	A	501	6/6	0.88	0.14	1.98	21,25,27,29	0
2	GOL	B	504	6/6	0.94	0.10	0.53	19,20,21,25	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.